

DET KGL. NORSKE VIDENSKABERS SELSKAB
MUSEET


GUNNERIA
40



Terje Klokk

MIRE AND FOREST VEGETATION FROM KLÆBU,
CENTRAL NORWAY

TRONDHEIM 1982



MIRE AND FOREST VEGETATION
FROM KLÆBU, CENTRAL NORWAY

by

Terje Klokk

University of Trondheim
The Royal Norwegian Society of Sciences and Letters, the Museum

ISBN 82-7126-308-0

ISSN 0332-8554

ABSTRACT

Klokk, Terje. 1982. Mire and forest vegetation from Klæbu, Central Norway. *Gunneria* 40: 1-71.¹

36 mire communities are subjectively arranged along the vegetation gradients poor - rich, mud bottom - hummock, and partly mire margin - mire expanse. There are 13 from ombrotrophic mire, 10 from poor fen, 4 from intermediate fen, 5 from rich fen, and 4 from extremely rich fen. Seven forest associations are recognized, viz. *Barbilophozio-Pinetum* Br.-Bl. et Sissingh 1939 em. K.-Lund 1967, *Chamaemoro-Piceetum* K.-Lund 1962, *Melico-Piceetum* K.-Lund 1962, *Equiseto palustris-Salicetum* ass. prov., *Alno (incanae)Prunetum* K.-Lund ex. Aune 1973, and *Ulmo-Tilieteum boreale* ass. prov. The *Vaccinium-Cornus suecica-Hylocomium splendens* community (Tab. IX) is also a forest community.

Ordination (RA) of forest analyses mainly confirms the subjective classification. The variation along the first and second axis, in the ordination diagram of forest analyses, is interpreted as a poor - rich gradient and a moisture gradient, respectively.

Exchangeable Ca^{++} and pH (mires only) are positively correlated with a poor - rich gradient.

Terje Klokk*, Botanical Institute, University of Trondheim, Bjørnsons gt. 12, N-7000 Trondheim, Norway.


* Present address, Applied Chemistry Division, Environmental Section, SINTEF, N-7034 TRONDHEIM-NTH, Norway.

¹ Botanical Series 12.




CONTENTS

INVESTIGATION AREA	7
GEOLOGY	8
CLIMATE	9
METHODS AND NOMENCLATURE	11
FLORISTIC ELEMENTS	13
The suboceanic element	14
Other elements	15
MIRE VEGETATION	16
Hydrotopographic complexes	16
Local variations in vegetation	16
SYNOPSIS OF THE MIRE COMMUNITIES	20
OMBROTROPHIC VEGETATION	22
Open bog	23
Hollow vegetation	23
Hummock vegetation	26
Tree-/shrub covered bog	28
POOR FEN	29
Open poor fen	30
Tree-/shrub covered poor fen	33
INTERMEDIATE FEN (OPEN)	34
RICH FEN	37
Open rich fen	38
Tree-/shrub covered rich fen	39
EXTREMELY RICH FEN	40
Open extremely rich fen	40
Tree-/shrub covered extremely rich fen	42
THE FOREST VEGETATION	42
Systematics of the forest communities	42
ASSOCIATION <i>CHAMAEMORO-PICEETUM</i>	43
ASSOCIATION <i>EQUISETO PALUSTRIS-SALICETUM</i>	44
ASSOCIATION <i>BARBILOPHOZIO-PINETUM</i>	45
ASSOCIATION <i>EU-PICEETUM</i>	46
ASSOCIATION <i>MELICO-PICEETUM</i>	48
ASSOCIATION <i>ALNO (INCANAEE)-PRUNETUM</i>	53
ASSOCIATION <i>ULMO-TILIETUM BOREALE</i>	54



GRADIENTS IN THE FOREST VEGETATION	58
SOIL SAMPLES	60
ACKNOWLEDGEMENTS	62
SUMMARY	63
REFERENCES	65



INVESTIGATION AREA

The investigated area includes the northern part of Klæbu rural district, 20 km south of Trondheim (Fig. 1). Most of the area lies between 150-400 m a.s.l., with the highest point 537 m a.s.l. and the lowest 99 m a.s.l. West of the river Nidelva the terrain gradually rises towards Vassfjellet (710 m a.s.l.). The clay soil area closest to the river, is cultivated. Elsewhere spruce forest dominates the landscape, with scattered stands of pine forest on shallow soil and borders of grey alder along the Nidelva and brooks. Pine is more frequent at altitudes of 400-500 m a.s.l., especially in the eastern part of the area, but the stands are small. The two most common mire complexes are excentric ombrotrophic bogs, in the lowest parts (150-200 m a.s.l.), and sloping fens.





Fig. 1. Survey map, investigation area hatched (x = precipitation stations, K = Klæbu, L = Løksmyr, V = Vennafjell).

GEOLOGY

This section is mainly based on Wolff (1968). The investigated area consists of cambro-silurian sedimented rocks, represented by the Støren-group and lower Hovin-group (Fig. 2).



KJÖLHAUGEN GROUP - RÖROS GROUP - UPPER HOVIN GROUP (UPPER ORDOVICIAN)

[diagonal lines] PHYLLITE, METAGRAYWACKES, WITH INCREASING AMOUNTS OF BIOTITE, HORNBLEND AND GARNET TOWARDS THE SOUTHEAST, PARTLY CONGLOM.

[dotted pattern] POLYGENOUS CONGLOMERATE

SULÅMO GROUP - LOWER HOVIN GROUP (MIDDLE ORDOVICIAN)

[horizontal lines] DARK SHALE AND RHYOLITE TUFF IN WEST, GREENSTONE IN EAST

[solid grey] GREY CALCAROUS SANDSTONE AND GREY TO DARK PHYLLITE

[diagonal lines with dots] VENNA, STOKKVOLA, LILLE FUNDSJØ AND SIMILAR CONGLOMERATES

FUNDSJØ GROUP - STØREN GROUP (LOWER ORDOVICIAN)

[wavy lines] GREENSTONES AND QUARTZKERATOPHYRES

[cross-hatch] GRANODIORITIC GNEISS

SONVATN GROUP - GULA SCHIST GROUP (CAMBRIAN)

[wavy lines] MICA SCHISTS, OFTEN WITH GARNET

Fig. 2. Geological map (after Wolff 1976).

The geological formations are younger from west to east. At Byneset the Støren-group dominates, with greenstone as the dominant rock. Towards the east the younger lower Hovin-group predominates. In the border zone between, a conglomerate with pebbles occurs frequently. The Hovin-group consists of fine grained sedimentary rocks such as sandstone and schists. This layer is deposited above the greenstone of the Støren group.

A wedge of greenstone passes lake Jonsvatnet and reaches the outlet of lake Selbusjøen. East of this wedge the rocks of the Hovin-group dominate. The Caledonian foldings caused the present positions of the various layers.

Within the investigated area the highest sea level after the last glacial age was approximately 180 m higher than the present level. At lower levels fine grained deposits, mostly clay dominate the landscape.

A marked moraine (the "Heimdal-morain") passes the area in the direction south-north (Carstens 1919). This moraine is covered by homogenous spruce forest.

CLIMATE

Within the investigated area there are two precipitation stations, Klæbu at $63^{\circ}18'N$ - $10^{\circ}28'E$ and Løksmyr at $63^{\circ}14'N$ - $10^{\circ}27'E$. Both stations are situated at low altitudes, 143 and 170 m a.s.l., respectively. Data from the Vennafjell station $63^{\circ}20'N$ - $10^{\circ}39'E$ (671 m a.s.l.) is included. These three stations together are considered to be representative for the area under investigation.

Neighbouring stations which also measure temperature are Tyholt Trondheim $63^{\circ}25'N$ - $10^{\circ}26'E$ (113 m a.s.l.) and Selbu $63^{\circ}12'N$ - $11^{\circ}07'E$ (197 m a.s.l.). The latter is considered more representative.

The humidity index of Martonné (Dahl 1950) is calculated,

$$H = \frac{\text{Yearly precipitation}}{\text{Yearly mean temperature } ^{\circ}\text{C} + 10}$$

The humidity index of Martonné of the Tyholt-Trondheim station is 57 and that of the Selbu station 58. Precipitation




Fig. 3. Precipitation norms for the period 1931-1960 for the stations Klæbu, Løksmyr and Vennafjell.




Fig. 4. Temperature norms for the period 1931-1960 for the stations Trondheim-Tyholt and Selbu.

values of the stations within the investigated area are higher than those of the Tyholt-Trondheim and Selbu stations. Based on precipitation at the station Løksmyr (Fig. 3) and a mean temperature between Tyholt-Trondheim and Selbu (Fig. 4), the estimated representative humidity index of Klæbu (humidity index of Martonné) is 66. This is a relatively high value considering the distance to the coast (Dahl 1950).

The hygroterm of Amman (Amman 1929)

$$A = \frac{\text{Yearly precipitation cm} \times \text{Yearly mean temperature } {}^{\circ}\text{C}}{\text{Mean temperature of warmest month} - \text{mean temperature of coldest month}}$$

gives values of 25 and 20 for Tyholt-Trondheim and Selbu, respectively, and a value of 26 corresponding to the humidity index of 66 as calculated above. This illustrates the suboceanic climate in the area.

Calculated from temperature figures of Tyholt-Trondheim and Selbu, July is the warmest month (mean 14,6 - 14,7[°]C) and January the coldest one (- 3,1 - - 4,2[°]C). The latter January temperature is probably more representative of the area.

METHODS AND NOMENCLATURE

555 relevés were recorded. The field methods are constant in each main type of vegetation, however, different sizes of relevés are used.

According to Fennoscandian mire tradition small relevés were used, 0.25 m² in open mires and 1 m² in shrub/tree covered mires. Larger relevés in shrub/tree covered mires are used to get a more satisfactory estimate of the degree of cover in shrub and tree layers. Moreover the mosaic is not so fine-meshed as in open mires.

In the forests (including the damp forests) relevés of 4 m² are used. Five relevés are recorded in each stand, both in mire and forest vegetation.

A stand is defined according to Dahl & Hadac (1949). Analysing by small relevés makes it possible to maintain a very strict homogeneity definition (Dahl 1967). Within the stand the relevés are subjectively distributed to encompass the possible variation

(Sjörs 1948).

Cover is noted according the the scale of Hult-Sernander, with some modifications. The main criticism against this scale is that cover degree 5 is too wide (Poore 1955). Cover degree 6 ($> 3/4$ cover) has been used in the present study (see also Steen 1956, Moen 1970, Fransson 1972). Species with cover degree 1, but only 1-2 individuals are noted with +.

Soil samples were analysed for pH (mires only), and exchangeable potassium, calcium, and magnesium. Potassium is calculated by means of flame photometry, and calcium and magnesium by atomicabsorbtion-spectrometry. pH was measured in water pressed from the peat. After precipitation for 12 hours pH was measured in the supernatant.

Similar stands are grouped together in Tables. The forest communities are grouped in a hierachial system, following the system of Kielland-Lund (1962, 1967 a, 1971, 1973). The mire analyses are calssified in communities, and the communities are arranged along the poor - rich and mud bottom - hummock gradients. Open mires are separated from tree/shrub ($> 10\%$ cover of shrub- and/or tree layer) covered mires. Some communities are divided into variants based on dominant relations. The mire communities usually correspond approximately to the association level, but sometimes correspond to a lower level.

Differential species and characteristic species are defined according to Braun-Blanquet (1964). Constant species are species with $> 80\%$ frequency.

Some of the subjective classifications are compared with numerical classifications obtained from the computer programme TABORD (Persson 1977, van der Maarel et al. 1978). The numerical treatment also includes use of the ordination technique reciprocal averaging (RA) (Hill 1973). For evaluation of this technique in relation to other ordination techniques see e.g. Orloci (1978), Whittaker & Gauch (1978), and van der Maarel (1979).

With some exceptions the nomenclature follows Lid (1974; vascular plants), Nyholm (1954-1969; mosses), Arnell (1956; liverworts), and Dahl & Krogh (1973; lichens). Species are not separated within the *Alchemilla vulgaris* group, *Hieracium*, and *Taraxacum*. *Empetrum nigrum* and *E. hermaphroditum* are not separated on sterile

collections. *Drepanocladus intermedius* (Lindb.) Warnst. is regarded as a distinct species (cf. Sjörs 1948, Fransson 1972, Malmer 1973). Species within the *Sphagnum Subsecunda* group are not separated. *Sphagnum fallax* includes *S. angustifolium*, *S. fallax* s.str., and *S. flexuosum*. *Rhytidadelphus calvescens/R. squarrosus* is handled collectively. Species are not separated within the liverwort genera *Calypogeia*, *Cephalozia*, *Cephaloziella*, *Lophozia*, *Pellia*, *Scapania*, and partly *Riccardia*. To facilitate the field work *Cladopodiella fluitans* and *Gymnocolea inflata* were not separated (cf. Flatberg 1970). Concerning '*Leiocolea borealis*' see Frisvoll & Moen (1981).

FLORISTIC ELEMENTS

The investigated area belongs to the northern spruce forest region of the boreale zone (Sjörs 1967).

A characteristic feature of the lowland flora of central parts of Trøndelag is the occurrence of several different floristic elements. The relatively humid climate is reflected in flora and vegetation with a large number of species with a suboceanic distribution.

A southern termophilous element is represented in south and south-west faced slopes with a warm local climate.

Some mountain species occur in Trøndelag at lower altitudes than in southern Norway (Flatberg & Sæther 1974). Some of these grow within the area at levels of 150-300 m a.s.l.

Few species with an eastern Fennoscandian distribution occur in the area.

The distribution of species and the categorising of different floristic elements are based on Dahl (1950), Nyholm (1954-69), Arnell (1956), Fægri (1960), Lye (1968), Størmer (1969), Hultén (1971), Gjærevoll (1972), Lid (1974), Flatberg & Sæther (1974), the authors' experiences during the field work, and data obtained from collections at the Museum of The Royal Norwegian Society of Sciences and Letters, Trondheim (TRH).

The suboceanic element

Species included in this element have a Fennoscandian sub-oceanic distribution and/or have their main distribution along the western coast of Norway (Flatberg 1970). The following species with a suboceanic distribution are found in the investigated area or its closest surroundings (w - weaker suboceanic tendency, * - species that belong to a thermophilous southern coastal element);

<i>Blechnum spicant</i>	* <i>Polystichum braunii</i>
<i>Carex hostiana</i>	<i>Ranunculus flammula</i>
<i>C. pulicaris</i>	<i>Succisa pratensis</i> (w)
<i>C. tumidicarpa</i>	<i>Thelypteris limbosperma</i>
<i>Erica tetralix</i>	<i>Bazzania trilobata</i>
* <i>Festuca altissima</i>	<i>Dicranum leioneuron</i> (w)
* <i>Galium odoratum</i>	<i>Eurhynchium striatum</i>
<i>Juncus bulbosus</i>	<i>Mnium hornum</i>
<i>J. conglomeratus</i>	<i>M. undulatum</i>
<i>J. effusus</i>	<i>Plagiothecium undulatum</i>
<i>Lycopodium inundatum</i>	<i>Racomitrium lanuginosum</i> (w)
<i>Myrica gale</i>	<i>Rhytidiodelphus loreus</i> (w)
<i>Narthecium ossifragum</i>	<i>Sphagnum imbricatum</i>
<i>Plantago lanceolata</i>	<i>S. quinquefarium</i>
	<i>S. strictum</i>

The suboceanic element is a geographic element including species with a different ecology, and there are presumably different ecological factors limiting their distribution. *Rhynchospora alba* and *R. fusca*, both occurring in the area are often regarded as sub-oceanic species, but are considered to be "lowland species" in the present study (see also Flatberg 1970).

Some species which are regarded as indicators of fen vegetation in more continental parts of Fennoscandia occur in oceanic areas in ombrotrophic vegetation (Skogen 1969, Flatberg 1976). *Carex pauciflora*, *Narthecium ossifragum*, *Sphagnum papillosum*, and *S. pulchrum* all grow in ombrotrophic mires in the area. *Racomitrium lanuginosum* is dominant in hummocks and the *Scirpus caespitosus-Sphagnum rubellum-S. tenellum* hollow community are oceanic features in the ombrotrophic vegetation (Skogen 1969, Flatberg 1970).

The mire communities show most resemblance to analogous communities in humid areas in Fennoscandia.

Other elements

Several thermophilous species with a southern distribution in Norway have their northern limit in Trøndelag. At their northern limits they grow on steep south facing slopes and screes with a favourable local climate and eutrophic soil (Gjærevoll 1972). The following species with a southern distribution in Norway grow within the investigated area (w - weaker thermophilous tendency);

<i>Corylus avellana</i>	<i>Hypericum hirsutum</i>
<i>Ulmus glabra</i>	<i>Lactuca muralis</i>
<i>Geum urbanum</i> (w)	<i>Polygonatum odoratum</i>
<i>Hepatica nobilis</i> (w)	<i>Satureja acinos</i>
<i>Humulus lupulus</i> (w)	<i>Viola mirabilis</i>

This group also includes species with varying ecology, probably belonging to several floristic elements.

Most species with an eastern Fennoscandian distribution are not found in the investigation area. However, the following species are exceptions:

<i>Picea abies</i>	<i>Eriophorum gracile</i>
<i>Carex buxbaumii</i>	<i>Ranunculus peltatus</i>
<i>C. chordorrhiza</i>	<i>Scheuchzeria palustris</i>
<i>Aconitum septentrionale</i>	<i>Viola rupestris</i>
	<i>Sphagnum majus</i>

In Trøndelag several mountain species grow at low altitudes, often on north facing rocks and screes (Dahl 1950, Flatberg & Sæther 1974). The following mountain species are found in localities lower than 300 m a.s.l. in the investigation area:

<i>Poa alpina</i>	<i>Oxyria digyna</i>
<i>Alchemilla alpina</i>	<i>Saxifraga cotyledon</i>
<i>Cerastium alpinum</i>	<i>Sedum rosea</i>

MIRE VEGETATION

Hydrotopographic complexes

Moen (1973) lists eight types of complexes for Norway, viz. four ombrotrophic and four minerotrophic. The two dominating complexes within the investigated area are excentric raised bogs and sloping fens. Most of the large excentric raised bogs are situated in the lower regions of the area. The sloping fens are rich, especially in the western parts of the area. In the eastern regions poor fens are common. "Ribbed" minerotrophic complexes occur scattered over the area.

Local variations in vegetation

Fennoscandian mire literature distinguishes between three main local "directions of variation" in vegetation, viz. the poor-rich gradient, the mud bottom - hummock gradient, and the mire expanse - mire margin gradient (Sjörs 1948, 1950, Ruuhijärvi 1960, Malmer 1962 a, Sonesson 1970, and Moen 1973).

The main gradient for classification in the present paper is the poor - rich gradient. The mire expanse - mire margin gradient distinguishes only between tree-/shrub covered mires and open mires.

The distribution of species along the three local gradients on Tabs. I, II, and III is based on the arrangement of the communities in the present paper.

Tab. I. Distribution of species groups along the poor - rich gradient.
 common, ----- scattered.

Bog	Fen			
	Poor	Intermed.	Rich	Extremely rich
<i>Carex hostiana</i>				
<i>C. lepidocarpa</i>				
<i>Schoenus ferrugineus</i>				
<i>Leiocolea rutheana</i>				
<i>Carex buxbaumii</i>				
<i>C. flava</i>				
<i>Eriophorum latifolium</i>				
<i>Thalictrum alpinum</i>				
<i>Calliergon trifarium</i>				
<i>Drepanocladus intermedius</i>				
<i>Sphagnum warnstorffii</i>				
<i>Leiocolea borealis</i>				
<i>Campylium stellatum</i>				
<i>Drepanocladus revolvens</i>				
<i>Scorpidium scorpioides</i>				
<i>Selaginella selaginoides</i>				
<i>Tofieldia pusilla</i>				
<i>Drepanocladus badius</i>				
<i>D. exannulatus</i>				
<i>Sphagnum subnitens</i>				
<i>S. sect. Subsecunda</i>				
<i>Viola palustris</i>				
<i>Carex lasiocarpa</i>				
<i>C. rostrata</i>				
<i>Eriophorum angustifolium</i>				
<i>Molinia caerulea</i>				
<i>Equisetum fluviatile</i>				
<i>Menyanthes trifoliata</i>				
<i>Pinguicula vulgaris</i>				
<i>Potentilla erecta</i>				
<i>Eriophorum vaginatum</i>				
<i>Rubus chamaemorus</i>				
<i>Calliergon stramineum</i>				
<i>Drepanocladus schulzei</i>				
<i>Sphagnum fallax</i>				
<i>S. fuscum</i>				
<i>S. lindbergii</i>				
<i>S. magellanicum</i>				
<i>S. nemoreum</i>				
<i>S. papillosum</i>				
<i>S. tenellum</i>				
<i>Cladopodiella fluit./Gymnoc.infl.</i>				
<i>Scheuchzeria palustris</i>				
<i>Sphagnum cuspidatum</i>				
<i>Andromeda polifolia</i>				
<i>Carex limosa</i>				
<i>Drosera anglica</i>				
<i>D. rotundifolia</i>				
<i>Myrica gale</i>				
<i>Oxycoccus microcarpus</i>				
<i>O. quadripetalus</i>				

Tab. II. Distribution of species groups along the mud bottom -
hummock gradient, _____ common, ----- scattered.

	Mud bottom	Carpet	Lawn	Hummock
<i>Equisetum fluviatile</i>	_____			
<i>Utricularia intermedia</i>	_____			
<i>Scheuchzeria palustris</i>	-----			
<i>Carex limosa</i>	_____			
<i>Juncus stygius</i>	_____			
<i>Rhynchospora alba</i>	_____			
<i>Drepanocladus exannulatus</i>	_____			
<i>D. schulzei</i>	_____			
<i>Sphagnum cuspidatum</i>	_____			
<i>S. lindbergii</i>	_____			
<i>Drosera anglica</i>	-----			
<i>Menyanthes trifoliata</i>	-----			
<i>Scorpidium scorpioides</i>	-----			
<i>Sphagnum majus</i>	-----			
<i>Carex lasiocarpa</i>	_____			
<i>C. rostrata</i>	_____			
<i>Eriophorum angustifolium</i>	_____			
<i>Cladopodiella fluit./Gymnoc.infl.</i>	_____			
<i>Sphagnum pulchrum</i>	_____			
<i>S. riparium</i>	_____			
<i>Carex chordorrhiza</i>	-----			
<i>Sphagnum subnitens</i>	-----			
<i>Scirpus caespitosus</i>	-----			
<i>S. hudsonianus</i>	-----			
<i>Selaginella selaginoides</i>	-----			
<i>Campyrium stellatum</i>	-----			
<i>Drepanocladus intermedius</i>	-----			
<i>Sphagnum papillosum</i>	-----			
<i>S. tenellum</i>	-----			
<i>S. compactum</i>	-----			
<i>Narthecium ossifragum</i>	_____			
<i>Molinia caerulea</i>	_____			
<i>Tofieldia pusilla</i>	_____			
<i>Drepanocladus badius</i>	-----			
<i>Oxycoccus microcarpus</i>	-----			
<i>O. quadripetalus</i>	-----			
<i>Calypogeia sp(p).</i>	-----			
<i>Dicranum leioneuron</i>	_____			
<i>Sphagnum rubellum</i>	_____			
<i>Lepidozia setacea</i>	_____			
<i>Ptilidium ciliare</i>	_____			
<i>Mylia anomala</i>	_____			
<i>Cephalozia sp(p).</i>	_____			
<i>Cephaloziella sp(p).</i>	_____			
<i>Calluna vulgaris</i>	_____			
<i>Empetrum spp.</i>	_____			
<i>Plurozium schreberi</i>	_____			
<i>Polytricum juniperinum var. gracilis</i>	_____			
<i>Sphagnum fuscum</i>	_____			
<i>Racomitrium lanuginosum</i>	_____			
<i>Cladonia alpestris</i>	_____			
<i>C. rangiferina</i>	_____			
<i>C. arbuscula</i>	_____			
<i>Drosera rotundifolia</i>	-----			
<i>Andromeda polifolia</i>	_____			
<i>Eriophorum vaginatum</i>	_____			

Tab. III. Distribution of species groups on open and tree/shrub covered mires. — common, ----- scattered.

	Mire margin	Mire expanse
<i>Alnus incana</i>		
<i>Agrostis canina</i>		
<i>Carex echinata</i>		
<i>C. nigra</i>		
<i>Deschampsia caespitosa</i>		
<i>Anemone nemorosa</i>		
<i>Cornus suecica</i>		
<i>Crepis paludosa</i>		
<i>Filipendula ulmaria</i>		
<i>Galium boreale</i>		
<i>Geranium sylvaticum</i>		
<i>Hieracium spp.</i>		
<i>Melampyrum pratense</i>		
<i>Saussurea alpina</i>		
<i>Calliergonella cuspidata</i>		
<i>Dicranum majus</i>		
<i>D. scoparium</i>		
<i>Hylocomium splendens</i>		
<i>Mnium pseudopunctatum</i>		
<i>M. seligeri</i>		
<i>Sphagnum fallax</i>		
<i>S. russowii</i>		
<i>Chiloscyphus polyanthus</i>		
<i>Obtusifolium obtusum</i>		
<i>Tritomaria quinquedentata</i>		
<i>Picea abies</i>		-----
<i>Pinus sylvestris</i>		-----
<i>Vaccinium myrtillus</i>		-----
<i>V. uliginosum</i>		-----
<i>V. vitis-idaea</i>		-----
<i>Potentilla erecta</i>		-----
<i>Trientalis europaea</i>		-----
<i>Aulacomnium palustre</i>		-----
<i>S. nemoreum</i>		-----
<i>Drosera anglica</i>		-----
<i>D. rotundifolia</i>		-----
<i>S. fuscum</i>		-----
<i>S. subnitens</i>		-----
<i>Carex limosa</i>		
<i>Juncus stygius</i>		
<i>Scirpus caespitosus</i>		
<i>Rhynchospora alba</i>		
<i>Scheuchzeria palustris</i>		
<i>Schoenus ferrugineus</i>		
<i>Utricularia intermedia</i>		
<i>Drepanocladus schulzei</i>		
<i>Racomitrium lanuginosum</i>		
<i>Scorpidium scorpioides</i>		
<i>Sphagnum compactum</i>		
<i>S. cuspidatum</i>		
<i>S. lindbergii</i>		
<i>S. pulchrum</i>		
<i>S. majus</i>		
<i>S. rubellum</i>		
<i>S. tenellum</i>		
<i>Cladopodiella fluit./Gymnoc. infl.</i>		
<i>Andromeda polifolia</i>		
<i>Calluna vulgaris</i>		
<i>Empetrum spp.</i>		
<i>Oxycoccus microcarpus</i>		
<i>O. quadripetalus</i>		
<i>Carex lasiocarpa</i>		
<i>C. pauciflora</i>		
<i>C. rostrata</i>		
<i>Eriophorum angustifolium</i>		
<i>E. latifolium</i>		
<i>E. virginatum</i>		
<i>Menyanthes trifoliata</i>		
<i>Molinia caerulea</i>		
<i>Pinguicula vulgaris</i>		
<i>Pleurozium schreberi</i>		
<i>Sphagnum warnstorfii</i>		
<i>Ptilidium ciliare</i>		
<i>Riccardia pinguis</i>		

SYNOPSIS OF THE MIRE COMMUNITIES

Ombrrophic mire

Open

Hollow

Mud bottom

Scheuchzeria palustris-Sphagnum cuspidatum
community

Carex limosa-Cladopodiella fluitans/Gymnocolea
inflata community

Mud bottom/carpet

Carex limosa-Rhynchospora alba-Drepanocladus
schulzei-Cladopodiella fluitans/Gymnocolea in-
flata community

Carpet

Eriophorum vaginatum-Sphagnum cuspidatum-S.
majus-S. pulchrum community

Sphagnum magellanicum-S. pulchrum community

Lawn

Scirpus caespitosus-Sphagnum rubellum-S. te-
nellum community

Hummock

Calluna vulgaris-Drosera rotundifolia-Rubus
chamaemorus-Sphagnum fuscum community

Calluna vulgaris-Pleurozium schreberi-Polytri-
chum juniperinum var. *gracilis* community

Dicranum leioneuron-Sphagnum nemoreum community

Racomitrium lanuginosum-Cladonia community

Cladonia community

Sphagnum fuscum community

Tree/shrub covered

Vaccinium-Cornus suecica-Hylocomium splendens
community

Minerotrophic mire

Poor fen

Open

Mud bottom

Carex limosa-Menyanthes trifoliata community

Carpet

Eriophorum angustifolium-Sphagnum pulchrum
community

Carex rostrata-Sphagnum riparium community

Drosera rotundifolia-Sphagnum magellanicum
community

Carex limosa-Sphagnum papillosum-S. majus
community

Carpet/lawn	<i>Scirpus caespitosus-Sphagnum compactum</i> - <i>Cladopodiella fluitans/Gymnocolea inflata</i> community
Lawn	<i>Scirpus caespitosus-Sphagnum tenellum</i> community <i>Molinia caerulea-Scirpus caespitosus-Narthecium</i> <i>ossifragum</i> community
Tree-/shrub covered	<i>Empetrum-Cornus suecica-Sphagnum fallax</i> community <i>Molinia caerulea-Narthecium ossifragum</i> community
Intermediate fen	
Open	
Mud bottom	<i>Equisetum fluviatile-Utricularia intermedia-</i> <i>Scorpidium scorpioides</i> community <i>Carex limosa-C. livida</i> community
Carpet	<i>Molinia caerulea-Sphagnum subnitens</i> community
Lawn	<i>Molinia caerulea-Scirpus caespitosus-Sphagnum</i> <i>papillosum-Sphagnum sect. Subsecunda</i> community
Rich fen	
Open	
Mud bottom	<i>Carex rostrata-Equisetum fluviatile</i> community
Mud bottom/carpet	<i>Scorpidium scorpioides</i> community
Lawn	<i>Carex dioica-Sphagnum subnites-S. warnstorffii</i> community <i>Molinia caerulea-Scirpus caespitosus-Campylium</i> <i>stellatum</i> community
Tree-/shrub covered	<i>Molinia caerulea-Potentilla erecta-Campylium</i> <i>stellatum</i> community
Extremely rich fen	
Open	
Carpet	<i>Carex limosa-Scirpus quinqueflorus-Scorpidium</i> <i>scorpioides</i> community
Lawn	<i>Campylium stellatum-Drepanocladus intermedius</i> community <i>Schoenus ferrugineus-Campylium stellatum-</i> <i>Drepanocladus intermedius</i> community
Tree-/shrub covered	<i>Potentilla erecta-Campylium stellatum-Drepano-</i> <i>cladus intermedius</i> community

OMBROTROPHIC VEGETATION

Large ombrotrophic mire complexes are common in the lowest parts of the area (150-200 m a.s.l.). They are all of the excentric raised bog type, with variously developed hummock banks. The lagg is narrow, approximately 1 m wide. Inside the lagg is a 10-30 m broad zone with pine forest and heather dominant in the field layer.

No outstanding floristic feature unites ombrotrophic vegetation as a whole (Fransson 1972). Only *Andromeda polifolia* and *Eriophorum vaginatum* grow throughout; along the mud bottom-hummock gradient as well as the mire expanse - mire margin gradient. *Drosera rotundifolia* is found along the mud bottom - hummock gradient, but it is a preferential mire expanse species.

The most common dominants in ombrotrophic vegetation are *Sphagnum cuspidatum* and *Carex limosa* in the lowest levels of hollows, *S. magellanicum* in carpets, *Scirpus caespitosus* and *S. tenellum* in lawns, and *Calluna vulgaris*, *Racomitrium lanuginosum*, *S. fuscum*, and *Cladonia* species in hummocks.

Although no outstanding floristic feature is common between hollows and hummocks, it is reasonable to emphasize the close interdependence between hollows and hummocks, together forming an ecological whole (see Du Rietz 1949). In Cajander (1913) and Nordhagen (1943) a separate phytosociological handling of hollow and hummock vegetation has obscured "Die Mineralbodenwasserzeigergrenze" (Du Rietz 1949). These authors have joined the ombrotrophic hollow vegetation with corresponding fen vegetation.

Compared with the regional complexes in the IBP-proposal (Malmer 1973) the complexes within the investigated area are closest to the *Rubellum-fuscum* complex, with *Sphagnum magellanicum* as an important hollow dominant. *Racomitrium lanuginosum* is a common hummock dominant on some mires (cf. *Racomitrium*-complex sensu Malmer 1973, see also Flatberg (1976)). Concerning the *Rubellum*-complex type "Komosse", Osvald (1923)) some of its oceanic species, *Erica tetralix* and *Sphagnum imbricatum* are lacking, except for *S. imbricatum* that occurs twice within the investigated area (K.I. Flatberg, pers. inform). Moreover, *Sphagnum rubellum* is not dominant in hummocks.

The *Cladonia* community and *Sphagnum fuscum* community form a transition between the mire expanse and mire margin vegetation.

The *Vaccinium-Cornus suecica-Hylocomium splendens* community is a pure margin community.

Open bog

Hollow vegetation

Scheuchzeria palustris-Sphagnum cuspidatum community (Tab VII).

The relevés are from a wet hollow mud bottom in the eastern part of the area (350 m a.s.l.).

The level of the ground water is high; *Sphagnum cuspidatum* floats in the water. *S. cuspidatum*-dominated communities are more common in ombrotrophic mires than in fens, and this species is one of the few preferentially ombrotrophic species.

The *Scheuchzeria palustris-Sphagnum cuspidatum* community is limited to small, wet mud bottom hollows on ombrotrophic mires.

Osvald (1923) reports a *Scheuchzeria palustris-Sphagnum cuspidatum* association from Komosse, in which sometimes only the two species naming the association occur. *S. cuspidatum* vegetation with *Scheuchzeria palustris* is described by Melin (1917), Sjörs (1948), Eurola (1962), Malmer (1962 a), and Fransson (1972). *S. cuspidatum* has a southerly Fennoscandia distribution (Malmer 1966), and *S. cuspidatum* dominated communities in bogs do not occur in Northern Finland (Eurola 1962).

Carex limosa-Cladopodiella fluitans/Gymnocolea inflata community (Tab. VII)

In mud bottoms with oxidized peat the hepaticas *Cladopodiella fluitans/Gymnocolea inflata* are among the most common dominants. The community has a patchy distribution, often with very small stands. Where the two hepaticas are not dense, the peat is covered by a layer of algae. *Sphagnum* species have scattered occurrences. The field layer is sparsely developed and may be absent. Species occurring are *Andromeda polifolia*, *Carex limosa*, and *Drosera anglica*.

The *Cladopodiella fluitans/Gymnocolea inflata-Drepanocladus fluitans* community of Flatberg (1970) is analogous to this community. An alga taxon is often used to name the community. Fransson (1972)

divides his *Cuspidatetum*-association into three subassociations, of which one is *zygogonietosum* (see also Sjörs 1948).

Carex limosa-Rhynchospora alba-Drepanocladus schulzei-
Cladopodiella fluitans/Gymnocolea inflata community (Tab. VII)

The transition between mud bottom and carpet is gradual. The *Sphagnum* mat is split up, and the field layer is sparse. Constant species are *Carex limosa*, *Rhynchospora alba*, *Drepanocladus schulzei*, and *Cladopodiella fluitans/Gymnocolea inflata*. *Eriophorum vaginatum* indicating a higher level than the two previously mentioned communities. Mat-forming *Sphagna* and species such as *Eriophorum vaginatum*, *Sphagnum lindbergii*, and *Drepanocladus schulzei* differ from these communities.

Two variants with *Sphagnum cuspidatum* and *S. papillosum* dominance, respectively, are separated. The *S. cuspidatum* variant is a transition community between mud bottom and carpet.

In eastern Fennoscandia *S. papillosum* and *S. pulchrum* grow only in fens (Ruuhiärv 1960). Both species occur in the *S. cuspidatum* variant, which is presumed to be an ombrotrophic community. Both are found in ombrotrophic mires in western Norway (Skogen 1966, Flatberg 1970, Moen & Selnes 1979) and in South-Western Sweden (Fransson 1972).

The *Eriophorum vaginatum-Sphagnum cuspidatum* association of Svensson (1965) is similar.

Eriophorum vaginatum-Sphagnum cuspidatum-S. majus-
S. pulchrum community (Tab. VII)

In this community the importance of *Sphagnum cuspidatum* diminishes and other *Sphagna* predominate, primarily *S. pulchrum* and *S. major*.

The ground water level is high, during most of the growing season water occurs a few cm below the top of the *Sphagnum* mat. *S. pulchrum* grows in a zone above the *cuspidatum* level in hollow pools. *S. majus* individuals grow scattered in the *pulchrum* mat.

Carex limosa and *Eriophorum vaginatum* are constans together with the three *Sphagnum* species mentioned. *S. pulchrum*, dominant in ombrotrophic carpets, is described by Skogen (1966), and *S. pulchrum* is frequently reported in bogs in western Norway (Flatberg 1976).

Sphagnum magellanicum-S. pulchrum community (Tab. VII)

Sphagnum magellanicum is one of the most common species in hollows in the area and the surrounding area (Klokk 1974). *S. magellanicum* has a wide amplitude along the mud bottom - hummock gradient, but has its main distribution in carpets. The ground water level is relatively high, 6-10 cm from the top of the *Sphagnum* mat during the growing season. The *S. magellanicum* mats are dense and other *Sphagnum* species grow sparsely. Vascular plants are not well established in these mats.

In addition to *S. magellanicum* the constants are *Andromeda polifolia*, *Eriophorum vaginatum*, and *S. pulchrum*.

The community has few differential species in comparison to other carpet communities, but it is best characterised by the dominance of *S. magellanicum*. *S. magellanicum* is reported as dominant in the upper hollow level in Rubello-Fuscion bogs (type "Skagerhult-mosse") (Du Rietz 1949) (see also Flatberg 1976). According to Sjörs (1948) *S. magellanicum* is common in large hollows in the western part of the Bergslagen area. Svensson (1965) and Fransson (1972) report *S. magellanicum*-dominated stands in a wide amplitude along the mud bottom - hummock gradient.

Scirpus caespitosus-Sphagnum rubellum-S. tenellum community (Tab. VII)

This is the lawn community of the hollows, and it is quantitatively the most important ombrotrophic community.

Sphagnum tenellum-dominated hollows are common in the investigated area. *Scirpus caespitosus* dominates the field layer and emphasizes the lawn character of the community. Other important field layer species are *Andromeda polifolia*, *Eriophorum vaginatum*, *Drosera anglica* and *D. rotundifolia*.

S. tenellum is dominant and constant in the bottom layer; *S. rubellum* is also constant. Several hepaticas, absent in the densest and most homogenous *Sphagnum* mats occur, viz. *Cephalozia* sp (p), *Cladopodiella fluitans/Gymnocolea inflata*, *Lepidozia setacea*, and *Ptilidium ciliare*.

S. tenellum-dominated hollow communities are common in Fennoscandia and are reported by Osvald (1923), Eurola (1962), Svensson (1965), Skogen (1966), Flatberg (1970), and Fransson (1972).

Hummock vegetation

Calluna vulgaris-Drosera rotundifolia-Rubus chamaemorus-Sphagnum fuscum community (Tab. VIII)

This is quantitatively the most important hummock community on the large excentric raised bogs in the lower parts of the area.

Calluna vulgaris and *Sphagnum fuscum* are dominants in the field and bottom layer, respectively. *Sphagnum rubellum* grows in the lower part of *fuscum* hummocks and is constant in the community. Other frequent species are *Empetrum* sp (p), *Oxycoccus microcarpus*, *Eriophorum vaginatum*, *Drosera rotundifolia*, *Rubus chamaemorus*, and *Mylia anomala*. *Scirpus caespitosus*, *Sphagnum fuscum*, *S. rubellum*, *Lepidozia setacea*, and *Mylia anomala* are preferentially differential species distinguishable from other hummock communities. Stand (B) is a *Myrica* variant of an ombrotrophic site in an ombro-minerotrophic complex. Occurrences of *Carex pauciflora*, *Myrica gale*, and *Calliergon stramineum* suggests this stand may not be ombrotrophic. In eastern Fennoscandia these species are considered to grow only in fens. Sjörs (1948) uses them to differentiate between species of fens and bogs. He also mentions that in South-West Sweden *Myrica* may grow in ombrotrophic sites (see also Skogen (1966) and Flatberg (1976) from western Norway, and Sjörs (1971)). Flatberg (1970) reports *Carex pauciflora* and *Calliergon stramineum* ombrotrophic in Nordmyra (10 km west of the investigated area).

Both Sjörs (1948) and Fransson (1972) assign all ombrotrophic hummock vegetation to one association. Fransson (1972) separates *Calluno-fuscetum* into five facies according to different dominants in the bottom layer. His *S. fuscum* facies are phytosociologically close to the *Calluna vulgaris - Drosera rotundifolia - Rubus chamaemorus - Sphagnum fuscum* community. The *Calluna vulgaris - Drosera rotundifolia - Rubus chamaemorus - Sphagnum fuscum* community of Flatberg (1970) is also a parallel.

Calluna vulgaris-Pleurozium schreberi-Polytrichum juniperinum var. gracilis community (Tab. VIII)

This community occurs on hummocks on some of the large excentric raised bogs, and is characterized by the dominance of *Pleurozium schreberi*. Other constants are *Andromeda polifolia*, *Calluna*

vulgaris, *Empetrum* sp(p), *Eriophorum vaginatum*, and *Polytrichum juniperinum* var. *gracilius*. The low importance of *Sphagnum* and *Cladonia* species is a negative characteristic of the community. *P. schreberi* and *P. juniperinum* var. *gracilius* are preferentially differential species distinguishing from other communities.

Pleurozium schreberi-dominated ombrotrophic mire expanse hummock vegetation is described by Sjörs (1946), Persson (1961), Sonesson (1970), Flatberg (1970, 1976), Moen et al. (1976), and Moen & Selnes (1979).

Dicranum leioneuron-Sphagnum nemoreum community (Tabe. VIII)

Some hummock tops are totally dominated by *Dicranum leioneuron*. This hummock vegetation is observed in only one of the eccentric raised bogs. *D. leioneuron* dominates hummocks in the same amplitude as *Pleurozium schreberi*, and they dominate alternate hummocks on the same mire. *D. leironeuron* and *Sphagnum nemoreum* are preferentially differential species distinguishing the community from other hummock communites.

The *Cetraria islandica* variant occurs at somewhat lower level as indicated by *Scirpus caespitosus*.

Racomitrium lanuginosum-Cladonia community (Tab. VIII)

This is a common hummock community which occurs at higher altitudes (350-500 m a.s.l.). In the large eccentric complexes at lower altitudes *Racomitrium lanuginosum* rarely dominates the hummocks, this indicates that the higher altitudes have a more humid climate (cf. precipitation values of the station Vennafjell, Fig. 3).

Cladonia alpestris, *C. rangiferina*, *C. arbuscula*, and *R. lanuginosum* are alternating dominants in the bottom layer. *Racomitrium* hummocks are large with some margin character indicated by *Vaccinium myrtillus* and *V. uliginosum*. A differential species distinguishable from other hummock vegetation is *R. lanuginosum*, and the three mentioned *Cladonia* species are preferential species. The two relevés 31 and 32 (Tab. VIII) are from dense stands of *Cladonia alpestris*, where other cryptogams are scarce. However, scattered individuals of *Pleurozium schreberi* and *Ptilidium ciliare* are characteristic of *C. alpestris* hummocks.

R. lanuginosum-dominated hummock vegetation is an oceanic

feature of ombrotrophic mire vegetation in Fennoscandia, reported only from coast areas or other humid areas (Skogen 1966, Flatberg 1970, 1976, Moen & Selnes 1979).

Tree-/shrub-covered bog

Cladonia community (Tab. IX)

This community is phytosociological closely related to the *Cladonia*-dominated mire expanse vegetation. *Racomitrium lanuginosum* is more scattered than in corresponding mire expanse vegetation (mire expanse in a geographic sense) (see also Flatberg 1970).

Three *Vaccinium* species indicate the margin character, but the phytosociological difference from the mire expanse *Racomitrium lanuginosum*-*Cladonia* community is not significant.

Sphagnum fuscum community (Tab. IX)

Sphagnum fuscum has a strong affinity with mire expanse. The margin character is weak, only indicated by *Vaccinium* species.

Near the margin of the mire *S. fuscum* grows in small flat hummocks, in time probably becoming pure margin vegetation. Phytosociologically the *Sphagnum fuscum* community may be listed to the *Calluna vulgaris*-*Drosera rotundifolia*-*Rubus chamaemorus* community.

Vaccinium-Cornus suecica-Hylocomium splendens community (Tab. IX)

This is the only pure mire margin community on ombrotrophic mire. The community usually occurs in a zone between the mire expanse and the mineral soil, but sometimes it occurs in depressions isolated from a mire expanse.

The tree layer consists of pine trees up to 15-20 m in height. *Picea abies* occurs scattered in the shrub layer, rarely higher than 1 m. *Calluna vulgaris*, *Empetrum hermaphroditum*, and the *Vaccinium* species dominate the field layer. The bottom layer is well developed with a mixture of spruce forest mosses and oligotrophic *Sphagna*.

The margin character is strong, and the following species are differential species distinguishing from the two previous communities and the mire expanse hummock vegetation: *Picea abies*,

Deschampsia flexuosa, *Cornus suecica*, *Dicranum majus*, *Hylocomium splendens*, and *Barbilophozia lycopodioides*.

The *Vaccinium-Cornus suecica-Hylocomium splendens* community may be assigned to *Vaccinio uliginosi-Pinetum* (Kleist 1929). Out of the 21 species Kielland-Lund (1973) assigns to this association, 18 are common in the *Vaccinium-Cornus suecica-Hylocomium splendens* community. Aune (1973) mentions associations described in the literature synonymous with *Vaccinio uliginosi-Pinetum*. In addition to those mentioned by Aune (1973), several papers on mire vegetation describe similar vegetation. From Finland i.a. Ruuhijärvi (1960) reports phytosociologically similar vegetation. As mentioned in his paper *Ledum palustre* and *Carex globularis* are common in this vegetation in eastern Fennoscandia (see also Sjörs 1948).

The *Pinus-Vaccinium* association of Sjörs (1948) and *Vaccinietum uliginosi* association of Fransson (1972) show good correlations with the described *Vaccinium-Cornus suecica-Hylocomium splendens* community.

Compared with analogous vegetation in Sjörs (1948), Ruuhijärvi (1960), Fransson (1972) (Tab. IX) shows a higher frequency of *Deschampsia flexuosa*, and the mosses *Hylocomium splendens*, *Pleurozium schreberi*, and *Ptilium crista-castrensis* are more frequent dominants than *Sphagnum*.

POOR FEN

In the eastern parts of the area slightly sloping poor fens predominate the mire vegetation. Some of these are former hay cut areas. Fens with a greater angle of inclination are usually rich.

Most of the species growing in poor fens also grow in bogs, the poor fens differ from the latter by stable occurrences of minerotrophic indicators. The most common indicators are *Carex lasiocarpa*, *C. rostrata*, *Molinia caerulea*, and *Menyanthes trifoliata*. All of them except for *Menyanthes*, are relatively indifferent along the mud bottom - hummock gradient.

Poor fens differ from intermediate and rich fens since they lack eutrophic species. Poor fens have no positive differential species distinguishable from richer fens.

The division between bogs and fens is stressed by the strict use of character species of minerotrophic vegetation. This is considered to be one of the most important dividing lines in mire vegetation (Du Rietz 1949). As mentioned by Flatberg (1970) situations may occur where a thick layer of ombrogenic peat causes ombrogenic wet of soil, but species with deep roots nevertheless reach ground water influenced by minerotrophy. These mires are regarded as fens (cf. "extreme poor fen" Du Rietz 1949) since indicators of minerotrophy are present.

Poor fens are negatively distinguishable from richer fens by the less exclusive eutrophic indicators of the intermediate fen. The "transitional poor fen" (övergångsfattigkärr") of Witting (1947) is regarded as intermediate fen (see also Du Rietz 1949).

The most common species in the tree/shrub layer of poor fens are *Betula pubescens* and *Picea abies*. The margin character of the *Empetrum-Cornus suecica-Sphagnum fallax* community is distinct. The *Molinia caerulea-Narthecium ossifragum* community is, however, a transitional community between mire margin and mire expanse.

Open poor fen

Carex limosa-Menyanthes trifoliata community (Tab. X)

The analyses are from wet flarks in a poor flark fen. The ground water was exposed, 0-10 cm deep water covering dark brown mud. The community represents the wet end of the mud bottom - hummock gradient. Mud bottom communities of this kind have no net peat production (Sjörs 1948).

Constants are *Carex lasiocarpa*, *C. limosa*, and *Menyanthes trifoliata*. *Utricularia intermedia* in stand B indicates more eutrophic conditions in this stand. It is a common problem to place communities extremely poor in species along the poor - rich gradient. The wet flark vegetation shows relatively little variation according to different nutrient condition of adjacent mire vegetation.

Eriophorum angustifolium-Sphagnum pulchrum community (Tab. X)

Both the separated variants represent lowest level of closed *Sphagnum* mats, the carpet level.

Sphagnum lindbergii is most frequent at higher altitudes in the area. The analyses of the *Sphagnum lindbergii* - *Drepanocladus schulzei* variant are recorded on an extremely poor fen, 440 m a.s.l. *Eriophorum angustifolium* and *Menyanthes trifoliata* are the only indicators of minerotrophy. Hepaticas have difficulties in establishing themselves in dense *S. lindbergii* mats.

S. lindbergii has a northern distribution in Fennoscandia (Sjörs 1948, Malmer 1966, Fransson 1972). In the "Sphagnum-Rimpi-weissmore" (Ruuhiärvi 1960) *S. lindbergii* gradually increases in importance towards the north.

The *Sphagnum lindbergii-Drepanocladus schulzei* variant is structurally similar to *Carex rotundata*-*D. schulzei* vegetation of Sonesson (1970), where *D. schulzei* and *S. lindbergii* dominate in a separated *S. lindbergii* variant, and *Sphagnum lindbergii*-vegetation of Elveland (1976).

The *Sphagnum pulchrum* variant is uninfluenced by altitude within the area (see also Fransson 1972). *S. pulchrum* has its main distribution in south-western parts of Fennoscandia (Waldheim & Weimark 1943, Sjörs 1948).

Carex rostrata-Sphagnum riparium community (Tab. X)

Sphagnum riparium rarely dominates in mire vegetation in the area. The analyses are based on a carpet stand adjacent to a small lake. Mean ground water level was 12 cm below the top of the *Sphagnum* mat on the day the analyses were recorded. The community is poor in species (cf. Sonesson 1970) and is homogenous. Except for scattered individuals of *Drepanocladus schulzei*, the moss mat is a pure population of *S. riparium*. The homogeneity is stressed by the fact that all four field layer species are constants.

S. riparium-dominated communities are described by e.g. Sjörs (1948), Persson (1961), and Sonesson (1970).

Drosera rotundifolia-Sphagnum magellanicum community (Tab. X)

This is a transition community between carpet and lawn. *Carex limosa* and *Drosera anglica* are preferentially differential species of carpet and mud bottom as opposed to lawn. *Sphagnum fuscum* and *S. rubellum* grow scattered on small elevations indicating a succession towards hummock vegetation. *S. magellanicum* is more dominant in bogs than in fens in the area (see also Klokk 1974).

Carex limosa-Sphagnum papillosum-S. majus community (Tab. X)

Sphagnum papillosum prefers lawns, but it is here dominant in a typical carpet community. The analyses are from a stand near a small lake, where *S. papillosum* mats form the lowest *Sphagnum* level. *Carex limosa*, *Drepanocladus schulzei*, and *Sphagnum majus* emphasize the carpet character.

The carpet association *Carex rostrata-pauciflora-Sphagnum papillosum* (Sjörs 1948) and parts of *Rhynchospora-papilletosum* (Fransson 1972) are similar vegetation types.

Scirpus caespitosus-Sphagnum compactum-Cladopodiella fluitans/Gymncolea inflata community (Tab. XI)

Sphagnum compactum and *Cladopodiella fluitans/Gymncolea inflata* dominated fens cover large areas at high altitudes in the area. *S. compactum* is the most common *Sphagnum* dominant in lawns on extremely poor fens. A characteristic feature of the community is a split *Sphagnum* mat, with small spots of open oxidized peat, often with *Cladopodiella fluitans/Gymncolea inflata*. This description also characterises the locality of *Lycopodium inundatum* (see also Sjörs 1948).

Increased frequency of *S. compactum* dominated communities at higher altitudes is mentioned by e.g. Ruuhijärvi (1960) and Fransson (1972). Both the *Sphagnetum compacti*-association of Fransson (1972) and the *Sphagnum compactum*-"Weissmoore" of Ruuhijärvi (1960) have the same floristic characteristics as the present community.

Scirpus caespitosus-Sphagnum tenellum community (Tab. XI)

Sphagnum tenellum and *S. papillosum* are the most common dominants in lawns of poor fens.

Constants are *Andromeda polifolia*, *Eriophorum vaginatum*, *Scirpus caespitosus*, *Drosera rotundifolia*, *Sphagnum tenellum*, and *Cladopodiella fluitans/Gymncolea inflata*. *Scirpus caespitosus* is the field layer dominant. The following species are preferentially differential species distinguishable from other lawn poor fen vegetation, *Empetrum* sp(p), *Oxycoccus microcarpus*, *O. quadripetalus*, *Eriophorum vaginatum*, *Scirpus caespitosus*, *Sphagnum tenellum*, *Ptilidium ciliare*, *Lepidozia setacea*, and *Mylia anomala*.

Sphagnum subnitens occurs in one stand. This species is in Fennoscandia often regarded as a characteristic species of intermediate fens. In the investigated area, *S. subnitens* also grows scattered in poor fens. In western Norway *S. subnitens* is also found in bogs (Skogen 1969, Flatberg 1976).

The main distribution of the *Sphagnum tenellum* variant occurs at a lower level than the *S. papillosum* variant, this is indicated by the occurrence of *Drosera anglica*, *Drepanocladus schulzei*, and by *Sphagnum majus* in the former and *S. rubellum* in the latter. The community is similar to *S. tenellum* dominated hollow vegetation.

Both Sjörs (1948) and Fransson (1972) mention two lawn communities in poor fens with *Scirpus caespitosus*, *Sphagnum papillosum*, and *S. tenellum* dominance. Although *Molinia caerulea* occurs in Table XI, the two communities without *Molinia*, viz. *Scirpus caespitosus-Carex pauciflora-Sphagnum tenellum-papillosum* association (Sjörs 1948) and *Trichophoro-papilletosum* (Fransson 1972), are most similar.

Molinia caerulea-Scirpus caespitosus-Narthecium ossifragum
community (Tab. XI)

Narthecium mats usually occur in patches in mire vegetation, but sometimes they cover the whole mire. This is a characteristic lawn community with dense firm mats of *Narthecium* and a sparse bottom layer. The community is poor in species and is homogenous. *Andromeda polifolia* has its highest mean cover value in this community.

Narthecium-dominated communities in Fennoscandia only occur in humid areas, Skogen (1976), Moen (1975), and Flatberg (1976) from western Norway, Fransson (1972) from south-western Sweden, and Sjörs (1948) and Fransson (1963) from western Jämtland, Central Sweden.

Tree-/shrub covered poor fen

Empetrum-Cornus suecica-Sphagnum fallax community (Tab. XII)

The most frequent field layer dominants are *Calluna vulgaris*, *Scirpus caespitosus*, and *Rubus chamaemorus*, and *Sphagnum fallax*

is the bottom layer dominant. The margin character is indicated by *Betula pubescens*, *Vaccinium* spp., *Cornus suecica*, *Dicranum majus*, *Pleurozium schreberi*, *Sphagnum fallax*, *S. russowii*, and *Obtusifolium obtusum*. These are also preferential or exclusive differential species distinguishing from poor fen mire expanse communities.

Betula-Eriophorum vaginatum-Carex pauciflora-fusca-Sphagnum parvifolium association (Sjörs 1948) and *Pauciflora-parvifolietum* association (Fransson 1972) are similar communities.

Molinia caerulea-Narthecium ossifragum community (Tab. XII)

The margin character is weak, indicated by *Vaccinium* spp. *Narthecium* totally dominates the field layer, and the bottom layer is sparse. Phytosociologically this forms a transition between mire margin and mire expanse, and the floristic difference against the mire expanse *Molinia caerulea-Scirpus caespitosus-Narthecium ossifragum* community is small.

INTERMEDIATE FEN (OPEN)

Intermediate fens are most common in eastern parts of the area at altitudes 300-400 m a.s.l.

Intermediate mud bottoms and carpets do not cover large continuous areas but occur in small flarks. On "ribbed" mires with low hummock banks ("ribs") the flarks often have intermediate vegetation and the "ribs" poor fen vegetation. Situations also occur with rich fen vegetation in the lower levels and intermediate fen vegetation in the higher levels, i.e. *Sphagnum subnitens* mat surrounded by *Scorpidium* or *Campylium* vegetation.

A characteristic feature of intermediate fens is the simultaneous occurrence of poor fen species and less exclusive rich fen species (Sjörs 1948, Flatberg 1970). The intermediate fen has no exclusive character species sensu stricto, but *Sphagnum subnitens* has its main occurrence there. The less exclusive rich fen species are differential species distinguishing from poor fens, viz. *Carex dioica*, *Selaginella selaginoides*, *Tofieldia pusilla*, *Campylium stellatum*, *Drepanocladus badius*, *D. exannulatus*, *D. revolvens* s.str., *Scorpidium scorpioides*, *Sphagnum* Sect. *Subsecunda* species, and *S.*

warnstorffii. Most of the field layer dominants also grow in poor fens, i.e. *Carex rostrata*, *Molinia caerulea*, and *Scirpus caespitosus*. Some species grow in poor and intermediate fens, but not in rich and extremely rich fens (Tab. II).

The amplitude of the intermediate fen along the poor-rich gradient is in accordance with Sjörs (1952). The transition towards poor fens is gradual within the area, sometimes with just one scattered indicator species of intermediate fens. This parallels the relationship of the extremely poor fen to ombrotrophic mire. Localization along the poor - rich gradient has been strictly defined according to the indicator value of the species indicating the highest value of eutrophy. Mud bottom and lawn communities are difficult to separate from poor fens. Many of the slightly sloping fens (3-4° slope) in the eastern parts of the area are "poor" intermediate fens.

In Fennoscandina papers before 1952, where vegetation is analysed along gradients, fens are classified as poor, medium rich, and extremely rich fens (i.e. Du Rietz 1949). In 1952 Sjörs wrote "... but in North Fennoscandia, there is often a kind of fen where some of the less exclusive species of rich fens mix with species of poor fens ...". Sjörs (1952) notes that several authors described communities with the species combination mentioned, e.g. Cajander (1913), Melin (1917), Nordhagen (1928), Kivinen (1935), and Sjörs (1946, 1948, 1950). Sjörs (1952) introduced the term "intermediate fen" for these mires. Definitions in the present paper of the less exclusive rich fen species are in accordance with Sjörs (1952), except for *Carex flava*, *Saussurea alpina*, and *Paludella squarrosa* which in the present paper are regarded as rich fen species.

The northerly distribution tendency of intermediate fens is also emphasized by Malmer (1973).

Equisetum fluviatile-Utricularia intermedia-Scorpidium scorpioides community (Tab. XIII)

Field and bottom layers are sparse. Differential species distinguishing from poor fen mud bottom are *Utricularia intermedia*, *Drepanocladus exannulatus*, and *Scorpidium scorpioides*. *S. scorpioides* has its main distribution in rich and extremely rich fens, but occurs scattered in intermediate fens. The floristic difference from rich mud bottom communities is small.

Carex limosa-C. livida community (Tab. XIII)

Carex livida-dominated flarks are only observed at higher altitudes in the eastern parts of the area.

Ground water level was exposed the day the analysis was recorded. The bottom layer is sparse. *Cladopodiella fluitans/Gymnocolea inflata* are the only bryophytes. The field layer is homogenous with *Carex livida* as the only dominant.

C. livida dominated mire vegetation is common in subalpine and alpine areas, see Nordhagen (1943). Sjörs (1948) mentions *c. livida* in all of his three mud bottom associations, but the most similar is the *Carex limosa-chordorrhiza-livida-mud* bottom-association. Almost identical is the *Carex limosa-C. livida* mud bottom reported by Elveland (1976).

Molinia caerulea-Sphagnum subnitens community (Tab. XIII)

The community is dominated by dense brown-violet mats of *Sphagnum subnitens*. *S. subnitens* has a clumped distribution, forming a mosaic with wetter surroundings. Stand F deviates by having a higher number of species. It should be regarded as a transitional community towards lawn.

Field layer dominants are *Molinia caerulea*, *Scirpus cæspitosus*, and *Menyanthes trifoliata*; with *Myrica* dominating in one stand. Indicator species of intermediate fen are not dominants, except for *S. subnitens*. Differential species distinguishing from poor fens are *Carex dioica*, *Selaginella selaginoides*, *Tofieldia pusilla*, *Campylium stellatum*, *Drepanocladus badius*, *D. revolvens* s.str. *Sphagnum* Sect. *Subsecunda* species, and *S. warnstorffii*. *Campylium stellatum* and *Drepanocladus revolvens* s.str. only grow scattered in intermediate fens. Preferentially differential species are *Viola palustris*, *S. subnitens*, and *Odontoschisma elongatum*.

The transition towards lawn is gradual, represented by stand F, and differential species distinguishing from intermediate lawn are few. *Carex limosa* is the only exclusive species and *Menyanthes* is a preferential differential species.

Flatberg (1970) describes several *S. subnitens* dominated communities in intermediate fens, where *S. warnstorffii* is codominant in most of them (see also Moen 1970). The *Molinia caerulea-Sphagnum subnitens* community is similar to *S. subnitens* dominated parts of

the *subfulvetum*-association of Fransson (1972). Sjörs' (1948) *Scirpus caespitosus-Molinia-Sphagnum papillosum*-association is also similar in spite of other dominants in the bottom layer.

Molinia caerulea-Scirpus caespitosus-Sphagnum papillosum-
S. Sect. Subsecunda community (Tab. XIII)

Intermediate lawn vegetation is common in the eastern part of the area, sometimes covering the whole mire. This vegetation has few indicator species of intermediate fen vegetation.

Frequent species which are also found in poor fens are *Molinia caerulea*, *Scirpus caespitosus*, *Drosera rotundifolia*, *Sphagnum papillosum*, and *Cladopodiella fluitans/Gymnocolea inflata*. Exclusive differential species distinguishing from poor fens are *Selaginella selaginoides*, *Campylium stellatum*, *Drepanocladus badius*, and *Sphagnum Sect. Subsecunda* species.

Similar vegetation is the *Scirpus caespitosus-Molinia-Sphagnum papillosum*-association (Sjörs 1948) and the *Scirpus caespitosus-Sphagnum papillosum-rubellum-subnitens* community (Flatberg 1970).

RICH FEN

Rich fens have many species in common with extremely rich fens, but they lack the exclusive character species of the latter. The field layer dominants are the same as in intermediate and poor fens, but the bottom layer dominants differ. *Scorpidium scorpioides*, *Campylium stellatum*, and *Drepanocladus* species replace the *Sphagna*. Differential (preferential-p) species distinguishing from intermediate fens are *Carex flava*, *Eriophorum latifolium*, *Thalictrum alpinum*, *Campylium stellatum* (p), *Drepanocladus revolvens* s.str. (p), *D. intermedius*, *Scorpidium scorpioides* (p), and *Leiocolea borealis*.

Until the separation of intermediate fens (Sjörs 1952), the rich mire vegetation was classified as "transition rich fen" ("övergångsrikkärr") and extremely rich fen (Witting 1947, Du Rietz 1949). In the present paper rich fen is used synonymously with "medium rich fen" (Sjörs 1967, Fransson 1972, Malmer 1973). Compared with the hierachial system of Nordhagen (1943), the rich fen occurs in several alliances, however, the main part is found in *Caricion*

canescentis-Goodenowii, and mud bottom and carpet communities in *Stygio-Caricion limosae*.

Open rich fen

Carex rostrata-Equisetum fluviatile community (Tab. XIV)

Ground water was exposed on the day the analyses were recorded. A brook runs close by and causes periodic limnic wet of soil. The community represents a transition towards pure limnic vegetation. The low number of species gives little data for the placing of the community along the poor - rich gradient. Although *Bryum pseudotriquetrum* is a preferential, extremely rich fen indicator, the community is best included in the rich fen category.

Scorpidium scorpioides community (Tab. XIV)

Scorpidium scorpioides occurs mainly in carpets and grows scattered in mud bottoms and lawns. Stands B, C, and D are transitions towards mud bottom with a discontinuous bottom layer. The *Scorpidium scorpioides* community includes all *Scorpidium* dominated vegetation on rich fen (Persson 1961, Sonesson 1970, Moen 1970).

In the bottom layer *Drepanocladus revolvens s.str.* is sometimes co-dominant. The field layer dominants are the same as in poor and intermediate fens, indicators of rich fen may be absent. The qualitative difference from intermediate mud bottom - carpet vegetation is not significant, but *S. scorpioides* and *D. revolvens s. str.* never dominate in intermediate fens.

The *Rhynchospora fusca* variant occurs as small islands in mosaic with mud bottom vegetation. *Odontoschisma elongatum* is here noted with its highest cover value. *R. fusca* seems to have the same amplitude as *R. alba* along the mud bottom - hummock gradient. The phytosociological relationship with extremely rich *Scorpidium* vegetation is obvious, but the *Scorpidium scorpioides* community lacks the exclusive indicators of extremely rich fens.

Carex dioica-Sphagnum subnitens-S. warnstorffii community (Tab. XIV)

This community is uncommon and covers only small areas. It

is an upper lawn community with scattered, lowgrowing *Calluna*. *Sphagnum warnstorffii* mats are similar in appearance to *S. subnitens* mats. The field layer species are mainly the less exclusive rich fen species, viz. *Carex dioica*, *Tofieldia pusilla*, *Drepanocladus badius*, and *Sphagnum subnitens*. Dominance of *S. warnstorffii* and constant occurrence of *Campylium stellatum* are reasons for including the community in the rich fen category.

In the series *Scorpidium-Drepanocladus-Campylietum-Tomen-typnum-Fuschetum* (Du Rietz 1949) the present community corresponds to *Tomentypnum* level. Flatberg (1970) separates three *S. warnstorffii* dominated communities, where the two intermediate fen communities belong to higher levels. Fransson (1972) includes corresponding vegetation in *Tomentypnetum* (see also Du Rietz 1949). Fransson (1972) emphasizes the northern distribution tendency of *S. warnstorffii*.

Molinia caerulea-Scirpus caespitosus-Campylium stellatum community (Tab. XIV)

The change in vegetation from *Scorpidium* to *Campylium* level is gradual. *Drepanocladus revolvens* s. str. occurs mainly in an intermediate zone between these levels. *Scirpus caespitosus* and *Campylium stellatum* are the most frequent dominants in rich fen lawn vegetation in the area. Differential species distinguishing from intermediate lawn vegetation are *Eriophorum latifolium*, *Thalictrum alpinum*, *Drepanocladus intermedius*, and *Leiocolea borealis*. *D. intermedius* mainly occurs in extremely rich fens.

Campylium dominated vegetation is described by Booberg (1939), Sjörs (1948), Ruuhijärvii (1960), Flatberg (1970), Moen (1970), and Fransson (1972).

Tree-/shrub covered rich fen

Molinia caerulea-Potentilla erecta-Campylium stellatum community (Tab. XIV)

The community is similar to the mire expanse *Molinia caerulea-Scirpus caespitosus-Campylium stellatum* community with common dominants. Mire margin species are differential species distinguishing from this community, viz. *Carex echinata*, *Saussurea alpina*,

Trientalis europaea, *Calliergonella cuspidata*, and *Tritomaria quinquentata*. *Potentilla erecta* is preferential mire margin species. Communities mentioned in connection with *Molinia caerulea-Scirpus caespitosus-Campylium stellatum* community also have relevance to the present community.

EXTREMELY RICH FEN

Character species of extremely rich fens are few in the lowland of Central Norway, and some of them are rare. Mud bottoms and carpets in particular have few character species. The dominants are common with rich fens, particularly the bottom layer dominants.

The dense *Schoenus ferrugineus* mats form a characteristic community, with a very sparse bottom layer. Exclusive extremely rich fen indicator species in the area are *Carex lepidocarpa*, *C. hostiana*, *Schoenus ferrugineus*, and *Leiocolea rutheana*. Preferential indicators are *Scirpus quinqueflorus*, *Triglochin palustre*, *Bryum pseudotriquetrum*, *Cinclidium stygium*, and *Drepanocladus intermedius*.

Du Rietz (1949) regards *C. lepidocarpa* and *Schoenus ferrugineus* as indicators of extremely rich fens. Extremely rich fen in the IBP proposal (Malmer 1973), corresponds to this category in the present paper. The extremely rich fen communities within the investigated area are not dealt with by Nordahgen (1943), but a lowland equivalent, *Schoenion ferruginei*, to *Caricion atrofuscae-saxatilis* is mentioned. Differential species of the former distinguishing from the latter are *Schoenus ferrugineus* and *Carex hostiana*, and the extremely rich fen in the present paper can be assigned to *Schoenion ferruginei*.

Open extremely rich fen

Carex limosa-Scirpus quinqueflorus-Scorpidium scorpioides community (Tab. XV)

The analysed stand grows near a *Schoenus* stand. The community is homogenous, poor in species, and with a sparse field layer. The only indicator of extremely rich fen is *Carex lepidocarpa*.

Carex limosa, *Drosera anglica*, *Menyanthes trifoliata*, and *Scorpidium scorpioides* characterize the wet habitat conditions, all of them being strong preferential differential species distinguishing from lawn vegetation.

Several authors, dealing with mire vegetation, unite the *Scorpidium* vegetation (Du Rietz 1949, Persson 1961, Sonesson 1970, Moen 1970). A phytosociologically close relationship with *Scorpidium scorpioides* community from rich fen is obvious.

Campylium stellatum-Drepanocladus intermedius community (Tab. XV)

This is quantitatively the most important extremely rich fen community in the area. The vegetation of the steep sloping fens in western parts of the area should be included in this community. *Scirpus caespitosus* is dominant. Other frequent field layer species also grow in poor, intermediate, and rich fens. The two bottom layer dominants are *Campylium stellatum* and *Drepanocladus intermedius*. Extremely rich fen indicators are few and have low frequencies, viz. *Carex hostiana*, *C. lepidocarpa*, and *Leiocolea rutheana*.

Scirpus caespitosus and *S. hudsonianus* are differential species distinguishing from the other extremely rich lawn community. Du Rietz (1949) separates a *Drepanocladetum* and a *Campylietum* association with different amplitudes along the mud bottom - hummock gradient. Fransson (1972) finds it difficult to separate these levels and unites them to a *Campylium-revolutum* association. Other authors describing *Drepanocladus/Campylium* vegetation are Sjörs (1948), Persson (1961), Fransson (1963), Flatberg (1970), and Moen (1970).

Schoenus ferrugineus-Campylium stellatum-Drepanocladus intermedius community (Tab. XV)

Schoenus vegetation forms two characteristic types, viz. dense continuous mats and single tussocks surrounded by wet oxidized peat.

The community is poor in species and the bottom layer is sparse or absent, this is caused by the growth form of *Schoenus*. Except for *Schoenus* all species are common in other rich and extremely rich lawn communities.

Schoenus dominated vegetation is described by i.a. Sjörs (1948), Flatberg (1970), and Moen (1970).

Tree-/shrub covered extremely rich fen

Potentilla erecta-Campylium stellatum-Drepanocladus intermedius
community (Tab. XV)

This is the mire community most rich in species. Characteristic for extremely rich mire margin vegetation is the occurrence of species from eutrophic forests (Sjörs 1948). *Crepis paludosa*, *Filipendula ulmaria*, *Calliergonella cuspidata*, and *Mnium seligeri* are frequent, and they are differential species distinguishing from corresponding rich margin vegetation. The margin character is further characterized by *Salix aurita*, *Carex echinata*, *Hierochloë odorata*, *Juncus articulatus*, *Equisetum palustre*, *Galium boreale*, *Trientalis europaea*, and *Fissidens adianthoides*.

The IBP proposal has no synonymous community, however, the present community has common features with "brunmossrika skogskärrkompleks" and "örtrika skogskärrkompleks" (Malmer 1973). *Betula-Filipendula Ulmaria-Sphagnum Warnstorffianum* association (Sjörs 1948) is closely related to the present community.

THE FOREST VEGETATION

Systematics of the forest communities

Class *Vaccinio-Piceeta*

Order *Cladonio-Vaccinetalia*

Alliance *Phyllodoco-Vaccinion*

Association *Vaccinio uliginosi-Pinetum* (see the *Vaccinium-Cornus suecica-Hylocomium splendens* community, Tab. IX)

Association *Barbilophozio-Pinetum*

Subassociation *cladonietosum*

Subassociation *hylocomietosum*

Order *Vaccinio-Piceetalia*

Alliance *Vaccinio-Piceion*

Association *Chamaemoro-Piceetum*

Association *Eu-Piceetum*

Subassociation *myrtilletosum*

Subassociation *dryopteridetosum*
Subassociation *blechnetosum*
Association *Melico-Piceetum*
Subassociation *convallarietosum*
Subassociation *typicum*
Subassociation *calliergonelletosum*
Class *Alnetea glutinosae*
Order *Salicetalia auritae*
Alliance *Alno-Salicion pentandrae*
Association *Equiseto palustris-Salicetum*
Class *Querco-Fagetea*
Order *Fagetalia sylvatica*
Alliance *Alno-Padion*
Association *Alno (incanae)-Prunetum*
Subassociation *aconitetosum*
Alliance *Tilio-Acerion*
Association *Ulmo-Tilieturn boreale*

ASSOCIATION *CHAMAEMORO-PICEETUM* KIELLAND-LUND 1962 (Tab. XVI)

This is a mesotrophic damp forest on peat ground of varying thickness. Stands belonging to this association grow in depressions with stagnating ground water or in border areas between mires and mineral soil.

The tree layer is sparse with low growing *Betula pubescens*, *Picea abies*, and *Pinus sylvestris*.

In the field layer *Vaccinium myrtillus*, *Cornus suecica*, and *Rubus chamaemorus* are constants and dominants. Preferential character species, which also have their main distribution in this association are *Juncus filiformis*, *Equisetum sylvaticum*, and *Listera cordata*.

Sphagnum fallax and *S. girgensohnii* dominate in the bottom layer. *S. centrale*, *S. magellanicum*, and *S. nemoreum* are local dominants.

Constants in the table are *Empetrum sp(p.)*, *Vaccinium myrtillus*, *V. vitis-idaea*, *Cornus suecica*, *Rubus chamaemorus*, *Hylocomium splendens*, *Plagiothecium undulatum*, *Pleurozium schreberi*, *Sphagnum fallax*, and *S. girgensohnii*.

There is a mixture of species from mineral soil and mire species, but some species occur primarily in this community.

The most closely related mire community is the poor fen margin *Empetrum-Cornus suecica-Sphagnum fallax* community. Differential species which distinguish from this community are *Deschampsia flexuosa*, *Gymnocarpium dryopteris*, *Hylocomium splendens*, *Ptilium crista-castrensis*, *Rhytidadelphus loreus*, and *Tritomaria quinque-dentata*.

Differential species which distinguish from the oligotrophic damp forest *Vaccinio uliginosi-Pinetum* are *Betula pubescens*, *Carex echinata*, *C. nigra*, *Juncus filiformis*, *Gymnocarpium dryopteris*, *Equisetum sylvaticum*, *Listera cordata*, *Maianthemum bifolium*, and *Polytrichum commune*.

In the IBP proposal (Kielland-Lund 1971) the mesotrophic damp forest belongs to *Chamaemoro-Piceetum* Kielland-Lund 1962 in the forest system, and to "vitmossrika" and "risrika skogskärrkomplex" in the mire system (Malmer 1971). The community seems common in the boreale coniferous region (Arnborg 1943, Malmström 1949, Samuelsson 1960, Kielland-Lund 1962, Sjörs 1971, Kjelvik 1978). In the subalpine region a parallel birch forest occurs (see e.g. Moen & Moen 1975, Moen 1976).

ASSOCIATION *EQUISETO PALUSTRIS-SALICETUM* ASS. PROV. (TAB. XVII)

Eutrophic damp forests are rare within the area. Stands that belong to this community are of two kinds. The most common occurrence is in wet habitats with stagnating ground water, forming a mosaic with tree covered tussocks. In such locations (stands A and B) only the wet area between the tussocks has been analysed. The community also occurs on wet slopes (stands C and D) with a somewhat lower ground water level. Further investigations will probably show the need for more than one association of rich damp forests.

The most frequent tree layer species are *Alnus incana* and low growing *Picea abies*. In the shrub layer *Salix* species predominate.

Comarum palustre, *Menyanthes trifoliata*, and partly *Filipendula ulmaria* dominate the field layer in the wet mosaic type. The bottom layer of this type is totally dominated by *Mnium* species.

Generally the "wet part" of the mosaic type is more homogeneous and less rich in species than the "drier" type which resembles wet stands of *Melico-Piceetum-calliergonelletosum*, and the transition towards the latter is gradual.

In the IBP system (Kielland-Lund 1971, 1973) *Equiseto palustris-Salicetum* corresponds to *Calamagrostio (purpureae)-Salicetum pentandrae*. Further investigations are needed to clarify the phytosociology of these forests.

ASSOCIATION *BARBILOPHOZIO-PINETUM* BR.-BL. ET SISSINGH 1939 EM.
KIELLAND-LUND 1967 (TAB. XVIII)

This is the most oligotrophic forest association on mineral soil. The pine forest stands of this community are small, occurring in areas with shallow soil. Within the investigation area the community is most frequent at altitudes above 300 m. *Pinus sylvestris* competes unfavourably with *Picea abies* on moraine materiale in lower areas.

Tree and shrub layers are sparse, with *Pinus sylvestris* and scattered incidences of *Betula pubescens* and *Picea abies*.

The field layer is dominated by dwarf shrub species and very few grasses and herbs. *Calluna* dominates in both subassociations. *Empetrum hermaphroditum* and *Vaccinium* species occur in *cladonietosum*, but in *hylocomietosum* they are constant and co-dominant. Exclusive differential species in the latter subassociation distinguishing from the former are *Linnea borealis*, *Deschampsia flexuosa*, *Cornus suecica*, and *Melampyrum pratense*.

Differences between the two subassociations are distinct in the bottom layers. In *cladonietosum* *Racomitrium lanuginosum* and *Cladonia* species dominate; in *hylocomietosum* they are replaced by *Eu-Piceetum* species.

Most species in *hylocomietosum* also grow in oligotrophic *Eu-Piceetum* forests. Differential species which distinguish these forests are *Calluna vulgaris* and *Empetrum hermaphroditum*.

Cladonietosum occurs on very shallow soil, often with only a thin layer of raw humus covering rocks, and frequently as fragments in *hylocomietosum* vegetation. In *hylocomietosum* the soil is deeper, but even here it only consists of raw humus. On deeper soil with

some mineral content *hylocomietosum* is replaced by *Eu-Piceetum myrtillietosum*. To the west of the distribution area of *Picea abies* forests the vicarious association of *Barbilophozio-Pinetum*, *Bazzanio-Pinetum* normally has a podsol profile or a podsol ranker (Aune 1973).

Kielland-Lund (1967 a) distinguishes between an eastern/northern and a vicarious western oligotrophic humid pine forest type in Norway, *Barbilophozio-Pinetum* and *Bazzanio-Pinetum*, respectively. Four of the differential species of *Bazzanio-Pinetum* (Kielland-Lund 1967 a) occur in my analyses, viz. *Blechnum spicant*, *Rhytidadelphus loreus*, *Barbilophozia barbata*, and *Racomitrium lanuginosum*. The latter is frequent in *cladonietosum*, but the others only occur once. Both differential species of *Barbilophozio-Pinetum* viz. *Barbilophozia lycopodioides* and *Orthocaulis floerkei* are frequent (*hylocomietosum*). *Andromeda polifolia* and *Dicranum majus* are considered to be differential species of *Bazzanio-Pinetum* (Aune 1973). Both species occur, the former in *cladonietosum* and the latter in *hylocomietosum*. The species content indicates a position between *Barbilophozio-Pinetum* and *Bazzanio-Pinetum*. Absence or low frequencies of characteristic *Bazzanio-Pinetum* species such as *Blechnum spicant*, *Sphagnum quinquefarium*, *Barbilophozia barbata*, and *Bazzania trilobata*, is the main reason for including the community in *Barbilophozio-Pinetum*.

The appearance of *cladonietosum* is similar to *Cladonio-Pinetum* K.-Lund 1967. In the opinion of the present author *Cladonio-Pinetum* does not occur in middle or western Central Norway, and *Cladonia* dominated stands (often fragments) occurring here must be included in *Barbilophozio-Pinetum* or *Bazzanio-Pinetum*. Omberg (1980) reports a parallel to *cladonietosum* from SW Norway, "*Racomitrio lanuginosi-Pinetum* prov. ass.".

ASSOCIATION EU-PICEETUM KIELLAND-LUND 1962 (TAB. XIX)

This is the dominating community in the area. The most common substratum is flat or slightly sloping moraine ground. On some of the steeper slopes it is replaced by *Melico-Piceetum*.

The tree layer is dense and homogenous, species other than *Picea abies* have scattered occurrence.

The shrub layer is often absent in dense spruce forests, but small individuals (<1,5 m) of *Sorbus aucuparia* are frequent.

The field layer is poor in species and homogenous (especially in *myrtilletosum*). *Vaccinium myrtillus* in *myrtilletosum* (sometimes also in *dryopteridetosum*) and small ferns in *dryopteridetosum* characterize the physiognomy. Sterile *Deschampsia flexuosa* is always present.

The bottom layer is well developed, totally covering the ground. There are no distinct differences between the two subassociations. *Hylocomium splendens* and *Rhytidadelphus loreus* are alternating dominants, and *Ptilium crista-castrensis* co-dominants.

Constants in the association table are *Picea abies*, *Vaccinium myrtillus*, *Deschampsia flexuosa*, *Dicranum majus*, *Hylocomium splendens*, *Plagiothecium undulatum*, and *Rhytidadelphus loreus*. Differential species distinguishing from *Barbilophozio-Pinetum* are *Gymnocarpium dryopteris*, *Maianthemum bifolium*, *Trientalis europaea*, and *Plagiothecium undulatum*.

Myrtilletosum vegetation grows on the flat moraines. The soil profile is a podsol. The raw humus layer is normally 5-10 cm thick. Below this is a bleached soil layer approximately 5 cm thick. *Myrtilletosum* has no exclusive differential species which distinguish from the two other subassociations in *Eu-Piceetum*.

In *Dryopteridetosum* the soil profile is a semi-podsol. The characteristic layers in a podsol are identifiable, but not distinctly separated. *Dryopteridetosum* vegetation occurs on slightly sloping moraine ground. Exclusive differential species distinguishing from *myrtilletosum* are *Thelypteris phegopteris*, *Oxalis acetosella*, *Anemone nemorosa*, and *Plagiochila major*.

Blechnetosum vegetation only occurs in the higher regions of the area. In the eastern parts the community is common at altitudes above 450 m a.s.l., but occurs scattered at lower altitudes in areas with long lasting snow cover. The tree layer is sparse with low growing *Picea abies* and *Betula pubescens*. Dominance of *Cornus suecica* and *Blechnum spicant* differs from *Myrtilletosum* and *Dryopteridetosum*. The species mentioned are generally more frequent in humid areas. *Cornus suecica* may also be favoured by better light conditions in *blechnetosum*. *Blechnetosum* mostly occurs in sloping terrains and the podsol profile is slightly developed, like that in *dryopteridetosum*.

The variation in vegetation from *myrtilletosum* to *dryopteridetosum* is interpreted as being caused by a moisture gradient. *Blechnetosum* occurs in the most humid areas (cf. station Vennafjell, Fig. 3).

Eu-Piceetum includes the mesotrophic *Vaccinium* and/or fern dominated spruce forests. Some authors also include corresponding western birch/pine forests in *Eu-Piceetum* (Omberg 1980, Rodvelt & Sekse 1980), others establish substituting associations (Kielland-Lund 1971, 1973 "*Corno-Pinetum*", Aune 1973 "*Corno-Betuletum*"). *Eu-Piceetum* and its substitute parallels are usually separated into three subassociations, viz. a poor *Vaccinium* dominated one (*myrtilletosum*), a small fern dominated one (*drypteridetosum*), and a tall fern dominated one (*athyrietosum*) (Kielland-Lund 1971, 1973, Aune 1973, Kjelvik 1978). Aune (1973) also separates a *thelypteridetosum* sub-association. Well defined stands of *athyrietosum* are not found within the investigation area. The few tall fern dominated slopes observed also have some eutrophic species, and this vegetation is included in *Melico-Piceetum*.

Cornus dominated vegetation within *Eu-Piceetum* is described by Omberg (1980) and Rodvelt & Sekse (1980) as a variant of *myrtilletosum*. *Cornus* predominates in *blechnetosum*, but *blechnetosum* is regarded as more eutrophic than *myrtilletosum*. It is floristically well separated from *myrtilletosum*, and is therefore regarded as a subassociation.

ASSOCIATION *MELICO-PICEETUM* KIELLAND-LUND 1962 (TAB. XX)

Melico-Piceetum includes the eutrophic spruce forest on mineral soil. In the area there are three distinct communities belonging to this association, which the present paper separates at subassociation level, viz. *convallarietosum*, *typicum*, and *calliergonelletosum*. The *convallarietosum* analyses are recorded from a steep south-west facing slope with some rocks. Stands with this vegetation occurring scattered adjacent to a scree area with *Ulmo-Tilietum* vegetation. The species content is a mixture of some thermophilous species and *Piceetea* species with different nutritiental demands. The soil layer is thin, consisting of brown aggregated mull and some stones.

The tree and shrub layers are sparse with *Betula pubescens*, *Picea abies*, *Pinus sylvestris*, *Populus tremula*, and *Sorbus aucuparia*.

The field layer is rich in species, although the mean cover value is low. Local character species are *Carex digitata*, *Festuca ovina*, and *Convallaria majalis*.

The moss layer is similar to that of oligotrophic/mesotrophic *Picea* forests. *Hylocomium splendens* is dominant; the only species indicating eutrophic conditions is *Rhytidadelphus triquetrus*.

The ordination diagram (Fig. 9) illustrates the close relationship with the subassociation *typicum*. Differential species distinguishing from the latter are *Corylus avellana*, *Convallaria majalis*, *Dactylis glomerata*, *Festuca ovina*, *Poa nemoralis*, *Galium boreale*, *Solidago virgaurea*, and *Vicia sylvatica*.

The subassociation *typicum* occurs on slopes with eutrophic, soligenous ground water. The soil layer is thick and consists of grey/brown to brown mull. This community often occurs in narrow borders on steep banks along brooks where the *typicum* vegetation gradually changes in direction towards *calliergonelletosum*.

Picea abies has almost optimal conditions in this community. *Alnus incana* is frequent in the tree layer. In the field layer lignous plants are replaced by grasses and herbs. *Vaccinium* species are frequent, but never dominant. Dominants are *Anemone nemorosa*, *Thelypteris phegopteris*, and *Geranium sylvaticum*.

Rhytidadelphus triquetrus is constant and dominant in the bottom layer and *R. loreus* is co-dominant.

Differential species distinguishing against *convallarietosum* are *Crepis paludosa*, *Thelypteris phegopteris*, *Geum rivale*, *Ptilium crista-castrensis*, *Rhytidadelphus loreus*, and *R. calvescens/squarrosum*. The variation in vegetation along the series *convallarietosum-typicum-calliergonelletosum* is interpreted as being caused by a moisture gradient.

Calliergonelletosum stands are only observed as fragments in steep brook ravines. *Picea abies* usually has optimal moisture conditions in this association, but on wet faces (stand A) the optimal water conditions are exceeded.

The field layer is characterized by tall herbs and the absence of *Vaccinium* species. Dominants are *Filipendula ulmaria* and

Lactuca alpina. *Aconitum septentrionale* is observed as a local dominant in fragments of *calliergonelletosum*, but does not occur in the table. The tall herbs mentioned (except for *Filipendula ulmaria*) and the absence of *Vaccinium* species distinguish the field layer from *Melico-Piceetum typicum*.

Brachythecium rivulare, *Calliergonella cuspidata*, and *Cirriphyllum piliferum* are alternating dominants in the bottom layer.

The species content resembles that of the grey alder forest (Fig. 9), and exclusive differential species are few. Further analyses would probably give several more *Piceetea* species. From tables in the present paper only *Caltha palustris*, *Calliergonella cuspidata*, *Mnium pseudopunctatum*, and *Chiloscyphus polyanthus* are differential species distinguishing from *Alno (incana)-Prunetum aconitetosum*. *Caltha palustris* occurs on wet facies in grey alder forests (Klokk 1980).

Kielland-Lund (1971, 1973) includes the eutrophic spruce forests in *Melico (nutantis)-Piceetum*. *Melico-Piceetum* also includes the calciphilous pine forests (subassociation *pinetosum*). Marker (1969) includes the latter community in *Melico-Pinetum*. Bjørndalen (1980 a) describes four geographically substituting associations of Scandinavian calciphilous pine forests. The northern *Epipacto atro-rubentis-Betuletum*, which often occurs in south facing screes, may be considered a substitute of *Ulmo-Tilietum* or other termophilous leaf forests (Bjørndalen 1980 a). The similarity with the "low herb spruce forest" of Kielland-Lund (1967 b) is considerable, but *convallarietosum* is assigned to the northern substitute of calciphilous pine forests mentioned (Bjørndalen 1980 a, 1980 b).

The subassociation *typicum* in the present paper has some hygrophilous species which do not occur in the southern parallel (Kielland-Lund 1967 b, 1971, 1973). The species content in southern descriptions is intermediate between *convallarietosum* and *typicum* in the present paper. The corresponding community in the classification-key of vegetation mapping in Central Norway is "spruce forest rich in grasses and herbs" (Moen et al. 1976).

Kielland-Lund (1971, 1973) and Kjelvik (1978) assigns the eutrophic tall herb spruce forests to *athyrietosum* in *Melico-Piceetum*. Bjørndalen (1978) separates this vegetation from *Melico-Piceetum*, and establishes an association *Aconito-Piceetum* (see also Kielland-Lund 1962). Bjørndalen (1980 a) also suggests a northern

association *Lactuco alpinae-Piceetum*. As mentioned above, the *Alno-Padion* character is considerable in the *calliergonelletosum* (cf. Bjørndalen 1978, 1980 a). The tall herb spruce forest occurs only fragmentarily within the investigation area and the analyses are few, but further investigations will probably indicate the necessity for an association including the tall herb spruce forests in Central Norway. At present the *calliergonelletosum* analyses are assigned to the *Melico-Piceetum* table.

The different kinds of the non thermophilous boreale tall herb vegetation are floristically similar. To test the classification of this vegetation numerical methods were applied on Fennoscandian tall herb vegetation including subalpine birch forests, spruce forests, and grey alder forests.

The RA-diagram in Fig. 5 is based on 20 tall herb communities including 144 randomly selected species of vascular plants. The variation on axis 1 is correlated with an altitudinal gradient, from the subalpine birch forest to the lowland grey alder forest. The diagram indicates the possibility of separating the three types of tall herb vegetation. The same data were run on the TABORD-program with a classification identical to that marked in the ordination diagram. The birch group (*Lactucion alpinae*) and the grey alder group (*Alno-Padion*) had several "characteristic species" at the significance level of 99.9%, but the spruce forest had none. Characteristic species of the birch forest were *Chamaenerion angustifolium*, *Cirsium heterophyllum*, *Luzula multiflora*, *Myosotis decumbens*, *Phleum commutatum*, and *Saussurea alpina*. Characteristic species of the grey alder forest were *Dactylis glomerata*, *Epilobium montanum*, *Geum urbanum*, *Prunus padus*, *Ranunculus repens*, *Roegneria canina*, and *Urtica dioica*. This illustrates the intermediate position of the spruce forest indicated by the variation along the first axis.




Fig. 5. Ordination diagram of 20 communities of tall herb forests from Scandinavia.

1: *Aconito-Piceetum typicum* (Bjørndalen 1978). 2: *Eupiceetum athyrietasum* (Bjørndalen 1978). 3: *Melico-Piceetum athyrietasum* (Kjelvik 1978). 4: *Aconito-Piceetum (Anemone hepatica "type")* (Kielland-Lund 1962). 5: *Aconito-Piceetum (Pyrola rotundifolia "type")* (Kielland-Lund 1962). 6: *Aconito-Piceetum (Athyrium alpestre "type")* (Kielland-Lund 1962). 7: *Poa remota-Aconitum septentrionale ass.* (Kielland-Lund 1962). 8: *Melico-Piceetum calliergonelletosum* (present paper). 9: *Alno (incanae)-Prunetum aconitetosum* (present paper). 10: *Aconitum septentrionale "förbundet"* (Arnborg 1943). 11: *Aconitum-skogar* (Malmström 1949). 12: "Högört - grälskog" (Sjörs 1960). 13: Tall-herb meadow forests (Holmen 1965). 14: "Högstaude björkeskog"-*Aconitum "samf."* (Moen 1978). 15: "Högstaude björkeskog"-*Geranium "samf."* (Moen 1978). 16: "Högstaude björkeskog" *Filipendula-Equisetum "samf"* (Moen 1978). 17: *Betuletum geraniosum subalpinum-Aconitum-Polygon. vertic. var.* (Nordhagen 1943). 18: *Alno (incanae)-Prunetum aconitetosum* (Klok 1980). 19: *Aconito-Alnetum "typisk var."* (Fremstad 1976). 20: *Aconito-Alnetum, Matteuccia "var."* (Fremstad 1976).

ASSOCIATION *ALNO (INCANAE)-PRUNTEUM* KIELLAND-LUND
EX. AUNE 1973 (TAB. XXI)

Grey alder forests in the area grow on clayey soil in ravines. Flat river bank terraces, a common habitat of grey alder forests in Central Norway (Aune 1973, Klokk 1978, 1980), are not observed within the investigated area. Most of the grey alder forests are adjacent to cultivated land and are influenced by grazing. *Alnus incana* is affected favourably by grazing for two reasons, firstly because of the bitter taste (Sjörs 1949), and secondly its resistance to trampling (Sjörs 1960, 1967). The grazing intensity is closely correlated to dominance of *Deschampsia caespitosa* (Steen 1958, Kielland-Lund 1962, Olsson 1975, Klokk 1980). Accordingly stand C in table XXI should be the stand most strongly influenced by grazing. Stand E shows the most common spring aspect in grey alder forests, with total dominance of *Anemone nemorosa*.

The grey alder forests are characterized by a dense tree layer, a luxuriant field layer which is medium rich in species, and a discontinuous moss mat in the bottom layer.

Constant in the table are *Alnus incana*, *Filipendula ulmaria*, *Oxalis acetosella*, *Rubus idaeus*, *Cirriphyllum piliferum*, and *Rhytidiodelphus triquetrus*. The grey alder forest normally differs from eutrophic spruce forests in the absence of *Picea* species such as *Vaccinium myrtillus*, *Deschampsia flexuosa*, *Dicranum majus*, and *Barbilopozia lycopodioides* (cf. Gjerlaug 1973, Kielland-Lund 1973) (notice, however, discussion of *calliergonelletosum* and Fig. 7). The present ravine grey alder forest belongs to the subassociation *aconitetosum* in *Alno (incanae)-Prunetum*. For further sociological discussion of *aconitetosum* see Klokk (1980). From Central Norway *Alnus incana* dominated forests are described by Aune (1973), Fremstad (1979), and Klokk (1980, 1981). For phytosociological discussion of *Alnus incana* hillside forests see also Fremstad & Øvstendal (1978).

ASSOCIATION *ULMO-TILIETUM BOREALE* ASS. PROV. (TAB. XXII)

The analyses are recorded in the only forest stand of *Ulmus glabra* in the area, a steep south/west facing scree slope.

The tree layer is sparse, but several species occur. The quantitatively most important species are *Ulmus glabra* and *Alnus incana*.

The field layer is luxuriant and heterogenous, with *Matteuccia struthiopteris* as the main dominant. *Pteridium aquilinum*, *Rubus idaeus*, and *Stachys sylvatica* are local dominants.

None of the species in the bottom layer are dominant (cf. Fremstad 1979) although some of them are frequent, viz. *Eurhynchium praelongum*, *Rhytidadelphus triquetrus*, *Rhodobryum roseum*, and *Thuidium philibertii*. Litter of *Matteuccia*, bare soil, and rocks predominate the ground.

The constants in the table are also common in non thermophilous grey alder forests, viz. *Circaeа alpina*, *Filipendula ulmaria*, *Matteuccia struthiopteris*, *Oxalis acetosella*, *Rubus idaeus*, and *Stellaria nemoreum*. Differential species distinguishing from *Alno (incanae)-Prunetum* are *Ulmus glabra*, *Galium odoratum*, *Scrophularia nodosa*, and *Viola mirabilis*.

Thermophilous deciduous forests are reported from Central Norway by Aune (1973), Holten (1978), Fremstad (1979), and Steen (1979). The thermophilous deciduous forests are assigned by Aune (1973), Fremstad (1979), and Steen (1979) to *Ulmo-Tilietum* (see also Seibert 1969, Bjørnstad 1971, Kielland-Lund 1971, 1973). Fremstad (1979) establishes a new association *Alno-Ulmetum glabrae* from Orkladalen, regarded as a northern substitute of *Alno-Fraxinetum*. According to tables in Fremstad (1979) *Alno-Ulmetum glabrae* includes thermophilous species such as *Galium odoratum*, *Hypericum hirsutum*, *Lathyrus vernus*, *Moehringia trinerva*, and *Viola mirabilis*. According to Fremstad (1979) most of the thermophilous deciduous forests in Central Norway belong to *Alno-Ulmetum glabrae* and very few of them to *Ulmo-Tilietum*. Aune (1973) prefers to assign his *Ulmus* forest from Hemne to a north-western race of *Ulmo-Tilietum*. Most of the thermophilous species characterizing the rich deciduous forest in Central Norway are at their northern limit of distribution. See floristic descriptions of rich deciduous forest localities in Cen-

tral Norway in Holten (1978). As mentioned by Aune (1973) and Fremstad (1979) further investigations are needed; the most luxuriant forests in Central Norway, in the middle and inner parts of the Trondheimsfjord area, are not yet investigated phytosociologically. Just as it is necessary to establish a boreal association of *Alno-Fraxinetum*, it would also be desirable to establish a northern association of *Ulmo-Tilietum*. Many of the thermophilous species characterizing *Ulmo-Tilietum* in southern Norway (Seibert 1969, Bjørnstad 1971, Kielland-Lund 1971, 1973) do not occur in Central Norway, and the floristic similarity between southern and Central Norwegian *Ulmo-Tilietum* analyses is low (Aune 1973, Fremstad 1979). *Ulmus glabra* and *Corylus avellana* grow in most of the Central Norwegian *Ulmo-Tilietum* localities and should be suitable taxa to identify the northern substitute of *Ulmo-Tilietum*. However, "*Corylo-Ulmetum glabare*" is suggested as the name of an oceanic luxuriant substitute of *Ulmo-Tilietum* (Blom 1980), characterized by some south-western species not occurring in Central Norway. Perhaps this name should be reserved for the northern association, and one of the south-western species used in the name of the western association (see differential species of "*Corylo-Ulmetum glabrae*" against *Ulmo-Tilietum* in Blom (1980)). At present the northern association of *Ulmo-Tilietum* is referred to as *Ulmo-Tilietum boreale*, to which the thermophilous forest in the present paper is assigned.

To test the possibility of maintaining at least two thermophilous *Ulmus* forest associations in Central Norway numerical methods were applied on 62 stand analyses from the papers of Aune (1973), Fremstad (1979), Steen (1979), Klokk (1980), and the present paper. In Fig. 6, 62 stands of *Alnus incana* and *Ulmus glabra* dominated forests from Central Norway are ordinated. If all the available analyses in a table were not used, a random selection was made. The ordination is based on 136 randomly selected species of vascular plants. The variation along axis 1 is interpreted as a temperature gradient from the non thermophilous *Alno (incanae)-Prunetum* to exclusive *Ulmo-Tilietum* communities.

The intermediate thermophilous character of *Alno-Ulmetum glabrae* (Fremstad 1979) is illustrated on the ordination diagram. The weak *Ulmo-Tilietum*-character of *Ulmo-Tilietum boreale* in the present paper, using strict definition of *Ulmo-Tilietum*, is emphasized. The data were also run on TABORD for separation into three




Fig. 6. Ordination diagram of 62 *Alnus incana* and *Alnus incana/Ulmus glabra* stand from Central Norway.

- *Ulmo-Tilietum* (Fremstad 1979)
- * *Ulmo-Tilietum* (Aune 1973)
- ▽ *Ulmo-Tilietum boreale* in the present paper
- △ *Alno-Ulmetum glabrae* (Fremstad 1979 (*typicum* var.))
- ▲ *Alno-Ulmetum glabrae* (Fremstad 1979 (*Matteuccia* var.))
- *Alno (incanae)-Prunetum aconiteto-sum* (Klok 1980)

clusters, and TABORD also identified a gradient corresponding to the first axis in the ordination diagram (Fig. 7 and 8, Tab. VI). The o-marks, outside the clusters represent stands from the residual group in TABORD. In both TABORD runs (cf. Fig. 7 and 8) the index used was "similarity ratio" and the threshold value was 0,3. In conclusion it seems possible to separate at least one less thermophilous association differing from *Ulmo-Tilietum* s.str. in Central Norway.




Fig. 7. The three TABORD clusters marked (quantitative index) in the ordination diagram, showing the correlation with the analyses in the papers mentioned in Fig. 6.




Fig. 8. The distribution of the TABORD-clusters in the ordination diagram. Qualitative index used in the classification.

Tab. VI. Distribution of 62 analyses of *Alnus incana* and *Alnus incana-Ulmus glabra* forests from Central Norway on the three TABORD-clusters marked in Fig. 7.

a: Quantitative index used in the classification.

b: Qualitative index used in the classification.

a		Cluster	Cluster	Cluster
		1	2	3
Ulmo-Tilietum Fremstad 1979				13
Ulmo-Tilietum Aune 1973				5
Ulmo-Tilietum Steen 1979		11		1
"Ulmo-Tilietum boreale"			2	
Alno-Ulmetum glabrae Fremstad 1979	5	7		3
Alno (incanae)-Prunetum aconitetosum Klokk 1980	1		13	
b		Cluster	Cluster	Cluster
		1	2	3
Ulmo-Tilietum Fremstad 1979				13
Ulmo-Tilietum Aune 1973				5
Ulmo-Tilietum Steen 1979	6	1		3
"Ulmo-Tilietum boreale"	2			
Alno-Ulmetum glabrae Fremstad 1979	14			1
Alno (incanae)-Prunetum aconitetosum Klokk 1980	2		10	

GRADIENTS IN THE FOREST VEGETATION

The RA ordination of 42 forest stand analyses (Fig. 9) is based on 132 vascular plants. Mean value of cover degree was calculated for each stand.

The RA diagram mainly confirms the subjective classification of forest communities, most of the communities are well separated in the ordination diagram. The poor pine forests have most of the dominant field layer species in common, and despite some moisture indicators in the damp forest, *Vaccinio uliginosi-Pinetum boreale* and *Barbilophozio-Pinetum* are not distinctly separated along any of




Fig. 9. RA ordination of 42 forest analyses. (EP = *Eu-Piceetum*, MP = *Melico-Piceetum*).

the two first axes. The close floristic relationship between the two tall herb communities *Melico-Piceetum calliergonelletosum* and *Alno (incanae)-Prunetum* is illustrated.

The variation described along the first axis is interpreted as a poor - rich gradient, with the poor pine and spruce forests to the right, the mesotrophic *Eu-Piceetum dryopteridetosum* in the middle, and the rich forests to the left (cf. also Tab. V). It is also interesting to observe the parallel sites between a mineral soil community and the assumed "corresponding" damp forest, viz. *Barbilophozio-Pinetum* and *Vaccinio uliginosi-Pinetum boreale*, *Eu-Piceetum* and *Chamaemoro-Piceetum*, and *Melico-Piceetum* and *Equiseto palustris-Salicetum*.

The variation along axis 2 is correlated with a moisture gradient. The rich damp forest *Equiseto palustris-Salicetum*, also the wettest of the damp forests, as indicated on Fig. 9, is well separated from the other rich forests. The appearance of *Barbilophozio-Pinetum* on the right of axis 1 is, however, in conflict with this interpretation; this is probably due to the large number of common field layer species between the two poor pine forests described. When interpreting the higher axis the general "arch" effect of RA also should be kept in mind (Hill & Gauch 1980).

In conclusion the main floristic variation between the described forest communities may be explained by different nutrient conditions (a poor - rich gradient) and different moisture conditions (a moisture gradient).

SOIL SAMPLES

The pH values from mires show good correlation with the poor - rich gradient (Fig. 10) (cf. Witting 1947, 1948, 1948, Malmer 1962 b). The pH samples are from lawn communities. The highest pH was measured in the *Schoenus ferrugineus-Campylium stellatum-Drepanocladus intermedius* community. Also exchangeable Ca^{++} is positively correlated with the poor - rich gradient (Malmer & Sjörs 1955, Malmer 1962 b). The K^+ and Mg^{++} values show little variation (Tab. IV).




Fig. 10. pH values of mire communities along the poor - rich gradient.

Tab. IV. Exchangeable Ca^{++} , K^+ , and Mg^{++} in upper peat layer (5-15 cm) of mire communities along the poor - rich gradient (g/kg dry weight).

	Ca	K	Mg
Open ombrotrophic bog (lawn)	0.7	0.6	1.9
Open poor fen (lawn)	1.8	0.3	1.0
Open intermediate fen (lawn)	2.5	0.3	0.6
Open extremely rich fen (lawn)	15.0	0.4	0.6

Tab. V. Exchangeable Ca^{++} , K^+ , and Mg^{++} in upper soil layer (5-15 cm) of the forest communities (g/kg dry weight).

	Ca	K	Mg
Chamaemoro-Piceetum	2.9	0.2	1.1
Equiseto palustris-Salicetum	13.9	0.1	0.9
Barbilophozio-Pinetum	1.1	1.3	2.3
Eu-Piceetum myrtillietosum	1.4	1.2	0.9
Eu-Piceetum dryopteridetosum	1.1	1.6	0.9
Melico-Piceetum typicum	4.1	0.6	0.6
Melico-Piceetum calliergonelletosum	6.9	0.1	0.1
Alno (incanae)-Prunetum aconitetosum	3.9	0.6	1.3
Ulmo-Tilietum boreale	7.4	0.4	1.1

Exchangeable Ca^{++} is also positively correlated with the poor - rich gradient in the forest vegetation, corresponding to the first axis on Fig. 6. The K^+ and Mg^{++} values are small and with little variation (Tab. V). Ca^{++} -, K^+ -, and Mg^{++} -values are based on only one sample for a representative stand of each community.

ACKNOWLEDGEMENTS

I wish to express thanks to cand.real. Asbjørn Moen for teaching me field methods at the first stage of this study. Cand. real. Bjørn Sæther has gives advice during the preparation of the manuscript. Mrs. Ruth Waadeland has drawn the figures.

SUMMARY

Phytosociological and ecological studies on mire and forest vegetation were carried out in Klæbu, Central Norway. The main part of the investigation area was between the altitudes 150-400 m a.s.l. The geology is dominated by cambro-silurian sedimented rocks. The climate is suboceanic with a July mean temperature of 14,6-14,7°C and a January mean of -3,1 - -4,2°C. The suboceanic climate is confirmed by the high number of species with a Fennoscandina suboceanic distribution in the area.

The mire vegetation is classified in 36 communities subjectively arranged along the gradients poor - rich, mud bottom - hummock, and partly mire margin - mire expanse. The latter gradient only distinguishes between open mires and tree/shrub covered mires. The boundary between ombratrophic and minerotrophic vegetation is stressed. Four types of minerotrophic mires are separated along the poor - rich gradient, viz. poor, intermediate, rich, and extremely rich fens. Three damp forests, viz. *Vaccinium-Cornus suecica-Hylocomium splendens* community (synonymous with *Vaccinio uliginosi-Pinetum*), *Chamaemoro-Piceetum*, and *Equiseto palustris-Salicetum* are separated along a poor - rich gradient.

The only pine forest community on mineral soil in the area is *Barbilophozio-Pinetum*, which grows on shallow soil, frequently occurring at altitudes above 300 m a.s.l. The dominant community on moraine ground is *Eu-Piceetum*, this is separated into three subassociations, viz. *myrtilletosum* with dominance of *Vaccinium myrtillus*, *dryopteridetosum* with dominance of *V. myrtillus* and small ferns, and *blechnetosum* with co-dominance of *Blechnum spicant* and *Cornus suecica*. *Blechnetosum* is frequent only at altitudes above 450 m.

The eutrophic spruce forests are assigned to the association *Melico-Piceetum*. The three subassociations *convallarietosum*, *typicum*, and *calliergonelletosum* represent a moisture gradient. Wet stands of *calliergonelletosum* gradually turn to rich damp forests. *Convallarietosum* grows on a steep slope adjacent to a *Ulmo-Tilietum boreale* stand. The tall herb dominated *calliergonelletosum* has several species in common with the grey alder forests (cf. Bjørndalen 1978, 1980 a). The grey alder forests in clay ravines are assigned to the subassociation *aconitetosum* in *Alno (incanae)-Prunetum* (see

Klokk 1980). The thermophilous *Ulmus glabra* forest, which only occurs on a south-west facing scree slope, is assigned to *Ulmo-Tilietum boreale* ass. prov.

Numerical methods, including the classification program TABORD (van der Maarel et al. 1978) and the ordination method reciprocal averaging (RA) (Hill 1973) were applied. The numerical treatment of *Ulmus glabra* and *Alnus incana* forests (also including data from other authors) indicates the possibility of separating an exclusive thermophilous community, an intermediate slight thermophilous one, and a non-thermophilous one (Figs. 6, 7, 8). RA was run on 20 Scandinavian communities including different kinds of tall herb vegetation. The main types, i.e. spruce forest, grey alder forest, and birch forest are well separated on the diagram (Fig. 5).

The pH values on the investigated mires are positively correlated with the subjectively arranged communities along the poor - rich gradient. Exchangeable Ca^{++} in the upper soil layer is positively correlated with the poor - rich gradient within the series: mires, damp forests, and forests on mineral soil, with the highest values in extremely rich mires, *Equiseto palustris-Salicetum*, and *Ulmo-Tilietum boreale*.

REFERENCES

- Amann, J. 1929. L'hygrothermie du climat, facteur déterminant la répartition des espèces atlantiques. *Rev. Bryol.* 2: 126-133.
- Arnborg, T. 1943. Granberget. En växtnöologisk undersökning av ett sydlandsk granskogområde med särskilt hänsyn till skogstyper och föryngring. *Norrländsk Handbibl.* 14: 1-282.
- Arnell, S. 1956. *Illustrated moss flora of Fennoscandia I. Hepaticae.* Lund. 308 pp.
- Aune, E.I. 1973. Forest vegetation in Hemne, Sör-Tröndelag, western Central Norway. *K. norske Vidensk. Selsk. Mus. Miscellanea* 12: 1-87.
- Bjørndalen, J.E. 1978. *Aconitum septentrionale* og *Lactuca alpina* som barskogsarter i Skien kommune, Telemark, (*Aconitum septentrionale* and *Lactuca alpina* in the coniferous forests of Skien, Telemark, SE Norway). *Blyttia* 36: 125-134.
- 1980 a. Klassifikasjon av urterike barskoger. Noen kritiske bemerkninger og foreløpige resultater vedrørende noen av skandinavisk plantesosiologis problembarn. In: Baadsvik, K., Klokk, T. & Rønning, O.I. (ed.): Fagmøte i vegetasjonsøkologi på Kongsvoll, 16.-18.3.1980. *K. norske Vidensk. Selsk. Mus. Rapp. Bot. Ser.* 1980-5: 118-126.
- 1980 b. Kalktallskoger i Skandinavien - ett förslag til klassifisering. *Svensk Bot. Tidskr.* 74: 103-122.
- Bjørnstad, A. 1971. A phytosociological investigation of the deciduous forest types in Søgne, Vest-Agder, South Norway. *Norw. J. Bot.* 18: 191-214.
- Blom, H.H. 1980. Plantesosiologiske undersøkelser av edellauvskog og beslektede samfunn på frisk mark i Ytre Hordaland. In: Baadsvik, K., Klokk, T. & Rønning, O.I. (ed.): Fagmøte i vegetasjonsøkologi på Kongsvoll, 16.-18.3.1980. *K. norske Vidensk. Selsk. Mus. Rapp. Bot. Ser.* 1980-5: 134-150.
- Booberg, G. 1930. Gisselåsmyren. En växtsociologisk och utvecklingshistorisk monografi över en jämtlandsk kalkmyr. *Norrländsk Handbibl.* 12: 1-329.
- Braun-Blanquet, J. 1964. *Pflanzensoziologie.* Wien. 865 pp. 3rd. ed.
- Cajander, A.K. 1913. Studien über die More Finlands. *Acta Forest. Fenn.* 2: 1-208.

- Carstens, C.W. 1919. Geologiske undersøkelser i Trondhjems omegn.
Norges Geol. Unders. 83: 1-51.
- Dahl, E. 1950. *Forelesninger over norsk plantogeografi.* Oslo.
114 pp.
- 1967. *Forelesninger i økologi ved Norges Landbrukskole.*
Oslo. 173 pp.
- Dahl, E. & E. Hadac. 1949. Homogeneity of plant communities. *Studia Bot. Cechosl.* 10: 159-176.
- Dahl, E. & H. Krog. 1973. *Macrolichens.* Oslo. 185 pp.
- Du Rietz, G.E. 1949. Huvudenheter och huvudgränser i svensk myr-vegetation. *Svensk Bot. Tidskr.* 43: 274-309.
- Elveland, J. 1976. Myrar på Storöen vid Norrbottenkusten. Coastal mires on the Storöen peninsula, Norrbotten, N. Sweden. *Wahlenbergia* 3: 1-274.
- Eurola, S. 1962. Über die regionale Einteilung der südfinnischen Moore. *Ann. Bot. Soc. 'Vanamo'* 33: 1-243.
- Flatberg, K.I. 1970. *Nordmyra, Trondheim. Aspekter av flora og vegetasjon.* I. Vegetasjonen. Cand.real. Thesis, Trondheim. 183 pp. (unpubl.).
- 1976. Myrundersøkelser i Sogn og Fjordane og Hordaland i forbindelse med den norske myrreservatplanen. *K. norske Vidensk. Selsk. Mus. Rapp. Bot. Ser.* 1976-8: 1-112.
- Flatberg, K.I. & B. Sæther. 1974. Botanisk verneverdigare områder i Trondheimsregionen. *K. norske Vidensk. Selsk. Mus. Rapp. Bot. Ser.* 1974-8: 1-51.
- Fransson, S. 1963. Myrvegetationen vid Rörvatnån i nordvästra Jämtland. *Svensk Bot. Tidskr.* 57: 283-332.
- 1972. Myrvegetation i Sydvästra Värmland. *Acta Phytogeogr. Suec.* 57: 1-135.
- Fremstad, E. 1976. *Vegetasjon og flora i rike løvskoglier i Orkladalen, Sør-Trøndelag.* Cand.real. Thesis. Bergen. 178 pp. (unpubl.).
- 1979. Phytosociological and ecological investigations of rich deciduous forests in Orkladalen, Central Norway. *Norw. J. Bot.* 26: 111-140, 5 Tabs.
- Fremstad, E. & D.O. Øvstedal. 1978. The phytosociology and ecology of grey alder (*Alnus incana*) forests in central Troms, north Norway. *Astarte* 11: 93-112.

- Frisvoll, A.A. & A. Moen. 1981. *Lophozia borealis* sp. nov., a rich fen hepatic from Fennoscandia. *Lindbergia* 6: 137-146.
- Fægri, K. 1960. Maps of distribution of Norwegian plants. I. Coast plants. *Bergen Univ. Skr.* 26: 1-134.
- Gjerlaug, H.C. 1973. Vegetasjonskartlegging av Lillehammer kommune. Cand.real. Thesis. Oslo. 283 pp. (unpubl.).
- Gjærevoll, O. 1972. *Plantegeografi*. Oslo. 186 pp.
- Hill, M.O. 1973. Reciprocal averaging. An eigenvector method of ordination. *J. Ecol.* 61: 237-249.
- Hill, M.O. & H.G. Gauch, Jr. 1980. Detrended correspondance analysis: An improved ordination technique. *Vegetatio* 42: 47-58.
- Holmen, H. 1965. Subalpine tall herb vegetation, site and standing crop. *Acta Phytogeogr. Suec.* 50: 240-248.
- Holten, J. 1978. Verneverdige edellauvskoger i Trøndelag. *K. norske Vidensk. Selsk. Mus. Rapp. Bot. Ser.* 1978-4: 1-199.
- Hultén, E. 1971. *Atlas över växternas utbredning i Norden*. Stockholm. 512 pp.
- Kielland-Lund, J. 1962. *Skogplantesamfunn i Skrukkelia*. Vollebekk. 98 pp.
- 1967 a. Zur Systematik der Kiefernwälder Fennoscandiens. *Mitt. Flor.-Soz. ArbGemein. N.F.* 11/12: 127-141.
 - 1967 b. Lågurtgranskogen og dens erstatningssamfunn på Furuberget. *Medd. Norske Skogforsøksv.* 85: 269-296.
 - 1971. A classification of Scandinavian forest vegetation for mapping purposes (draft). *IBP i Norden* 7: 13-43.
 - 1973. A classification of Scandinavian forest vegetation for mapping purposes. *IBP i Norden* 11: 173-207.
- Kivinen, E. 1935. Über Elektrolythalt und Reaktion der Moorwässer. *Maatalouskeoleitoksen Maatutkimusosaito Agrogeol. Julkaisuja* 38: 1-71.
- Kjelvik, L. 1978. *Barskogsvegetasjon i Øvre Forradalsområdet, Nord-Trøndelag*. Cand.real. Thesis. Trondheim. 131 pp. (unpubl.).
- Klokk, T. 1974. Myrundersøkelser i Trondheimsregionen i forbindelse med den norske myrreservatplanen. *K. norske Vidensk. Selsk. Mus. Rapp. Bot. Ser.* 1974-1: 1-30.
- 1978. *Myricaria*-krattene langs elvene i Trøndelag. (The *Myricaria germanica* thickets along the rivers in Trøndelag, Central Norway.) *Blyttia* 36: 153-161.

- Klokk, T. 1980. River bank vegetation along lower parts of the rivers Gaula, Orkla, and Stjørdalselva, Central Norway. *K. norske Vidensk. Selsk. Skr.* 4: 1-71.
- 1981. Classification and ordination of river bank vegetation from middle and upper parts of the river Gaula, Central Norway. *K. norske Vidensk. Selsk. Mus. Skr.* 2: 1-43.
- Lid, J. 1974. *Norsk og svensk flora*. Oslo. 808 pp. 2nd ed.
- Lye, K.A. 1968. *Moseflora*. Oslo. 140 pp.
- Maarel, E. van der 1979. Multivariate methods in phytosociology, with references to the Netherlands. Pp. 163-225 in: Werger, M.J.A. (ed.): *The study of vegetation*. Junk, The Hague.
- Maarel, E. van der, Janssen, J.G.M. & J.M.W. Louppen. 1978. TABORD, a program for structuring phytosociological tables. *Vegetatio* 38: 143-156.
- Malmer, N. 1962 a. Studies on mire vegetation in the Archaean Area of southwestern Götaland (South Sweden). I. Vegetation and habitat conditions on the Åkhult mire. *Opera Bot.* 7 (1): 1-322.
- 1962 b. Studies on mire vegetation in the Archaean Area of southwestern Götaland (South Sweden). II. Distribution and seasonal variation in elementary constituents on some mire sites. *Opera Bot.* 7 (2): 1-67.
- 1966. *De svenska Sphagnum-arternas systematik och ekologi*. 3rd ed. rev. M. Sonesson. Lund. 46 pp. (mscp).
- 1973. Riktlinjer för en enhetlig klassificering av myrvegetationen i Norden. *IBP i Norden* 11: 154-172.
- Malmer, N. & H. Sjörs. 1955. Some determinations of elementary constituents in mire plants and peat. *Bot. Not.* 109: 46-80.
- Malmström, C. 1949. Studier över skogstyper och trädslagsfördeling inom Västerbottens län. *Medd. Statens Skogf. Inst.* 37: 1-231.
- Marker, E. 1969. A vegetation study of Langøya, Southern Norway. *Nytt Mag. Bot.* 16: 15-44.
- Melin, E. 1917. Studier över de norrländske myrmarkernas vegetation. *Norrländsk Handbibr.* 7: 1-428.
- Moen, A. 1970. *Myr- og kildevegetasjon på Nordmarka - Nordmøre*. Cand.real. Thesis. Trondheim. 245 pp. (unpubl.).

- Moen, A. 1973. Landsplan for myrreservater i Norge. *Norsk Geogr. Tidsskr.* 27: 173-193.
- 1975. Myrundersøkelser i Rogaland. Rapport i forbindelse med den norske myrreservatplanen. *K. norske Vidensk. Selsk. Mus. Rapp. Bot. Ser.* 1975-3: 1-126.
- 1976. Botaniske undersøkelser på Kvikne i Hedmark, med vegetasjonskart over Innerdalen. *K. norske Vidensk. Selsk. Mus. Rapp. Bot. Ser.* 1976-2: 1-100, 1 pl.
- Moen, A. & B.F. Moen. 1975. Vegetasjonskart som hjelpemiddel i arealplanleggingen på Nerskogen, Sør-Trøndelag. *K. norske Vidensk. Selsk. Mus. Rapp. Bot. Ser.* 1975-5: 1-168, 1 pl.
- Moen, A., L. Kjelvik, S. Bretten, S. Sivertsen & B. Sæther. 1976. Vegetasjon og flora i Øvre Forradalsområdet i Nord-Trøndelag, med vegetasjonskart. *K. norske Vidensk. Selsk. Mus. Rapp. Bot. Ser.* 1976-9: 1-135, 1 pl.
- Moen, A. & M. Selnes. 1979. Botaniske undersøkelser på Nord-Fosen, med vegetasjonskart. *K. norske Vidensk. Selsk. Mus. Rapp. Bot. Ser.* 1979-4: 1-96, 1 pl.
- Moen, B.F. 1978. Vegetasjonsøkologiske studier av subalpin skog på Nerskogen, Sør-Trøndelag. Cand.real. Thesis. Trondheim. 149 pp., 6 Tabs. (unpubl.).
- Nordhagen, R. 1928. Die Vegetation und Flora des Sylenegebietes. I. Die Vegetation. *Skr. Norske Vidensk.-Akad. Naturv. Kl.* 1927 (1): 1-612.
- 1943. Sikilsdalen og Norges fjellbeiter. *Bergens Mus. Skr.* 22: 1-607.
- Nyholm, E. 1954-69. *Illustrated moss flora of Fennoscandia. II. Musci.* Lund. 799 pp.
- Olsson, G. 1975. Inverkan av betning och annan skötsel på hagmarkers vegetation. *Svensk Bot. Tidskr.* 69: 393-404.
- Omberg, A. 1980. Noen furuskoger i Ulvik i Hardanger. In: Baadsvik, K., Klokk, T. & Rønning, O.I. (ed.): Fagmøte i vegetasjonsøkologi på Kongsvoll, 16.-18.3.1980. *K. norske Vidensk. Selsk. Mus. Rapp. Bot. Ser.* 1980-5: 92-102.
- Orloci, L. 1978. *Multivariate analyses in vegetation research.* 2nd ed. Junk. The Hague. 451 pp.
- Osvald, H. 1923. Die Vegetation des Hochmoores Komosse. *Sv. Växtsoc. Sällsk. Handl.* 1: 1-436.

- Persson, Å. 1961. Mire and spring vegetation in an area north of Lake Torneträsk, Torne Lappmark, Sweden. I. Description of vegetation. *Opera Bot.* 6 (1): 1-187.
- Persson, S. 1977. Datorprogram för bearbetning av vegetationsdata. I. Klassifikationsprogram - dokumentation och handhavande. *Medd. Avd. Ekol. Bot. Lunds Univ.* 33: 1-68.
- 1978. Datorprogram för bearbetning av vegetationsdata. 2. Ordinationsprogram. Dokumentation och handhavande. *Medd. Avd. Ekol. Bot. Lunds. Univ.* 34: 1-64.
- Poore, M.E.D. 1955. The use of phytosociological methods in ecological investigations. I. Practical issues involved in an attempt to apply the Braun-Blanquet system. *J. Ecol.* 43: 245-269.
- Post, L. von & E. Granlund. 1926. Södra Sveriges torvtillgångar. *Sv. Geol. Unders. Ser. C.* 335: 1-127.
- Rodvelt, O. & L. Sekse. 1980. Eit forslag til sosiologisk inndeling av blåbærdominert vegetasjon i Sør-Noreg. In: Baadsvik, K., Klokk, T. & Rønning, O.I. (ed.): Fagmøte i vegetasjonsøkologi på Kongsvoll, 16.-18.3.1980. *K. norske Vidensk. Selsk. Mus. Rapp. Bot. Ser.* 1980-5: 103-117.
- Ruuhiärvi, R. 1960. Über die regionale Einteilung der nordfinnischen Moore. *Ann. Bot. Soc. 'Vanamo'* 31 (1): 1-360.
- Samuelsson, A. 1960. Skogsvegetationen inom en del av Garpenbergs kronopark. *Svensk Bot. Tidskr.* 54: 69-120.
- Seibert, P. 1969. Über das *Aceri-Fraxinetum* als vikariierenden Gesellschaft des *Galio-Carpinetum* am Rande der Bayerischen Alpen. *Vegetatio* 17: 165-175.
- Sjörs, H. 1946. Myrvegetationen i övre Långanområdet i Jämtland. *Ark. Bot.* 33 (6): 1-96.
- 1948. Myrvegetation i Bergslagen. *Acta Phytogeogr. Suec.* 21: 1-299.
- 1949. Hagar och slotterängar i södra Dalarna. *Natur i Dalarna*: 300-314.
- 1950. Regional studies in North Swedish mire vegetation. *Bot. Not.* 2: 173-222.
- 1952. On the relation between vegetation and electrolytes in North Swedish mire waters. *Oikos* 2: 241-258.

- Sjörs, H. 1960. Kärvaxtfloran och vegetationstyper vid Ångermanälven mellan Nämforsen och Moforsen. *Svensk Bot. Tidskr.* 54: 121-175.
- 1967. *Nordisk växtgeografi*. Stockholm. 244 pp. 2. ed.
- 1971. *Ekologisk botanik*. Stockholm. 296 pp.
- Skogen, A. 1966. Flora og vegetasjon i Ørland herred, Sør-Trøndelag. *K. norske Vidensk. Selsk. Mus. Trondheim Årb.*: 1-124.
- 1969. Trekk av noen oceaniske myrers vegetasjon og utvikling. *Myrers økologi og hydrologi*. IHD 1: 88-95.
- Sonesson, M. 1970. Studies on mire vegetation in the Torneträsk area, Northern Sweden. III. Communities of the poor mires. *Opera Bot.* 26: 1-120.
- Steen, E. 1956. Undersökningar över betingens inflytande i tre naturbeten. *Statens jordbruksförsök Med.* 49: 1-146.
- 1958. Betesinflytelser i svensk vegetation. *Statens jordbruksförsök Medd.* 86: 1-82.
- Steen, O.M. 1979. Verneverdige områder i Melhus kommune, Sør-Trøndelag. Cand.real. Thesis. Trondheim. 131 pp. (unpubl.).
- Størmer, P. 1969. *Mosses with a western and southern distribution in Norway*. Oslo. 288 pp.
- Svensson, G. 1965. Vegetationsundersökningar på Store mosse. *Bot. Not.* 118: 49-86.
- Waldheim, S. & H. Weimark. 1943. Bidrag till Skånes Flora. 18. Skånes myrtyper. *Bot. Not.* 1943: 1-40.
- Whittaker, R. & H.G. Gauch, Jr. 1978. Evaluation of ordination techniques. Pp. 277-336 in: Whittaker, R.H. (ed.): *Ordination of plant communites*. 2nd ed. Junk, The Hague.
- Witting, M. 1947. Kationbestemningar i myrvatten. *Bot. Not.* 4: 287-304.
- 1948. Preliminärt meddelande om fortsatta kationbestämningar i myrvatten sommaren 1947. *Svensk Bot. Tidskr.* 42: 116-135.
- 1949. Kalsiumhalten i några nordsvenska myrvatten. *Svensk Bot. Tidskr.* 43: 715-739.
- Wolff, F.C. 1968. Litt om geologien i Trondheimsområdet. *Trondheim Turterreng*: 51-63.
- 1976. *Geologisk kart over Norge, begrunnslart Trondheim 1:250 000*. Norges geologiske Undersøkelser.

Tab. VII. Open ombrotrophic mire (mud bottom, carpet, lawn).

	Scheuchz. palustr.- commu.	Car.lim.- Clad.flu./ Gymn.inf.- commu.	Car. lim.- Rhync. al.- Drep. schul.- Clad. flu./ Gymno. in.- commu.	Erioph. vag.- Sph. cusp.- S. majus- S. pulchrum- commu.	Sph. magell.- S. pulchrum- commu.	Scirpus caespitosus- S. rubellum- S. tenellum-commu.
Stand						
Relevé No. (0,25 m ²)	1 2 3 4	5 6 7	8 9 10 11	12 13 14 15 16	17 18 19 20 21	22 23 24 25 26 27 28 29 30 31 32 33
Slope in grades	0 0 0	0 0 0	0 0 2 1	0 0 1 1 1	0 3 0 1 2	0 0 0 1 0 0 2 0 3 2 2 0
Aspect	- - -	- - -	- S S	- NW NW NW	- W - S W	- - - S - - S - E E E -
Date	21/7-73	25/7-73	25/7-73	31/8-73	3/8-73	20/6-72 12/7-72 13/7-72
pH	4.3-4.3	4.4-4.5	4.4-4.4 4.5	4.3-4.4	4.4	4.0-4.1 4.2 3.9-4.0
No. of vasc. plants	1 1 1 1	2 3 4	4 3 5 5	2 2 3 2 2	5 3 2 3 2	4 7 6 5 6 3 5 6 5 4 5 6
No. of cryptogams	1 1 1 1	1 1 1	4 4 8 8	4 4 4 3 4	2 5 3 3 2	9 7 5 9 7 6 5 6 6 9 4 6
Tot. num. of species	2 2 2 2	3 4 5	8 7 13 13	6 6 7 5 6	7 8 5 7 5	13 14 11 14 13 9 10 12 11 13 9 12
Andromeda polifolia	.	.	1 1	.	1 1	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
Betula nana
Oxycoccus quadrifetalus	1	.
Pinus sylvestris	1 1
Carex limosa	.	.	1 1 1	1 1 1 1 1	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	.
Eriophorum vaginatum	.	.	.	1 1 .	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	.
Scirpus caespitosus	4 4 4 2 2 2 2 2 1 2 2 2 4	.
Drosera anglica	.	.	1	1 .	.	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
D. rotundifolia	.	.	1	1 .	.	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
Rhynchospora alba	.	.	1 1	2 2 1 1	.	.
Rubus chamaemorus	.	.	.	+	.	.
Scheuchzeria palustris	2 2 2 2	1 .	.	1 1	.	.
Dicranum leioneuron	.	.	.	1	.	1 1 . 1 1
Drepanocladus schulzei	.	.	1 1 1 1	.	1 1 1	.
Sphagnum cuspidatum	6 6 6 6	.	5 5 . 1	3 3 4 4 4	1 .	.
S. fallax
S. fuscum	.	.	1 1	.	.	.
S. lindbergii	.	.	3 3 . 1	.	1	.
S. magellanicum	.	.	.	1 1 1 .	6 6 6 6 6	.
S. papillosum	.	.	.	5 5	.	1 1 1 1 1 4 1 1 1
S. majus	.	.	.	1 1 1 1 1	.	.
S. pulchrum	.	.	.	3 3	6 6 5 5 5	1 1 2 2 2
S. rubellum	.	.	.	1 1	.	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
S. tenuilem	.	.	.	1 .	.	6 6 6 5 5 6 6 5 6 1 6 4
Cephalozia sp(p).	1 1 1 1 1 1 . 1 . 1
Cephaloziella sp.	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
Cladop.fluit./Gymnoc.inflata	.	6 6 6	3 4 2 2	.	.	1 1 1 . 1 1 1 1 1 1 1 1 1 1 1
Lepidozia setacea	1 1 . 1 1 . 1 1 . 1 1 . 1
Mylia anomala	1 . 1
Ptilidium ciliare	1 1 1 1
Cladonia squamosa	1 1
Algae/bare peat	.	2 2 2	2 2

Additional species: Empetrum sp. 23(+), Calliergon stramineum 22(1), Calypogeia sp. 31(1), Cladonia rangiferina 23(+).

Tab. VIII. Open ombrotrophic mire (hummock).

Tab. IX. Tree/shrub-covered embotrophic site.

Stand	Cladonia - commu.												Sph. fuscum-commu.		Vaccinium - Cornus suecica - Hylocomium splendens - commu.													
	S. fusc.- S. nem.- var.						S. fusc.- S. nem.- var.						S. fusc.- S. nem.- var.						S. fusc.- S. nem.- var.									
Relevé No. (1 m ²)	1 2 3	4 5 6	7 8 9	10 11	12 13	14 15	16 17	18 19	20	21 22	23 24	25	26	27 28	29 30	31 32	33 34	35 36	37 38	39 40								
Slope in grades	12 6	3 1 8 4	3 2 4 0	1 16 0	8 6 12	3	6 12 2	8 2 2	15 4	5 6 0	0 2	4	7	8 6 5	8 6 3 4													
Aspect	E SW	E W NW	E SE	- S	N - W W	W W W	W W W	S S SE	EW	E W W W	N N N N	N	W W W W															
Date	15/8-72	10/7-72	13/7-72		15/8-72	10/7-72	27/7-73	21/6-72		10/7-72	15/8-72		14/8-72															
pH	3.7-4.0	3.8-4.2	4.1-4.2		3.6-4.2	3.8-4.2		4.0																				
No. of vasc. plants	7 7 5	8 9 9	12 7 10	8 10	9 11	8	10 10 10	13 13	13 12	12	11 10	11 13	10	10 11	11 8	8	13 13	13 14	11									
No. of cryptogams	6 7 6	12 11 13	10 9 12	11 7	6 11 1	3	4 15	4 5 3	8 6 6	5 10	7 9 12	10 10	4	5 8	5 5	10 11	11 8	9										
Tot. num. of species	13 14 11	20 20 22	22 16 22	19 17	15 19 11	12	15 23	14 15 13	21 19 19	17 22	18 19 23	23 20	14 16 19	13 13	23 24	24 22	20											
A <i>Pinus sylvestris</i>	.	.	.	2	.	.	.	3	.	4	2	3	3	2	3	.	5	2	2	.	5	3	3	.	1	.		
B <i>Picea abies</i>	.	.	.	3	.	.	.	1	.	1	1	1		
<i>Pinus sylvestris</i>	.	.	.	3	.	.	.	1	.	1	1	1		
C <i>Andromeda polifolia</i>	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1			
<i>Betula nana</i>	1 2 3	4 5 6	4 2 3	4 4	3 3	1 3	2	5	3 4	2	3	2	3	3	3	4	1 3 2	1 2	4 5 4	4 3								
<i>Calluna vulgaris</i>	.	1 2	1	2	.	.	.	2	.	2	.	1	1	1	1	1	1	1	1	2	2	3	3	3	3	3		
<i>Empetrum hermaphroditum</i>		
<i>E. nigrum</i>		
<i>E. spp(p).</i>	2 1 2	1	.	2	1	2	3	3	3	2	.	1	1	1	1	1	1	2	2	2	2			
<i>Oxycoccus microcarpus</i>		
<i>O. quadripetalus</i>	.	1	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1			
<i>Picea abies</i>	.	.	.	+	+	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1			
<i>Pinus sylvestris</i>	.	.	.	+	+	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1			
<i>Vaccinium myrtillus</i>	.	1	.	.	.	2	.	2	1	1	1	3	3	2	2	1	2	1	2	2	2	3	2	2	2			
<i>V. uliginosum</i>	2	1	1	.	1	.	1	1	1	2	3	3	2	2	1	1	1	2	1	2	2	2	3	1	2			
<i>V. vitis-idaea</i>	1 1 1	.	.	1	.	1	1	1	2	3	3	2	2	1	1	1	2	2	2	2	2	3	2	2	2			
<i>Carex pauciflora</i>	.	.	1	+	1	1	.	1			
<i>Deschampsia flexuosa</i>	1	1	+	1	1	1	1	1	1	1			
<i>Eriophorum vaginatum</i>	2 1 2	1 1 1	1 1 2	2 1	1 1 1	2	1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1			
<i>Scirpus cespitosus</i>	.	1	1	2	2	2	3		
<i>Cornus suecica</i>	.	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1			
<i>Drosera rotundifolia</i>	.	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1			
<i>Melampyrum pratense</i>		
<i>Rubus chamaemorus</i>	1	1	3	2	3	1	1	1	2	3	2	2	2	3	2	3	3	1	3	4	3	3	3	2	3			
<i>Dicranum leiomorphum</i>	.	1 1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1			
<i>D. majus</i>	1 2 1			
<i>D. scoparium</i>	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1			
<i>D. dilatatum</i>	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1			
<i>Hylocomium splendens</i>	1	2	2	1	1	2	3	1	1	1	1	4	4	5	5	6	5	5	4	3	2	2	1	5	5			
<i>Pleurozium schreberi</i>	6 2	2	1	1	2	1	1	2	3	1	1	1	5	6	4	1	1	1	3	4	1	5	5	1	2			
<i>Polytrichum juniperinum</i>			
<i>var. gracilis</i>			
<i>Ptilium crista-castrensis</i>	1 1 1	1	1 1 2	1	1	1	1	3	5	4	.		
<i>Racomitrium lanuginosum</i>	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	2	2	6	1	1	1	1			
<i>Sphagnum fallax</i>	.	2	2	1	2	1	2	2	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6			
<i>S. fuscum</i>			
<i>S. magellanicum</i>			
<i>S. mooreae</i>			
<i>S. rubellum</i>	.	2	1	1	1	1	1	1	1	1	1	6	4	4	4	2	2	3	3	3	2	1	6	2	1			
<i>S. russowii</i>	1	1	1	1	1	1	1	1	1	1	1	1	1	1			
<i>Barbilophozia lycopodioides</i>	1	1	1	1	1	1	1	1	1	1			
<i>Calypogeia spp(p.)</i>	.	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1			
<i>Cephalozia spp(p.)</i>	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1			
<i>Lepidozia setacea</i>	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1			
<i>Mylia anomala</i>	.	2	2	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1			
<i>Ptilidium ciliare</i>	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1			
<i>Cladonia alpestris</i>	.	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1			
<i>C. rangiferina</i>	2	5	4	1	4	2	1	5	4	3	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1			
<i>C. arbuscula</i>	1	3	5	2	4	1	1	4	3	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1			

Additional species: *Betula pubescens* 38(1), *Calliergon glaucum* 17(1), *Dicranum fuscescens* 2(1), *Dicranum polysetum* 20(1), *Sphagnum compactum* 17(1), *S. gregsonii* 2(1), *S. papillosum* 2(1), *Brabillophozia barbata* 27(1), 30(1), 35(1), 38(1), *Cephalozia spp(p.)* 1(1), 17(1), *Cladopodiella fluitans/Gymnocalyx inflata* 13(1), *Lophozia spp(p.)* 1(1), 6(1), 28(1), *Obtusifolium obtusum* 36(1), *Orthocaulis attenuatus* 3(1), *Orthocaulis cf. binsteadii* 28(1), *Orthocaulis kunzeanus* 40(1), *Cladonia furcata* 36(1).

Tab. X. Open poor fen (mud bottom, carpet)

Tab. XI. Open poor fen (carpet/lawn, lawn).

	Scirpus caesp. - Sph. comp. - Cladop. fluitans/ Gymnac. inflata - comm.												Scirpus caespitosus - Sphagnum tenellum - comm.												Molinia caerulea- Scirp. caespito- Narthecium ossi- comm.									
	Lycopodium inund.-var. var.			Sci. caesp.- Sph. comp.-Cl. fluit./Gymn. inf. var.			Sphagnum papillosum-var.												Sphagnum tenellum-var.												Molinia caerulea- Scirp. caespito- Narthecium ossi- comm.			
Stand	A	B	C	D	E	F	G	H																										
Kelive No. (0,25 m ²)	1 2 3	4 5 6	7 8	9 10 11 12	13 14 15 16	17 18 19 20	21 22 23 24	25 26 27	28 29 30 31 32																									
Slope in grades	1 3 3	4 4 4	4 4	10 13 10 7	9 6 7	6 4 6 3 6	4 4 4 4 4	4 6 6	6 6 6 5 6																									
Aspect	S S S	E E E	E E	NE NE NE NE	NE NE NE NE	NE NE NE NE	N N N N	N NE NE	W W W W																									
Date	7/8/73	1/8/73	14/7/72	17/7/73	19/7/73	24/7/73	17/7/73	21/7/73																										
pH	4.7-5.1	4.0-4.1	4.2-4.5	4.2	-	4.3	4.2	-	4.1-4.5																									
No. of vasc. plants	10 7 8	5 7 9	6 5	12 14 13 15	12 8 6	9 8 7 6	8 5 5 5	5 6 7	5 6 5 6																									
No. of cryptogams	2 4 5	6 4 5	4 4	11 10 8	9 6 7	7 9 6 7	6 7 6 5	4 5 4	0 0 0 0																									
Tot. num. of species	12 11 13	11 11 14	10 9	23 24 21 24	18 15 16	18 14 14 13	14 12 9 10	9 11 11	5 6 5 6																									
Andromeda polifolia	1 1 1	1 1 1	1 1 1	1 1 1 1	1 1 1 1	1 1 1 1	1 1 1 1	1 1 1 1	3 3 2 1 2																									
Betula nana	1	1	1	1	1	1	1	1	1																									
B. pubescens	1	1	1	1	1	1	1	1	1																									
Calluna vulgaris	1	1	1	1	1	1	1	1	1																									
Empetrum nigrum	1	1	1	1	1	1	1	1	1																									
Oxycoccus microcarpus	1	1	1	1	1	1	1	1	1																									
O. quadripetalus	1	1	1	1	1	1	1	1	1																									
Pinus sylvestris	1	1	1	1	1	1	1	1	1																									
Carex limosa	1	1	1	1	1	1	1	1	1																									
C. rostrata	1	1	1	1	1	1	1	1	1																									
Eriophorum angustifolium	1	1	1	1	1	1	1	1	1																									
E. vaginatum	1	1	1	1	1	1	1	1	1																									
Molinia caerulea	1 1 1	3 3 3	3 3 3	3 2 3	2 2 1	4 3 3 3 3	3 4 2	1 1 1	1 1 1																									
Scirpus caespitosus	1 1 1	3 3 3	3 3 3	3 2 3	2 2 1	4 3 3 3 3	3 4 2	1 1 1	1 1 1																									
Drosera anglica	1 1 1	1 1 1	1 1 1	1 1 1	1 2 1	1 1 1 1	1 1 1 1	1 1 1	1 1 1																									
Lycopodium inundatum	5 4 4	4 4 4	4 4 4	3 1 2	1 1	1 1 1 1	1 1 1 1	1 1 1	1 1 1																									
Menyanthes trifoliata	1 2 5	1 1 1	1 1 1	2 2	2	1 1 1 1	1 1 1 1	1 1 1	5 5 5 5 5																									
Narthecium ossifragum	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1																										
Rubus chamaemorus	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1																										
Schuchteria palustris	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1																										
Trientalis europaea	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1																										
Calliergon stramineum	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1																										
Dicranum leibnizii	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1																										
D. schulzei	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1																										
Polytrichum juniperinum	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1																										
S. gracilis	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1																										
Sphagnum compactum	2 5 5	5 5 6	5 5 5	5 5 5	5 5 5	1 1 1 1	1 1 1 1	1 1 1 1																										
S. falax	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1 1																										
S. magellanicum	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1 1																										
S. majus	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1 1																										
S. papillosum	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1 1																										
S. pulchrum	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1 1																										
S. rubellum	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1 1																										
S. subnitens	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1 1																										
S. tenellum	3 4 3	1 4	1 1 1	1 1 1	1 1 1	2 1 1 2	6 6 6 6	6 5 5																										
Calypogeia sp. p.	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1 1																										
Cladop. fluit./Gymnac. inflata	5 5 2	4 4 4	4 4	1 1 1	1 1 1	1 1 1 1	1 1 1 1	2 2 2 2	3 1 1																									
Lepidozia setacea	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1 1																										
Mylia anomala	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1 1																										
Odontoschisma elongatum	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1	1 1 1 1	1 1 1 1	1 1 1 1																										
Ptilidium ciliare	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1 1																										
litter	1	1	1	1	1	1 1 1 1	1 1 1 1	1 1 1 1	5 5 5 5 5																									

Additional species: Vaccinium uliginosum 19(1), Carex lasiocarpa 9(1), Carex pauciflora 10(1), Cornus suecica 13(1), Pinguicula vulgaris 12(1), Potentilla erecta 12(1), Drepanocladus radicans 15(1), Sphagnum fuscum 9(1), Cephalozia sp. 13(1).

Tab. XII. Tree shrub covered poor fen.

	Empetrum - Cornus suecica - Sphagnum fallax - commu.																			Molinia caerulea - Narthecium ossifragum - commu.					
Stand	A					B					C					D					E				
Relieve No. (1 m ²)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
Slope in grades	2	4	3	0	4	6	8	6	4	5	2	3	0	8	3	4	5	6	7	6	2	8	6	8	10
Aspect	E	E	E	-	E	E	E	E	E	NE	NE	-	E	E	E	E	E	E	E	W	W	W	W	W	
Date	5/7-72					10/7-72					7/8-72					7/8-72					9/8-72				
pH	4.4-4.5										4.2-4.6					4.1-4.2					4.0				
No. of vasc. plants	18	17	14	14	18	17	14	11	11	9	16	19	18	17	18	10	11	15	13	9	7	9	10	10	9
No. of cryptogams	10	8	7	9	5	13	12	16	10	7	9	12	9	9	10	10	11	13	16	11	4	1	1	3	2
Tot. num. of species	28	25	21	23	23	30	26	27	21	16	25	31	27	26	28	20	22	28	29	20	11	10	11	13	11
A Pinus sylvestris	1
B Betula pubescens	.	.	.	1
C Andromeda polifolia	1	3	3	3	2	1	1	1	1	1	1	1	1	1	1	2	2	2	1	1	1	1	1	1	1
Betula pubescens	1	1	1	1	.	1	+	+	1	.	1	.	1	.	.	.	1	.	.	.	4	4	5	2	
Calluna vulgaris	1	1	.	1	2	2	1	3	1	3	2	3	1	1	2	.	.	1	1	.	
Empetrum hermaphroditum	.	1	.	1	1	2	2	1	1	2	.	1	2	.	.	1	1	.	
E. sp(p).	2	2	1	1	1	1	1	1	1	1	1	2	2	1	1	2	.	1	2	.	.	1	1	.	
Lycopodium annotinum	1	1	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
Oxycoccus microcarpus	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
O. quadrripetalus	1	1	1	1	1	1	1	1	1	1	2	1	1	1	1	3	2	1	1	1	
Picea abies	1	1	.	+	1	.	.	.	1	1	.	.	1	
Pinus sylvestris	.	1	.	+	1	.	.	.	1	.	1	.	1	.	1	1	.	1	
Vaccinium myrtillus	.	1	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
V. uliginosum	2	1	.	1	1	1	1	1	1	1	1	1	1	1	1	2	2	2	2	3	1	1	2	2	
Carex nigra	1	1	1	1	1	1	1	
C. pauciflora	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
C. rostrata	1	1	1	2	1
Eriophorum angustifolium	2	2	2	1	2	1	1	1	1	1	2	1	2	2	2	.	.	.	1	1	1	1	1	1	
E. vaginatum	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	3	3	1	1	2	.	2	3	2	2
Molinia caerulea	1	1	2	2	2	2	.	1	2	2	.	2	2	2	1	2	
Scirpus caespitosus	2	3	1	1	2	2	2	3	1	3	2	2	1	3	4	4	4	.	2	2	2	1	2	.	
Cornus suecica	1	1	1	.	1	2	1	1	3	3	1	1	2	2	
Dactylorhiza maculata	.	.	.	1	.	1	.	1	.	1	1	1	1	1	1	1	1	1	
Drosera rotundifolia	1	1	1	1	1	2	1	2	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
Narthecium ossifragum	5	2	2	2	2	2	2	1	2	.	1	1	1	1	1	1	
Potentilla erecta	.	.	.	1	2	2	2	2	2	2	1	2	.	1	1	1	1	1	1	
Rubus chamaemorus	3	3	2	1	1	1	1	1	1	1	2	1	1	1	1	2	2	1	1	1	.	1	1	1	
Trientalis europaea	1	.	.	1	1	2	1	1	1	2	1	
D Aulacomnium palustre	1	2	1	1	1	1	
Calliergon stramineum	1	1	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
D. majus	3	2	1	1	
D. undulatum	1	1	1	1	1	1	1	1	1	1	1	1	2	1	1	
Pleurozium schreberi	1	1	1	1	1	1	1	1	1	1	1	2	3	1	1	.	1	1	.	.	
Polytrichum juniperinum	1	.	.	.	3	2	1	.	1	1	4	1	3	
var. gracilius	1	.	.	1	.	.	1	.	
Sphagnum compactum	6	6	6	5	6	1	2	1	1	1	2	1	6	5	2	5	1	1	1	
S. fallax	1	1	.	1	1	1	2	4	1	1	1	2	2	2	1	1	1	6	
S. fuscum	5	2	4	1	1	1	5	4	5	4	4	1	1	1	2	
S. magellanicum	1	1	2	1	1	2	1	2	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
S. nemoreum	1	1	1	1	1	1	1	1	1	1	1	5	2	1	2	
S. papillosum	1	1	1	2	1	1	2	2	6	1	1	1	1	1	1	1	1	1	4	
S. rubellum	1	1	1	1	1	2	2	2	1	1	2	.	1	1	1	1	1	1	2	3	1	.	.	.	
S. russowii	1	.	6	.	.	1	.	1	1	1	1	1	1	1	1	1	.	.	1	
S. temulum	1	.	6	.	.	1	.	1	1	1	1	1	1	1	1	1	.	.	1	
Calypogeia sp(p).	1	1	.	1	.	1	1	1	1	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
Cephalozia sp(p).	.	.	.	1	.	1	1	1	1	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
Cladop. fluit./Gymnoc. inflata	1	1	1	1	1	.	1	.	1	.	1	1	1	1	1	1	1	1	1	1
Lepidozia setacea	.	1	.	.	1	1	1	1	1	.	1	1	1	1	1	1	1	1	3	2	2	1	1	1	
Mylia anomala	1	.	1	.	1	1	2	1	1	.	1	1	1	1	1	1	1	1	3	2	2	1	1	1	
Ptilidium ciliare	1	.	1	.	1	.	1	1	1	.	1	1	1	1	1	1	2	2	

Additional species: Alnus incana B 12(2), Vaccinium vitis-idaea 1(1), 16(1), Carex echinata 13(1), 15(1), Carex panicea 14(1), Nardus stricta 14(1), Melampyrum pratense 16(1), 17(1), Pinguicula vulgaris 7(1), Selaginella selaginoides 12(1), 15(1), 16(1), 18(1), 19(1), Odontoschisma elongatum 8(1), Scapania sp(p). 4(1), 18(1).

Tab. XIII. Open intermediate form

Additional species: *Betula nana* 22(1), 27(1), *Erythronium* sp(p), 40(1), 41(1), *Lycopodium annotinum* 38(1), 39(1), *Vaccinium vitis-idaea* 40(1), *Equisetum palustre* 2(1), *Rubus chamaemorus* 40(1), *Sphagnum compactum* 34(1), *Sphagnum fuscum* 26(2), 41(1), *Sphagnum lindbergii* 40(1), 41(1), *Sphagnum pulchrum* 34(1), 35(1), *Mylia anomala* 30(1).

卷之四

Tab. XV. Estimated yield test.

Tab. XVI. Ass. *Chamaemoro-Piceetum* K.-Lund 1962.

Stand	A					B					C					D					E					
Relevé No. (4 m ²)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	
Slope in grades	14	12	13	12	14	11	6	11	8	6	7	5	5	0	1	9	7	7	3	11	0	2	1	3	1	
Aspect	E	E	E	E	N	NW	NW	W	NW	SE	S	SE	-	SW	W	W	W	W	W	-	E	W	N	E		
Date	16/6-72		4/6-72			11/7-72				14/7-72						14/7-72					10/8-72					
pH	4.0-4.2										4.2						4.2									
No. of vasc. plants	22	26	22	19	18	15	14	14	14	15	11	11	9	9	13	11	13	10	10	11	20	18	18	16	20	
No. of cryptogams	7	10	12	13	13	9	13	15	12	16	12	10	11	14	13	16	14	14	15	10	12	14	12	16	14	
Tot. num. of species	29	36	34	32	31	24	27	29	26	31	23	21	20	23	26	27	27	24	25	21	32	32	30	32	34	
A <i>Betula pubescens</i>	1	1	.	.	.	3	.	5	.	.	2	3	1	
<i>Picea abies</i>	1	.	3	1	1	.	1	1	.	3	2	1	1	1	2	2	1	2	
<i>Pinus sylvestris</i>	2	1	1	.	3	.	.	2	
B <i>Betula pubescens</i>	.	1	1	.	1	1	.	.	.	3	.	5	.	.	2	3	1	
<i>Picea abies</i>	.	1	1	.	1	1
C <i>Betula pubescens</i>	.	1	1	.	1	1	1	2	1	3	1	
<i>Calluna vulgaris</i>	1	.	.	.	1	1	1	.	1	1	1	1	1	1	1	1	1	1	2	1	1	
<i>Empetrum</i> sp(p).	1	1	1	1	1	1	1	1	1	1	2	2	2	1	1	1	1	1	1	1	1	1	1	1	1	
<i>Linnæa borealis</i>	1	1	1	1	1	.	.	.	1	1	1	1	1	
<i>Lycopodium annotinum</i>	.	1	.	.	1	1	1	1	1	1	.	1	.	1	
<i>Picea abies</i>	1	1	1	1	+	.	1	.	1	
<i>Pinus sylvestris</i>	.	.	1	
<i>Sorbus aucuparia</i>	.	1	1	1	1	1	
<i>Vaccinium myrtillus</i>	4	3	3	4	4	5	4	4	4	4	5	6	5	5	5	1	5	4	4	4	3	3	4	3		
<i>V. uliginosum</i>	1	1	1	1	1	1	2	.	2	1	1	2	1	1	1	1	1	1	1	1	1	1	1	1	1	
<i>V. vitis-idaea</i>	1	1	1	1	1	1	1	2	1	2	1	1	1	1	1	1	1	1	1	1	1	1	1	2	1	
<i>Carex echinata</i>	1	1	1	1	1	1	1	.	.	1	.	
<i>C. nigra</i>	.	1	1	1	1	1	1	3	3	1	1		
<i>Deschampsia flexuosa</i>	1	1	1	1	1	1	1	1	1	1	1	1		
<i>Equisetum sylvaticum</i>	1	1	1	1	1	.	.	1	1	1	1	.	.	1	1		
<i>Eriophorum angustifolium</i>	1	1	1	1	1	.	.	1	1	
<i>E. vaginatum</i>	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1		
<i>Luzula pilosa</i>	1	1	1	1	1		
<i>Cornus suecica</i>	1	3	2	3	2	3	2	4	4	4	4	1	1	1	1	2	4	3	2	2	4	4	4	4	3	
<i>Gymnocarpium dryopteris</i>	1	1	1	1	.	.	1	.	1	.	1	2	.		
<i>Listera cordata</i>	1	1	1	1	1	1	.	.	1	1	1	
<i>Maianthemum bifolium</i>	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1		
<i>Melampyrum pratense</i>	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	.	.	.	1	1	1	1	1	1		
<i>Potentilla erecta</i>	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	.	.	.	1	1	1	1	1	1		
<i>Rubus chamaemorus</i>	2	2	1	2	1	5	5	4	5	5	5	5	4	4	5	4	5	4	4	4	3	2	3	4	4	
<i>Tribulus europaeus</i>	1	1	1	.	1	.	1	1	1	1	.	1	1	1	1	1	1	1	1		
D <i>Aulacomnium palustre</i>	1	.	.	.	1	1	1	1	1	.	1	1	1	1	1	1	1	1	1		
<i>Calliergon stramineum</i>	1	1	.	1	.	1	1	1	1	1	1	1	2	1	1	1	1	1	1	.	1	1	1	.		
<i>Dicranum majus</i>	.	1	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	.	1	1	1	.		
<i>D. scoparium</i>		
<i>Hylocomium splendens</i>	.	1	1	1	1	.	1	5	1	1	3	1	1	1	3	1	1	1	1	1	1	2	2	1		
<i>Mnium pseudopunctatum</i>		
<i>Plagiothecium undulatum</i>	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1		
<i>Pleurozium schreberi</i>	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1		
<i>Polytrichum commune</i>	1	2	3	1	1	.	1	1	1	1	.	1	1	1	2	2	4	1	1		
<i>P. juniperinum</i>	.	1	1	.	1	.	1	1	
var. <i>gracile</i>		
<i>Ptilium crista-castrensis</i>	.	1	.	1	1	1	.	1	1	1	1	1	1	1	1	1	1	1	1	1	2	1	2	1		
<i>Rhytidiodelphus loreus</i>	1	1	1	1	1	.	1	1	1	1	1	1	1	1	1	1	2	1	1	1	1	1	1	1		
<i>Sphagnum centrale</i>	.	.	1	3	1	1	5	2	1	1	1	1	1	1	1		
<i>S. fallax</i>	6	5	2	3	6	6	5	1	1	6	3	3	3	3	2	2	2	1	1	2	5	4	5			
<i>S. greggianum</i>	1	1	3	2	2	2	1	1	1	1	4	3	3	2	2	2	3	3	1	2	6	5	3	2		
<i>S. magellanicum</i>	.	.	.	1	2	1	1	2	6	5		
<i>S. nemoreum</i>	.	.	2	1	1	1	1	1	1	.	.	1	3	3	.	2	1	1	1	1	1	1	1	.		
<i>S. russowii</i>	1	1	2	1	1	.	1	1	1	.	1	1	1	1		
Barbilophozia lycopodioides	1	.	1	1	.	1	.	1	.	.	2	.	1	.	.	.		
Calypogeia sp(p).	.	1	1	1	.	1	.	1	1	.	1	.	1	1	1	.	1	1	.	.	1	.	.	.		
Cephalozia sp(p).	.	.	.	1	.	1	1	1	.	1	.	1	1	1	.	1	1		
Obtusifolium obtusum	.	.	1	1	.	1	.	1	.	1	.	1	1	1	1	1	1	1	1	.	1	1	1	1		
Ptilidium ciliare	1	.	.	1	.	.	1	.	1	.	1	1	1	1	1	1	1	1	1		

Additional species: *Andromeda polifolia* 15(1), *Oxycoccus quadrifolatus* 2(1), 17(1), *Agrostis tenuis* 21(1), 23(1), 24(1), 25(1), *Carex pauciflora* 17(1), *Juncus filiformis* 2(1), 3(1), 21(1), 25(1), *Molinia caerulea* 12(1), *Nardus stricta* 1(1), 2(1), 6(1), *Blechnum spicant* 2(1), 3(1), 15(1), 16(1), 17(1), *Dactylorhiza maculata* 1(1), 7(1), 11(1), *Melampyrum* sp(p), 3(1), 4(1), *Dicranum polysetum* 19(1), 22(1), 24(1), 25(1), *Dicranum undulatum* 23(1), *Polytrichum* cf. *formosum* 13(1), *Rhytidiodelphus calvescens*/*squamulosus* 11(1), *Rhytidiodelphus trigetrus* 11(1), *Cephaloziella* sp. 16(1), *Harpanthus flotowianus* 15(1), 21(1), 24(1), 25(1), *Orthocaulis kunzeanus* 21(1), *Pellia* sp. 24(1), *Tritomaria quinqueidentata* 9(1), 10(1), 22(1), 25(1).

Tab. XVII. Ass. *Equisetum palustris-Salicetum* ass. prov.

Stand	A				B				C	D
Relevé No.	1	2	3	4	5	6	7	8	9	10
Area m ²	4	4	4	4	4	4	4	4	2	25
Aspect	-	-	-	-	-	-	-	-	E	N
Slope in grades	0	0	0	0	0	0	0	0	10	4
Date	9/7-73				9/7-73				31/8-78	21/8-78
No. of vascular plants	14	10	10	10	10	12	11	8	14	21
No. of cryptogams	4	6	4	5	8	6	6	5	9	11
Tot. number of species	18	16	14	15	18	18	17	13	23	32
A <i>Alnus incana</i>	2	1	.	2	.	.	1	.	1	.
<i>Betula pubescens</i>	1	2	.	.	.
<i>Picea abies</i>	1	.	1	.	.	1	1	.	.	1
<i>Salix pentandra</i>	2	.	.
B <i>Alnus incana</i>	.	.	2	1
<i>Betula pubescens</i>	1
<i>Picea abies</i>	1
<i>Salix glauca</i>	2	.	1	.	.
<i>S. myrsinifolia</i>	1	2	2
<i>S. pentandra</i>	.	.	.	1	1	1	1	1	.	.
C <i>Calamagrostis purpurea</i>	1	1	.	.	.	1	1	1	.	.
<i>Carex canescens</i>	.	1	3	1
<i>C. echinata</i>	1	1
<i>C. nigra</i>	1	1	1	1	1	2
<i>Deschampsia caespitosa</i>	1	1
<i>Equisetum palustre</i>	1	1	2	1	1	1	1	1	2	3
<i>E. pratense</i>	1	1	1	1	.	.
<i>Juncus articulatus</i>	1	1
<i>J. filiformis</i>	2	2
<i>Anemone nemorosa</i>	1	1
<i>Caltha palustris</i>	2	1	1	1	.	1	.	.	2	1
<i>Comarum palustre</i>	5	5	5	5	1	1	1	2	.	.
<i>Filipendula ulmaria</i>	1	.	1	1	2	2	3	3	.	.
<i>Galium palustre</i>	1	1	1	1	1	.	1	.	2	.
<i>Geum rivale</i>	1	.	.	.	1	1	.	.	.	4
<i>Menyanthes trifoliata</i>	4	4	4	4	5	5	4	4	.	.
D <i>Bryum pseudotriquetrum</i>	1	1	1	1	.	.
<i>Calliergon giganteum</i>	1	3	1	3	.	.	1	.	.	2
<i>Calliergonella cuspidata</i>	2	1	1	1	1	.
<i>Climaciun dendroides</i>	1	.	.	.	1	2
<i>Hylocomium splendens</i>	1	1
<i>Mnium pseudopunctatum</i>	1	1	1	1	3	3	3	3	.	1
<i>M. rugicum</i>	3	1	3	1	1	1	1	1	.	1
<i>M. seligeri</i>	5	5	5	5	5	5	5	5	.	2
<i>Rhytidiodelphus calvescens</i>	2	3
<i>Sphagnum squarrosum</i>	.	1	.	1	.	1
<i>Riccardia pinguis</i>	.	1	1	.	.

Additional species: *Sorbus aucuparia* A 1(1), *Salix aurita* B 4(1), *Salix phyllicolia* 5(1), *Carex vaginata* 10(1), *Luzula multiflora* 10(1), *Poa palustris* 10(1), *Poa trivialis* 10(1), *Cirsium palustre* 9(1), *Galium uliginosum* 10(1), *Myosotis scorpioides* 9 (1), *Polygonum viviparum* 10(1), *Ranunculus acris* 10(1), *Ranunculus repens* 9(1), *Rumex acetosa* 10(1), *Trifolium repens* 10(1), *Tussilago farfara* 9(1), *Atrichum undulatum* 10(1), *Calliergon stramineum* 9(1), *Cirriphyllum piliferum* 9(1), *Drepanocladus uncinatus* 10(1), *Philonotis fontana* 10(2), *Sphagnum girgensohnii* 9(1), *Sphagnum teres* 9(1), *Chiloscyphus polyanthus* 5(1), *Pellia* sp. 5(1), *Cladonia cristata* 10(1).

Tab. XVIII. Ass. *Barbilophozio-Pinetum* Br.-Bl. et Sissing em. K.-Lund 1962.

Stand	cladonietosum												hylocomietosum												
	Racomitrium lanuginosum						Cladonia-var.						Racomitrium lanuginosum						Cladonia-var.						
Relevé No. (4 m ²)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
Slope in grades	13	0	18	0	10	18	20	20	17	20	8	3	7	4	12	10	8	7	17	7	16	12	9	14	12
Aspect	S	-	W	-	W	SW	S	S	S	SE	E	SE	S	S	S	S	E	W	SW	NE	NE	E	E	NE	
Date	10/7-72					16/8-73					10/7-72					11/7-72					14/7-72				
No. of vasc. plants	4	7	4	7	7	5	3	3	3	3	10	9	9	11	10	8	10	10	11	9	9	9	8	12	8
No. of cryptogams	6	5	5	4	6	5	7	6	5	5	6	6	6	8	6	9	10	10	11	7	9	12	9	8	9
Tot. num. of species	10	12	10	11	13	10	10	9	8	8	16	15	15	19	16	17	20	20	22	16	18	21	27	20	17
A <i>Betula pubescens</i>	2	5	2	3	2	.	2	2	2	4	4	2	4	.	1
Pinus sylvestris	4	3	2	5	2	3	2	.	2	2	2	4	4	2	4	.	.
B <i>Picea abies</i>	1	.	.	1	.	1	.	1
Pinus sylvestris	.	.	.	1
C <i>Andromeda polifolia</i>	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	3	4	5	3	3	3	2	3	2
<i>Calluna vulgaris</i>	3	4	3	1	2	4	4	4	4	4	2	1	1	1	1	1	3	4	5	3	3	3	2	3	2
<i>Empetrum hermaphroditum</i>	.	1	.	4	1	1	1	1	1	1	3	5	2	2	3	4	3	5	3	3	4	4	3	3	
E. sp(p).	1	.	1	3
<i>Linnæa borealis</i>	1	1	1	1	1	.	1	
Pinus sylvestris	1	1	.	.	
Vaccinium myrtillus	.	1	.	1	2	1	2	2	2	3	2	3	1	1	3	3	3	4	4
V. uliginosum	.	1	1	1	1	1	1	1	1	1	1	1	2	1	3	2	1	3	2	1	3	4	2	3	
V. vitis-idaea	1	1	1	1	1	3	2	1	1	1	3	4	2	2	2	2	2	3	2	2	1	1	1	1	
Deschampsia flexuosa	1	1	1	1	1	.	1	1	.	1	1	1	1	1	
Cornus suecica	3	2	1	1	2	1	3	2	1	1	1	1	1	1	
Melampyrum pratense	2	2	1	1	1	1	1	1	1	1	1	1	1	1	
Rubus chamaemorus	.	1	1	.	1	
D <i>Dicranum fuscescens</i>	1	1	1	1	1	
D. majus	1	1	1	1	1	.	1	1	1	1	1	1	1	1	
D. scoparium	1	1	1	1	1	1	1	2	1	1	1	1	1	1	1	1	1	1	1	
Hylocomium splendens	1	6	6	6	6	6	6	5	6	5	6	5	3	2	2	
Pleurozium schreberi	.	1	1	.	1	3	3	3	3	1	1	1	1	1	1	1	2	3	2	2	3	2	2	2	
Polytrichum commune	1	1	1	1	1	1	1	2	1	1	1	2	1	1	
Ptilium crista-castrensis	1	1	1	1	1	1	2	1	1	1	1	2	1	1	
Racomitrium lanuginosum	6	6	2	6	4	1	1	1	1	.	1	2	4	5	5
Sphagnum nemoreum	
Barbilophozia lycopodioides	1	.	1	.	1	1	1	.	1	1	1	.	.	.	
Calypogeia sp(p).	1	1	.	.	1	1	
Orthocaulis floerkei	1	1	1	1	1	1	1	1	.	.	1	.	.	
Ptilidium ciliare	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
Cladonia alpestris	3	3	5	2	4	5	5	5	5	5	1	1	1	1	1	1	1	1	
C. rangiferina	1	1	2	1	2	3	4	4	3	3	1	1	1	1	1	1	1	1	1	
C. arbuscula	1	1	1	1	1	2	2	2	1	1	1	

Additional species: *Sorbus aucuparia* C 18(1), 24(1), *Blechnum spicant* 24(1), *Rhytidadelphus loreus* 15(1), 23(1), *Sphagnum russowii* 25(1), *Barbilophozia barbata* 19(1), *Obtusifolium obtusum* 18(1), *Scapania* sp. 22(1), *Tritomaria quinquedentata* 18(1), 25(1), *Hepaticae* sp(p), 19(1), 22(1), *Cladonia furcata* 7(1), *Cladonia uncialis* 2(1).

Tab. XIX. Ass. Eu-Piceetum K.-Lund. 1962

Additional species: *Alnus incana* A 28(2), *Calluna vulgaris* 41(1), 50(1), *Anthoxanthum odoratum* 13(1), 23(1), 36(1), *Nardus stricta* 41(1), 44(1), *Dryopteris filix-mas* 36(2), 38(1), 39(2), *Ceratodon sylvaticum* 28(1), *Hieracium* sp. 24(1), *Orthilia secunda* 34(1), *Solidago virgaurea* 14(1), *Bryocomum umbratum* 36(1), 37(1), 39(1), 40(1), #3(1), *Chlorophloia* sp. 27(1), *Lophocolea* cf. *bidentata* 8(1), 32(1), 36(1), 37(1), 38(1).

Tab. XX. Ass. *Melico-Piceetum* K.-Lund 1962.

	convallarietosum							typicum							calliergonellietosum																															
Stand	Releve No. (4 m ²)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31														
Slope in grades	35	35	44	40	39	42	42	40	15	W	W	W	W	W	E	E	E	E	E	2	0	0	0	2	4	5	4	5	7	20	15	31														
Aspect	W	W	W	W	W	W	W	W	E	E	E	E	E	E	E	E	E	E	E	S	-	-	-	W	E	E	E	E	N	N	N															
Date	22/8/73	10/7/72	20/6/72	5/7/72																	10/7/72	9/7/73	9/7/73																							
No. of vasc. plants	25	22	25	25	24	25	23	21	17	25	22	21	26	26	25	27	34	23	22	17	19	17	17	12	9	11	10	10	14	15	14															
No. of cryptogams	2	10	5	5	5	4	4	13	7	16	8	11	6	9	18	13	6	11	9	9	10	9	9	9	8	10	9	7																		
Tot. num. of species	27	32	30	30	29	30	27	25	30	32	38	29	37	32	34	45	47	29	33	26	28	27	26	21	18	20	18	20	23	22	21															
A	<i>Alnus incana</i>																																													
	<i>Betula pubescens</i>	4	1	3	2	3			2											2	2	2	3		3	2	3	3																		
	<i>Picea abies</i>	2						4	4	2			2	3	5	5	1	1	1	4	2	2		1	1	3																				
	<i>Sorbus aucuparia</i>								2																																					
B	<i>Alnus incana</i>																	1	1		1		1	1	2																					
	<i>Corylus avellana</i>																																													
	<i>Picea abies</i>					1	1																																							
	<i>Sorbus aucuparia</i>					1																																								
C	<i>Alnus incana</i>																	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1												
	<i>Betula pubescens</i>																		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1											
	<i>Lycopodium annotinum</i>																		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1											
	<i>Picea abies</i>																		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1											
	<i>Sorbus aucuparia</i>	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1														
	<i>Vaccinium myrtillus</i>																		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1											
	<i>V. vitis-idaea</i>	1	2	1	2	2	2	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1														
D	<i>Anthonoxanthum odoratum</i>																	1																												
	<i>Carex digitata</i>	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	2	1	1	1	1	1	1	1	1	1	1														
	<i>Dactylis glomerata</i>	1	1	1	1	1	1	1																																						
	<i>Deschampsia caespitosa</i>	1																																												
	<i>Equisetum pratense</i>																																													
	<i>Elymus sibiricus</i>																																													
	<i>Festuca ovina</i>																																													
	<i>Luzula pilosa</i>																																													
	<i>Melica nutans</i>	2	1	2	1	2	1	2	2	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1														
	<i>Poa nemoralis</i>																																													
E	<i>Alchemilla spp.</i>																	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1											
	<i>Anemone nemorosa</i>	1	1	1						1	1	2	3	2	4	4	2	1	2	1	1	2	2	3	3	1	1	1	1	1	2	2	2													
	<i>Athyrium filix-femina</i>																																													
	<i>Blechnum spicant</i>																																													
	<i>Carex palustris</i>																																													
	<i>Cirsium heterophyllum</i>																																													
	<i>Convallaria majalis</i>																																													
	<i>Crepis paludosa</i>																																													
	<i>Chrysosplenium alternifolium</i>																																													
	<i>Dryopteris filix-mas</i>																	2																												
	<i>Gymnocarpium dryopteris</i>	1	1															4	3	1	1		1	1	1	2		2	1	2	2	2	2	1	1											
	<i>Thelypteris phegopteris</i>																	2	4	2	2	1	5	3	3	5																				
F	<i>D. carthusiana</i>																																													
	<i>Filipendula ulmaria</i>																	1	3																											
	<i>Fragaria vesca</i>	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1													
	<i>Gaultheria shallon</i>																																													
	<i>Geum urbanum</i>																																													
	<i>Hieracium acris</i>																																													
	<i>Rubus repens</i>																																													
	<i>R. sativus</i>																																													
	<i>R. triquetus</i>	2	3	3	4	4	5	4	3	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6												
	<i>Rhododendron roseum</i>																	1	1	1	1																									
G	<i>Barbilephoria barbata</i>	1	1	1	1	1	1	1										1																												
	<i>Lycopodioides</i>	1																																												
	<i>Chiloscaphus polyanthus</i>																																													
	<i>Plagiochila cf. amplexoides</i>																																													
	<i>P. major</i>	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	<i>Lophocolea cf. bidentata</i>																	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1								
	<i>Peltaria sp(p.)</i>																																													
	<i>Pellia sp(p.)</i>																																													
	<i>Additional species:</i>	<i>Pinus sylvestris A 3(2), Betula pubescens B 5(1), Salix caprea C 2(2), Salix alba C 1(1), Salix sp. C 10(1), Agrostis tenuis 1(1), Calamagrostis epigejos 17(1), Carex pallescens 16(1), Carex vaginata 17(2), Deschampsia flexuosa 11(1), Equisetum pratense 15(1), Solidago virgaurea 11(1), Festuca ovina 16(1), Malpighia sp(p.) 1(1), Campanula rotundifolia 3(1), Cornus suecica 16(1), Dryopteris affinis 1(1), Lathyrus muralis 7(1), 8(1), Melampyrum sylvaticum 14(1), </i>																																												

Tab. XXI. Ass. *Alno (incanae)-Prunetum* K.-Lund ex Aune 1973.

Stand	A					B					C					D					E				
Relevé No. (4 m ²)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
Slope in grades	24	13	16	20	20	22	26	20	26	32	32	26	21	25	30	18	20	20	22	20	19	18	20	21	20
Aspect	W	W	W	W	W	W	W	W	W	S	S	S	S	S	W	W	W	NW	N	N	N	N	N	N	
Date	13/7-72		13/7-72			20/7-72				10/7-73						8/6-73									
No. of vasc. plants	13	12	12	14	15	13	14	11	10	13	14	13	11	14	10	17	15	15	13	11	12	12	11	12	12
No. of cryptogams	5	5	5	5	5	5	5	5	5	6	5	6	5	5	6	7	7	7	7	4	5	6	6	6	
Tot. num. of species	18	17	17	19	20	18	19	16	15	18	20	18	17	10	15	23	22	22	20	18	16	17	17	18	18
A <i>Alnus incana</i>	5	5	5	5	5	3	5	5	5	5	5	5	5	5	5	5	5	1	2	4	3	4	4	4	4
<i>Picea abies</i>	5	3	.	5	5
<i>Prunus padus</i>	1	.	1
B <i>Alnus incana</i>	1	1	.	2	.	4	3	1	1
<i>Picea abies</i>	1	.	.	1
<i>Prunus padus</i>	1	3	.	.	.	1
<i>Sorbus aucuparia</i>	1
C <i>Alnus incana</i>	1	1	1	.	1	.	1	1	.	1	1	1	1	1	1	1	.	.
<i>Prunus padus</i>	1	1	.	.	1	1	.	1	.	1	.	.	.	1	.	.	1
<i>Sorbus aucuparia</i>	.	.	.	1	1	.	.	1
Deschampsia caespitosa	1	2	1	1	1	1	1	1	1	1	3	2	3	4	2	2	1	1	1	1	.
Aconitum septentrionale	1	3	4	2	4	.	1	1	.	1	.	1	.	1	.	1	1	1	1	1	6	6	6	6	6
Anemone nemorosa	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Circaea alpina	1	1	1	1	1	1	1	1	1	1	2	2	4	2	1	1	1	1	1	1	1
Crepis paludosa	.	1
Chrysosplenium alternifolium	1	1	1	1	.	1	.	1	.	1	.	1	.	1	.	1	1	2	2
Thelypteris phlegopteris	1	1	.
Epilobium montanum	1	.	.	1	1	1	1	1	1	3	3	3	3	3	3	3
Equisetum pratense	3	3	3	3	3	1	1	1	1	1
E. sylvaticum	4	2	2	2	2	1	1	1	1	1	1
Filipendula ulmaria	4	3	4	4	1	4	2	5	4	4	3	3	3	3	3	1	1	1	1	1
Fragaria vesca	2	1	1	1	3	
Geranium sylvaticum	.	.	2	1	1	2	1	1	1	3	1	1	1	1	1	
Geum rivale	1	1	1	1	1	2	2	1	2	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Gymnocarpium dryopteris	1	2	.	.	1	1	1	1	1	1	2	.	.	1	1	1
Lactuca alpina	1	2	.	.	1	1	1	1	1	1	2	5	5	2	2	2
Oxalis acetosella	1	1	1	1	1	2	2	1	2	1	4	4	3	3	4	3	3	3	2	2	1	1	1	1	1
Paris quadrifolia	1	1	1	1	1	1	.	1	.	1	1	1	1	1	1	1
Ranunculus acris	.	.	.	1	1	1	1	1	1	1
R. repens	2	1	2	1	2	1	1	1	1	1	1
Rubus idaeus	2	1	1	2	1	2	2	1	1	2	3	2	1	4	.	5	5	5	5	5	2	1	1	2	1
Stellaria nemorum	.	.	1	1	1	1	1	1	1	1	1	1	1	1	.	2	2	2	2	2
Veronica chamaedrys	1	1	.	1	1	
D <i>Brachythecium reflexum</i>	1	1	1	1	1	1	3	1	1	1	1
B. rivulare	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Cirriphyllum piliferum	5	5	2	5	3	3	6	3	3	1	1	1	1	1	5	5	5	5	5	5	3	3	3	3	3
Euryhydnium praelongum	2	2	1	1	3	2	1	1	3	3	1	1	1	1	1	1	1	1	1	1	1
Mnium affine	1	1	1	1	1	1	1	1	1	1	1
M. undulatum	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Rhytidadelphus calv. /squarros	1	1	1	1	1	3	2	1	1	1
R. loeius	1	1	1	1	1	1	1	1	1	.	.	.
R. triguetrus	2	3	3	2	1	4	1	2	1	1	5	5	4	1	5	.	3	2	2	2	.	2	1	2	3
Rhodobryum roseum	1	1	1	1	1
Plagiochila major	1	1	1	1	1	1	1	1	1	

Additional species: *Betula pubescens* A 16(1), *Picea abies* A 6(5), 8(3), 10(5), B 14(1), *Sorbus aucuparia* B 6(1), C 5(1), 6(1), 11(1), 17(1), *Carex vaginata* 18(1), *Melica nutans* 13(1), *Athyrium filix-femina* 1(1), 14(1), *Dryopteris assimilis* 21(1), 23(1), *Dryopteris filix-mas* 4(1), *Dryopteris carthusiana* 19(1), *Galium aparine* 14(1), 16(1), 17(1), *Geum urbanum* 6(1), *Maianthemum bifolium* 16(1), 17(1), 18(1), *Solidago virgaurea* 16(1), 17(1), *Stachys sylvatica* 5(4), *Taraxacum* sp. 11(1), *Viola palustris* 16(1), 17(1), *Viola riviniana* 11(1), 12(1), 15(1), *Thuidium philibertii* 11(1), *Plagiochila cf. asplenoides* 1(1).

Tab. XXII. "Ulmo-Tilietum boreale" ass. prov..

Stand Relevé No. (4 m ²)	A												B			
	1	2	3	4	5	6	7	8	9	10	11	12	13			
Slope in grades	25	30	35	42	40	40	42	43	40	44	37	38	40			
Aspect	W	W	W	W	W	W	W	W	W	W	W	W	W			
Date							8/6-73							10/7-73		
No. of vasc. plants	13	14	18	14	13	9	12	16	19	14	14	13	15			
No. of cryptogams	6	2	7	6	3	5	5	8	1	1	8	5	5			
Tot. num. of species	19	16	25	20	16	14	17	24	20	15	22	18	20			
A	<i>Alnus incana</i>	.	.	.	5	3	.	.	3	4	.	4	.			
	<i>Prunus padus</i>	.	.	.	4	4	.	.	2			
	<i>Ulmus glabra</i>	4	4	5	5			
B	<i>Alnus incana</i>	1			
	<i>Ulmus glabra</i>	1	.	.	.	1			
C	<i>Dactylis glomerata</i>	.	.	1	1	1	1	.	.	.		
	<i>Deschampsia caespitosa</i>	.	1	3	1	1	1	2	2	1		
	<i>Melica nutans</i>	.	1	1	1	.	.	1			
	<i>Roegneria canina</i>	.	1	2	1	1	1	1	1	1	.	.	1	.		
	<i>Aconitum septentrionale</i>	2	2	.	.	1	1	.	.	.		
	<i>Anemone nemorosa</i>	3	3	3	1	1	1		
	<i>Campanula latifolia</i>	1	2	.	.	.	1	1		
	<i>Circaeа alpina</i>	1	1	2	1	1	1	1	1	1	2	1	1	.		
	<i>Cystopteris fragilis</i>	1	.	1	1	.	1	.		
	<i>Filipendula ulmaria</i>	1	1	1	1	1	2	2	2	.	1	1	1	.		
	<i>Galium odoratum</i>	2	2	2	2	.		
	<i>Geranium robertianum</i>	.	1	1	1	.	.	.			
	<i>G. sylvaticum</i>	.	1	.	1	.	.	1	.	1	.	.	.			
	<i>Matteuccia struthiopteris</i>	5	5	.	5	5	5	5	5	3	.	6	6	6		
	<i>Oxalis acetosella</i>	1	1	1	1	1	1	1	1	1	1	1	1	1		
	<i>Pteridium aquilinum</i>	1	1	2	2	4	5	.	.	.		
	<i>Rubus idaeus</i>	1	1	1	4	4	3	4	3	2	.	1	.	1		
	<i>Stachys sylvatica</i>	.	2	5	5	.	1	1	2		
	<i>Stellaria nemorum</i>	2	.	.	1	1	1	1	1	1	1	3	3	2		
	<i>Urtica dioica</i>	.	.	.	1	1	1	1	1		
	<i>Valeriana sambucifolia</i>	.	.	1	1	.	.	.	1	1	1	.	.	.		
	<i>Viola mirabilis</i>	.	.	1	1	1	.	1	1	1	.	.	.	1		
D	<i>Atrichum undulatum</i>	1	1	1
	<i>Brachythecium reflexum</i>	.	.	1	1	1	.	.
	<i>B. starkei</i>	1	1	1
	<i>Cirriphyllum piliferum</i>	.	.	.	1	.	1	1	1	.	1
	<i>Eurhynchium praelongum</i>	1	1	1	1	.	.	.	1	1	1	.	1	.	1	.
	<i>E. striatum</i>	.	1	.	1	1	.	1	.	1	.
	<i>E. angustirete</i>	.	.	2	1	.	1	.
	<i>Mnium affine</i>	1	1	1	1
	<i>Rhytidadelphus triquetrus</i>	1	.	1	1	1	.	.	1	.	.	1	.	1	.	.
	<i>Rhodobryum roseum</i>	1	.	.	1	1	.	.	1	.	.	1	1	.	.	.
	<i>Thuidium philibertiae</i>	1	.	1	1	1	1	1	1
	rocks/bare soil	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6

Additional species: *Salix caprea* A 11(2), *Sorbus aucuparia* B 3(1),
Carex digitata 3(1), *Chrysosplenium alternifolium*
11(1), 12(1), *Epilobium montanum* 3(1), 13(1),
Fragaria vesca 7(1), 8(1), *Geum urbanum* 11(1),
12(1), *Melandrium rubrum* 1(1), *Paris quadrifolia*
9(1), 10(1), *Scrophularia nodosa* 13(1), *Veronica*
chamaedrys 3(1), *Vicia sylvatica* 9(2), 10(1),
Brachythecium cf. mildeanum 1(1), 11(1), *Hylocomium*
splendens 3(1), 12(1), *Mnium undulatum* 9(1),
Rhytidadelphus loreus 3(1), *Conocephalum conicum*
11(1), *Porella platyphylla* 11(1).