

## REFERENCES

- ALEXANDER, A. J., 1957. The courtship and mating of the scorpion, *Opistophthalmus latimanus*. *Proc. Zool. Soc. Lond.*, **128**: 529–544.
- 1959. Courtship and mating in the buthid scorpions. *Proc. Zool. Soc. London*, **133** (1): 145–169.
- BALINSKY, B. I., 1962. Patterns of animal distribution on the African Continent. *Ann. Cape Prov. Mus.* **2**: 299–309.
- BARNARD, W. S., 1965. 'n Kaart van die Klimaatstreke van Suidwes-Afrika. *J. S.W.A. Sc. Soc.*, **18–19**: 74–84.
- BIRULA, A., 1903. Sur une nouvelle espèce de scorpions provenant de l'île Madagascar. *Ann. Mus. St. Petersb.* **7** (1): x–xi (Russian, with Latin diagnoses of n.g. and sp.).
- CAPORIACCO, L. DI, 1932. Arachnidi. *Boll. Zool., Napoli* **3** (5): 233–238.
- CARTHY, J. D., 1968. The pectines of scorpions. *Symp. Zool. Soc. Lond.* No. **23**: 251–261.
- COATON, W. G. H. & SHEASBY, 1972. Preliminary report on a survey of the termites (Isoptera) of South West Africa. *Cimbebasia Memoir* No. **2**: 1–129.
- COUZIJN, H. W. C., 1976. Functional anatomy of the walking-legs of scorpionida with remarks on terminology and homologization of legs segments. *Neth. J. Zool.* **26** (4): 453–501.
- CROSSKEY, R. W. & WHITE, G. B., 1977. The Afrotropical Region. A recommended term in zoogeography. *J. nat. Hist.*, **1977**, **11**: 541–544.
- DAVIDSON, D. T., 1965. Penetrometer measurements. In C. A. Black (ed.-in-chief). *Methods of soil analysis* Part 1: 472–484. No. 9, Agronomy Series, American Soc. of Agronomy. Inc. Madison, Wisconsin.
- DAVIS, B. J., 1964. Disc electrophoresis—II. Method and application to human serum protein. *Ann. N.Y. Acad. Sci.*, **121**, Art. 2: 404–436.
- FRANCKE, O. F., 1974. Nota sobre los generos *Opistophthalmus* Peters y *Nepabellus* nom. nov. (Scorpionida, Scorpionidae) e informe sobre el hallazgo de *O. lepturus* en la isla del Coco, Costa Rica. *Brenesia* **4**: 31–35.
- GOYFFON, M. & LAMY, J., 1973. Une nouvelle sous-espèce d'*Androctonus australis* L. (Scorpion, Buthidae): *Androctonus australis garzoni* N. SSP. Characteristiques morphologiques, écologiques et biochimiques. *Bull. Soc. Zoo. Fr.* Tome **98**, No. I, 1973: 137–144.
- GOYFFON, M., LE FICHOUX, J. & LAMY, J., 1970. Hétérogénéité du protéinogramme de l'hémolymphe des scorpions *Androctonus australis* L., *Androctonus mauretanicus* Pocock et *Androctonus amoreuxi* Aud. et Sav. en gel de polyacrylamide. *C.R. Séan. Soc. Biol.* **164** (2): 374–378, 2 figs.
- GOYFFON, M., STOCKMANN, R. & LAMY, J., 1973. Valeur taxonomique de l'électrophorèse en disques des protéines de l'hémolymphe chez le Scorpion: étude du genre *Buthotus* (Buthidae). *C.R. Acad. Sc. Paris. T. 277 Serie D*: 61–63.
- HEMPRICH, F. G. & EHRENBURG, C. G., 1828 and 1831. Icones et descriptions corporum naturalium novorum aut minus cognitorum quae ex itineribus per Lybiūm, Egyptūm, Nubīm, Dongolan, Syriūm, Arabīm et Abyssinīm. *Sym. Phys. Arach. Scorp.* No. **10**. Plates published in 1828 and text in 1831. (Original text not seen and page numbers not available in either Sherborn, C. D., 1926. *Index Animalium*, II, or Woodward, cat. Libr. B.M. (N.H.) 515).
- HEMPRICH, F. G., 1829. Symbolae Physicae seu Icones et descriptions Insectorum quae ext itinere per Africam borealem et Asiam occidentalem Animalia articulata, Arachnoidea, Scorpiones. *Berolini ex officina Academica*. 12 pp., 2 pls.
- HEWITT, J., 1913. The Percy Sladen Memorial Expedition to Great Namaqualand 1912–13. Records and descriptions of the Arachnida of the collection. *Ann. Transv. Mus.* **4**: 146–159, pl. 15, figs.
- 1918. A survey of the scorpion fauna of South Africa. *Trans. Roy. Soc. South Africa* **6**: 89–192, pls xix–xxxii.
- 1931. A new Solifuge and Scorpion from S.W.A. *Ann. S. Afr. Mus.* **30** (1): 93–99, 3 text figs.
- 1934. On several Solifuges, Scorpions and a trapdoor spider from S.W.A. *Ann. Transv. Mus.* **15** (3): 401–412, pl. 7, text figs.
- 1935. Scientific results of the Vernay Lang Kalahari Expedition Mar.–Sept. 1930. The Trap-door spiders, Scorpions and Solifuges. *Ann. Transv. Mus.* **16** (3): 459–479, 4 pls. 2 text figs.
- KARSCH, F., 1879a. Scorpilogische Beiträge II. *Mitt. Münchener Ent. Ver.* **3**: 97–136.
- 1879b. West-afrikanische Arachniden, gesammelt von Herrn Stabsarzt Dr Falkenstein. *Zeitschr. Ges. Nat. wiss., Berlin* (3) **IV**: 329–373.
- 1891. Arachniden von Ceylon und von Minikoy. *Berl. Ent. Zeits.* **36** (2): 267–310, 3 pls.
- KELLY, K. L. & DEANE, B. J., 1965. *The ISCC-NBS method of designating colors and a dictionary of color names*. Nat. Bur. Standards circ 553 (reprint ed.): V + 158. ISCC-NBS color-name charts illustrated with centroid colours (suppl. to NBS circ. 553).
- KOCH, C. L. 1837. *Die Arachniden*, Nürnberg **4**: 1–135.
- 1845a. *Die Arachniden*, Nürnberg **11**: 1–174, figs.
- 1845b. *Die Arachniden*, Nürnberg **12**: 1–166, figs.

- KÖPPEN, W., 1931. *Grundriss des Klimakunde*. Berlin.
- KRAEPELIN, K., 1894a. Revision der Skorpione. II. Scorpionidae und Bothriuridae. *Jahrb. Hamburg Wissensch. Anst.* 11: 1-248.
- 1894b. Nachtrag zu Theil I der Revision der Skorpione. *Jahrb. Hamburg Wiss. Anst.* 12: 75-96.
- 1897. Neue und weniger bekannte Skorpione. *Jahrb. Hamb. Wiss. Anst.* 13: 121-146, 1 pl.
- 1898a. Die Skorpione Ost-Afrikas. *Thierw. Ost-Afrika* 4 (5): 1-5 text figs.
- 1898b. Neue Pedipalpen und Skorpione des Hamburger Museums. *Mitt. zool. st. Inst. Hamb.* 15: 1-6, 1 fig.
- 1899. Scorpiones und Pedipalpi. Das Tierreich 8: 1-265, 94 figs.
- 1901. Catalogue des Scorpiones des collections du Museum d'Histoire Naturelle de Paris. *Bull. Mus. Hist. nat. Paris* 7: 265-274.
- 1908. Skorpione und Solifugen Ergebnisse e. Forsch Sudafrika. Bd 1, Lief 2, *Densch. med. Nat. Ges. Jena* 13: 247-282.
- LAMORAL, B. H., 1972. New and Little known Scorpions, Solifuges and Spiders from the Namib Desert, S.W.A. *Madoqua Ser. II*, 1 (54-62): 117-131.
- 1973. The Arachnid fauna of the Kalahari Gemsbok National Park. Part 1. A revision of the species of 'Mole Solifuges' of the genus *Chelypus* Purcell, 1901. (Family Hexisopodidae). *Koedoe* 16: 83-102.
- 1976. *Akentrobuthus leleupi*, a new genus and species of humicolous scorpion from eastern Zaire, representing a new subfamily of the Buthidae. *Ann. Natal Mus.* 22 (3): 681-691.
- 1977. *Parabuthus kalaharicus*, a new species of scorpion from the Kalahari Gemsbok National Park in the Republic of South Africa. (Buthidae, Scorpionida). *Koedoe* 20: 101-107.
- 1978. Soil hardness, an important and limiting factor in burrowing scorpions of the genus *Opistophthalmus* C. L. Koch, 1837. (Scorpionidae, Scorpionida). *Symp. zool. Soc. Lond.* (1978). No. 42: 171-181.
- 1978. Systematics and bionomics of the scorpions of South West Africa (Arachnida, Scorpionida). Ph. D. Thesis, 2 volumes, University of Natal, Pietermaritzburg.
- LAMORAL, B. H. & REYNARDS, S. C., 1975. A catalogue of the scorpions described from the Ethiopian Faunal Region up to December 1973. *Ann. Natal Mus.* 22 (2): 489-576.
- LAWRENCE, R. F., 1927. Contribution to a knowledge of the fauna of S.W.A. V. Arachnida. *Ann. S. Afr. Mus.* 25 (1): 1-75, 4 pls.
- 1928. Contribution to a knowledge of the fauna of S.W.A. VII. Arachnida (Part 2). *Ann. S. Afr. Mus.* 25 (2): 217-312, 4 pls. 1 text fig.
- 1946. A collection of scorpions and Solifugae in the Transvaal Museum with notes on two Natal Solifuges. *Ann Trans. Mus.* 20 (4): 399-408, text-figs 1-3.
- 1949. A collection of Opiliones and Scorpions from North-East Angola made by Dr A. de Barros Machado in 1948. *Pub. Cult. Comp. Diam. Angola*: 1-20, text figs 1-8.
- 1955. Solifugae, Scorpions and Pedipalpi, with checklists and keys to South African families, genera and species. Results of the Lund Univ. Exped. in 1950-51. *S. Afr. Animal Life* 1: 152-262. Figs 1-3.
- 1959. A collection of Arachnida and Myriapoda from the Transvaal Museum. *Ann. Transv. Mus.* 23: 363-386, figs 1-8.
- 1961. New scorpions and solifuges from S.W.A. and Angola. *K. fysiogr. Sallsk. Lund Forh.* 31 (15) 1961: 147-160, figs 1-3.
- 1962. Solifuges, Scorpions and Chilopoda of the Namib Desert. *Ann. Transv. Mus.* 24: 213-222, figs 1-2.
- 1965. New and little known arachnids from the Namib Desert, S.W.A. *Sci Pap. Namib Desert Res. Stn.* 27: 1-12, figs 1-6.
- 1966. New and little known scorpions and Solifuges from the Namib Desert and S.W.A., *Sci. Pap. Namib Desert Res. Stn.* 29: 1-11, 5 figs.
- 1967. Additions to the fauna of S.W.A.: Solifuges, Scorpions and Pedipalpi. *Sci. Pap. Namib Desert Res. Stn.* 34: 1-19, 8 figs.
- 1969. A new genus of Psammophile scorpion and new species of *Opistophthalmus* from the Namib Desert. *Sci. Pap. Namib Desert Res. Stn.* 48: 105-115, 4 figs.
- MAURY, E. A., 1969. *Tityus bahiensis* (Perty 1834) en la Argentina (Scorpiones, Buthidae). *Physis*, 29 (78): 159-164.
- MAURY, E. A. & SAN MARTIN, P. R., 1973. Revalidacion del genero Timogenes Simon 1880 (Scorpiones, Bothriuridae). *Physis*, Sec. C. 32 (84): 129-140.
- MAYR, E., LINSLEY, E. G., & USINGER, R. L., 1953. *Methods and principles of systematic Zoology*. McGraw-Hill Book Company, New York.
- MONARD, A., 1930. Matériaux de la mission scientifique suisse en Angola. Scorpions. *Bull. Soc. neuchâtel Sci. nat.* 54: 37-43, fig.
- 1937. Scorpions, Solifuges et Opilions d'Angola. *Rev. suisse Zool.* 44 (13): 251-270.
- NEWLANDS, G., 1970. A re-examination of some Southern African Scorpion species. *Ann. Transv. Mus.* 26 (10): 199-210, 4 figs.

- NEWLANDS, G., 1972a. A description of *Hadogenes lawrencei* n. sp. (Scorpionidae) with a check list and key to the S.W.A. species of the genus Hadogenes. *Madoqua Ser. II*, 1 (54–62): 133–140.
- 1972b. Notes on Psammophilous scorpions and a description of a new species. *Ann Transv. Mus.* 27 (12): 241–254, figs 1–3, pl. 22.
- 1972c. Ecological adaptations of Kruger National Park Scorpionids (Arachnida: scorponides). *Koedoe* 15: 37–48.
- 1973. Zoogeographical factors involved in the trans-atlantic dispersal pattern of the genus *Opisthacanthus* Peters. *Ann. Transv. Mus.* 28 (7): 91–98.
- PETERS, W., 1861. Ueber eine neue Eintheilung der Skorpione. *Deutsche K. Akademie Wiss. Zu Berlin, Monatsber.* 507–516.
- 1862. Ueber eine neue Eintheilung der Skorpione. *Monber. Ak. Berlin*: 507–516.
- POCOCK, R. I., 1890a. A revision of the genera of scorpions of the family Buthidae with descriptions of some S. African species. *Proc. zool. Soc. Lond.* 1890: 114–141, 2 pls.
- 1890b. Descriptions of two new species of scorpions brought by Emin Pasha from the inland parts of East Africa. *Ann Mag. nat. Hist.* (6) 6: 98–101.
- 1893. Notes on the classification of Scorpions, followed by some observations upon Synonymy with descriptions of new genera and species. *Ann. Mag. nat. Hist.* (6) 12: 303–330, 2 pls.
- 1895. On the Arachnida and Myriapoda obtained by Dr Anderson's collector during Mr T. Bent's Expedition to the Hadramaut, South Arabia. *J. Linn. Soc. (Zool.)* 25: 292–316, pl. IX.
- 1896. On the species of the South African scorpion *Opistophthalmus* contained in the collection of the British Museum. *Ann. Mag. nat. Hist.* (6) 17: 233–248, pl. x.
- 1899. Descriptions of some new species of Scorpions. *Ann. Mag. nat. Hist.* (7) 3: 411–420.
- 1900. Some new Arachnida from Cape Colony. *Ann. Mag. nat. Hist.* (7) 6: 316–333.
- 1901. Descriptions of some new African Arachnida. *Ann. Mag. nat. Hist.* (7) 7: 284–287.
- PURCELL, W. F., 1898. Descriptions of new South African Scorpions in the collection of the South African Museum. *Ann. S. Afr. Mus.* 1 (i): 1–32, pls i–iv.
- 1899a. On the species of *Opistophthalmus* in the collection of the South African Museum, with descriptions of some new forms. *Ann. S. Afr. Mus.* 1: 131–180.
- 1899b. New South African Scorpions in the collection of the South African Museum. *Ann. S. Afr. Mus.* 1: 433–438.
- 1901. On some South African Arachnida belonging to the orders Scorpiones, Pedipalpi and Solifugae. *Ann. S. Afr. Mus.* 2: 137–225, 10 figs.
- ROEWER, C. F., 1943. Über einen neu erworbene Sammlung von Skorpionen des Natur Museums Senckenberg. *Senck. Frankfurt* 26 (4): 205–244, pls 1–6.
- SAN MARTIN, P. R., 1969. Estudio sobre la compleja estructura del esqueleto esclerificado del organo paraxil del genero *Brachiosternus* (Bothriuridae—Scorpionida). *Bol. Soc. Biol. Concepcion* 41: 13–30.
- SCHINZ, H., 1887. Exploration dans le Sud Ouest de l'Afrique. *L'Afrique explorée et civilisée* 6 & 7: 175–188.
- SIMON, E., 1872a. Etudes sur les Scorpions. *Rev. Mag. zool.* (2) 23: 51–59, 97–101, pl. 6.
- 1872b. Arachnides de Syrie, rapportés par M. Charles Piochard de la Brûlerie (Scorpions et Galéodes). *Annals. Soc. ent. Fr.* (5) 2: 247–264.
- 1879. Les Arachnides de France. VII. Contenant les Ordres des Chernetes, Scorpiones et Opiliones. Pages 1–332, pls XVII–XXIV. Paris.
- 1887. Arachnides recueillis dans le sud de l'Afrique par le Dr Hans Schinz. XXVIII. Etudes Arachnologiques, 20<sup>e</sup> Mémoire. *Annls. Soc. ent. Fr.* 7: 369–384.
- STAHNKE, H. L., 1970. Scorpion nomenclature and mensuration. *Ent. News.* 81: 297–316, 10 figs.
- THORELL, T., 1876. On the classification of scorpions. *Ann. Mag. nat. Hist.* (4) 17: 1–15.
- 1877. Etudes Scorpionologiques. *Aiti. Soc. Ital. Sci. Nat.* 19: 75–272.
- TORE-BUENO DE LA, J. R., 1950. *A glossary of Entomology*. Brooklyn, N.Y.: Brooklyn Entomological Society.
- VACHON, M., 1948. Etudes sur les Scorpions. *Arch. Inst. Pasteur Alger.* 26 (1): 25–90, figs 1–86.
- 1948. Etudes sur les Scorpions. Description de scorpions du Nord de l'Afrique. *Ibid.* 26 (2): 162–208, figs. 87–147.
- 1948. *Idem*. *Ibid.* 26 (3): 289–316, figs 148–187.
- 1948. *Idem*. *Ibid.* 26 (4): 441–481, figs 188–241.
- 1949a. *Idem*. *Ibid.* 27 (1): 66–100, figs 242–292.
- 1949b. *Idem*. *Ibid.* 27 (2): 134–169, figs 293–363.
- 1949c. *Idem*. *Ibid.* 27 (3): 281–288, figs 364–371.
- 1949d. *Idem*. *Ibid.* 27 (4): 334–396, figs 372–476.
- 1952a. La réserve naturelle intégrale du Mt Nimba I. Scorpions. Mission M. Lamotte en Guinée (1942). *Mém. F.A.N.*, No. 19. *Rés. Mont Nimba* 1952: 9–15, figs 1–12.
- 1952b. *Etudes sur les Scorpions*: 1–482, figs 1–690. *Inst. Pasteur Algérie*.
- 1955. Remarques préliminaires sur la faune des Scorpions du Soudan oriental. *Bull. Mus. Hist., nat. Paris* (2) 27: 371–373.
- 1961. Scorpions. Le parc national du Niokolo-Koba, Fasc. 2. *Mem. Inst. Franc. Afr. noire* 62: 31–32.

- VACHON, M., 1967. Le grand scorpion du Sénégal: *Pandinus gambiensis*, Pocock 1899 doit être considéré comme une véritable espèce et non comme sous-espèce de *Pandinus imperator* C. L. Koch 1842. *Bull. Inst. fr. Afr. noire* 29A (4): 1534–1537, 11 figs.
- 1969. Complément à la description d'*Isometrus madagassus*, Scorpion Buthidae. *Senckenberg. biol.* 50 (5/6): 417–420.
- 1969. *Groshus griveaudi*, nouvelle espèce de Scorpion Buthidae malgache. *Bull. Mus. Hist. nat. Paris* 41 (2): 476–483.
- 1973. Etude des caractères utilisés pour classer les familles et les genres de Scorpions (Arachnides). *Bull. Mus. Hist. nat. Paris* (3) Zool. 104: 857–958, figs 1–243.
- VACHON, M. & STOCKMANN, R., 1968. Contribution à l'étude des Scorpions africains appartenant au genre *Buthotus* Vachon, 1949 et étude de la variabilité. *Monitore zool. ital. Firenze* 2 suppl.: 81–149, 69 figs.
- WERNER, F., 1902. Die Skorpione, Pedipalpen und Solifugen in der zoologischvergleichend-anatomischen Sammlung der Universität Wien. *Verh. kk. zool-bot. Ges. Wien* 52: 595–608.
- 1916. Über einige Skorpione und Gliederspinne des Naturhistorischen Museums in Wiesbaden. *Jahr. nass. ver. Natk.* 69: 79–97.
- 1933. Die von Dr Fritz Haas auf der Schomburgk-Afrik-Expedition 1931–32 gesammelten Skorpione. *Senck. biol. Frankfurt* 15 (5–6): 323–324, 3 text figs.
- 1936. Neu Eingänge von Skorpionen im Zoologischen Museum in Hamburg. *Festschrift f. Embrifk. Strand Riga* 2: 171–193, 12 text figs.

#### GAZETTEER OF LOCALITY NAMES

The majority of localities listed are in Namibia and the sequence of data is as follows: *locality* (followed by a lot number if a farm), *magisterial district* in brackets, abbreviated *country name* if outside Namibia, *latitude and longitude*. The names and boundaries of the magisterial districts used are those of 1972. Localities outside Namibia are not followed by a magisterial district reference.

#### Key to abbreviations

- A = Angola  
 B = Botswana  
 C = Cape Province, South Africa  
 Mt, Mts = Mountain(s)  
 riv. = river (banks or vicinity)

- Aandster 147, (Maltahöhe), 25°24'S 15°58'E  
 Aar 16, (Lüderitz), 26°46'S 16°28'E  
 Aarpan, C, 27°23'S 20°43'E  
 Agate Beach, (Lüderitz), 26°36'S 15°09'E  
 Aha Mts, Bushmanland, 19°47'S 21°00'E  
 Ai-Ais, (Karasburg), 27°56'S 17°27'E  
 Albrechtshöhe 44, (Karibib), 21°56'S 16°08'E  
 Alte Feste, Windhoek (Windhoek), 22°34'S 17°06'E  
 Ameib 60, (Karibib), 21°49'S 15°40'E  
 Amichab Mts, (Walvis Bay), 23°10'S 15°32'E  
 Andara, Kavango, 18°02'S 21°29'E  
 Anenous, C, 29°17'S 17°34'E  
 Angra Fria, (Skeleton Coast Park), 18°20'S 11°58'E  
 Annabis 677, Damaraland, 20°00'S 14°38'E  
 Annisfontein, Richtersveld, C, 28°25'S 16°53'E  
 Arigap riv., (Karasburg), 28°29'S 18°43'E  
 Aris 29, (Windhoek), 22°43'S 17°07'E  
 Arnhem 222, (Windhoek), 22°43'S 18°06'E  
 Aroab, (Keetmanshoop), 26°48'S 19°39'E  
 Auas Mts, Aris 29, (Windhoek), 22°42'S 17°09'E  
 Augrabies Falls, C, 28°35'S 20°23'E

- Auob and Nossob riv. confluence, C, 26°26'S 20°38'E  
 Auob riv. 11 km N of Twee Rivieren, C, 26°22'S 20°36'E  
 Auob riv. 34 km S of Mata Mata, C, 25°59'S 20°16'E  
 Aus, Etosha Game Park, (Outjo), 19°15'S 16°15'E  
 Aus, (Lüderitz), 26°40'S 16°16'E  
 Auspanplatz (Windhoek), 22°41'S 17°07'E  
 Avis Dam, (Windhoek), 22°35'S 17°07'E  
 Avro 194, (Windhoek), 23°32'S 17°42'E  
 Awasib dunes, (Lüderitz), 25°15'S 15°43'E  
 Bakers Bay, (Lüderitz), 27°42'S 15°32'E  
 Barby 26, (Bethanien), 25°50'S 16°35'E  
 Bayip, C, 25°18'S 20°09'E  
 Beacon 39, Owambo, 17°20'S 17°45'E  
 Belda 361, (Karasburg), 28°27'S 18°01'E  
 Bergkranz 370, (Windhoek), 22°52'S 16°14'E  
 Berseba, Namaland, (Keetmanshoop), 25°59'S 17°47'E  
 Berseba, 10 km S of, Namaland, (Keetmanshoop), 26°07'S 17°46'E  
 Berseba, 10 km W of, Namaland, (Keetmanshoop), 26°02'S 17°42'E  
 Berseba, 48 km W of, Namaland, (Keetmanshoop), 26°07'S 17°27'E  
 Bitterwater, Etosha Game Park, (Outjo), 18°57'S 15°08'E  
 Blässkranz 7, Naukluft Mts, (Maltahöhe), 24°08'S 16°15'E  
 Blinkoog 30, (Karasburg), 27°40'S 19°04'E  
 Bloeddrif, Orange riv. Richtersveld, C, 28°21'S 16°48'E  
 Bloedkopkie, (Swakopmund), 22°49'S 15°24'E  
 Brakwater 48, (Windhoek), 22°23'S 17°04'E  
 Brandberg Basiskamp, Damaraland, 21°08'S 14°33'E  
 Brandberg Mts, Northern Valley, Damaraland, 21°05'S 14°40'E  
 Brandberg Wes Myn, Damaraland, 21°08'S 14°33'E  
 Bruckaros, Namaland, (Keetmanshoop), 25°53'S 17°47'E  
 Cape Cross, (Swakopmund), 21°43'S 13°58'E  
 Cape Cross, 5 km N of, (Swakopmund), 21°43'S 13°56'E  
 Cape Fria, (Skeleton Coast Park), 18°26'S 12°01'E  
 Chamais, (Lüderitz), 27°51'S 15°43'E  
 Chamais Gate, (Lüderitz), 27°52'S 15°44'E  
 Chitapua, A, 14°±15'S 15°±15'E  
 Churutabis 108, (Bethanien), 27°22'S 17°28'E  
 Claratal 18, (Windhoek), 22°50'S 16°51'E  
 Daan Viljoen Game Reserve, (Windhoek), 22°36'S 16°47'E  
 De Jager 279, (Gobabis), 23°13'S 18°44'E  
 De Waal 22, (Gobabis), 23°22'S 18°53'E  
 Djab 26, (Windhoek), 23°16'S 16°12'E  
 Döbra 49, (Windhoek), 22°24'S 17°06'E  
 Dolondolo, A, 13°±45'S 13°±00'E  
 Donkins Bay, C, 31°55'S 18°16'E  
 Dreylingen farm 48, (Bethanien), 26°10'S 17°01'E  
 Düdoabib Ost 57, (Windhoek), 23°13'S 17°39'E  
 Duineveld (Rehoboth), 23°55'S 17°32'E  
 Dunedin Star, Skeleton Coast, 18°13'S 11°56'E  
 Eava 383, (Gobabis), 22°06'S 18°58'E  
 Elandshoek 771, Otavi Highlands, (Tsumeb), 19°26'S 17°42'E  
 Emeritus 123, (Karibib), 23°00'S 15°46'E  
 Epukiro Reserve 268, (Gobabis), 21°43'S 19°05'E  
 Erongo Mts, (Omaruru), 21°40'S 15°40'E  
 Erongo West 83, (Omaruru), 21°40'S 15°50'E  
 Eros 69, (Windhoek), 22°32'S 17°07'E  
 Espinheira, A, 16°46'S 12°24'E  
 Excelsior 286, (Windhoek), 22°27'S 17°38'E  
 Fazenda Valle, Sobreiro, Moçamedes, A, 15°±00'S 12°±15'E  
 Finkenstein 71, (Windhoek), 22°35'S 17°16'E  
 Fish River Canyon, (Karasburg), 27°37'S 17°37'E  
 Fonteine 717, Damaraland, 20°20'S 14°06'E  
 Fransfontein, Damaraland, 20°13'S 15°00'E  
 Frischgewaagd 289, (Windhoek), 22°32'S 17°50'E  
 Gaerdes F. Nature Park, (Okahandja), 21°52'S 17°03'E  
 Gamkarab 176, (Outjo), 20°01'S 16°23'E  
 Ganab, (Walvis Bay), 23°07'S 15°35'E  
 Gasib, Etosha Game Park, (Outjo), 19°19'S 16°14'E  
 Gautscha Pan, Bushmanland, 19°48'S 20°32'E  
 Gemsbokvlakte, Etosha Game Park, (Outjo), 19°13'S 16°07'E

Ghaub 47, Otavi Highlands (Tsumeb), 19°29'S 17°41'E  
 Ghobab 381, (Rehoboth), 23°26'S 17°21'E  
 Goais 13, 8 km S of Helmeringhausen, (Bethanien), 25°56'S 16°57'E  
 Gobabeb, Namib Desert Park, (Walvis Bay), 23°34'S 15°03'E  
 Göllschau 20, (Windhoek), 23°18'S 16°33'E  
 Goodhouse, C, 29°S 18°13'E  
 Goreangab Dam, (Windhoek), 22°32'S 17°02'E  
 Gorob Mine, Namib Desert Park, (Walvis Bay), 23°33'E 15°25'E  
 Gorrasis 99, (Maltahöhe), 25°24'S 15°58'E  
 Grootberg 191, Damaraland, 19°46'S 14°15'E  
 Groot Kolk, C, 24°54'S 20°06'E  
 Grosse Bucht, (Lüderitz), 26°45'S 15°06'E  
 Grünau, (Karasburg), 27°47'S 18°20'E  
 Guibes, or Kuibis (Bethanien), 26°41'S 16°52'E  
 Guruchab, (Keetmanshoop), 26°30'S 18°50'E  
 Halali, Etosha Game Park, (Tsumeb), 19°02'S 16°28'E  
 Hanusa, Namaland, (Mariental), 25°15'S 17°42'E  
 Hardap Dam, (Mariental), 24°30'S 17°52'E  
 Hartmannberge, Kaokoland, 17°30'S 12°23'E  
 Hazeldene, (Outjo), 19°22'S 14°38'E  
 Heide 407, (Rehoboth), 23°31'S 17°22'E  
 Hentiesbaai, (Swakopmund), 22°08'S 14°18'E  
 Heusis 329, (Windhoek), 22°37'S 16°42'E  
 Hoachanas 120, (Mariental), 23°56'S 18°04'E  
 Hoanib riv. mouth, Skeleton Coast Park, 19°27'S 12°44'E  
 Hoas 273, (Outjo), 19°55'S 14°45'E  
 Hoasas 16, (Otiwarongo), 20°20'S 16°36'E  
 Hobas 374, 16 km E of Great Fish riv. canyon, (Karasburg), 27°38'S 17°46'E  
 Hoffnung 66, (Windhoek), 22°27'S 17°11'E  
 Hohenheim 24, (Windhoek), 23°16'S 16°22'E  
 Homob, Etosha Game Park, (Outjo), 19°03'S 16°13'E  
 Hondeklipbaai, C, 30°19'S 17°16'E  
 Hoogland 132, (Maltahöhe), 24°45'S 16°14'E  
 Hottentot Bay, Diamond Area, (Lüderitz), 26°07'S 14°57'E  
 Houmoed, C, 26°22'S 20°31'E  
 Huabmond, Skeleton Coast Park, (Swakopmund), 20°55'S 13°26'E  
 Huams 67, (Maltahöhe), 25°36'S 17°03'E  
 Huns 106, (Bethanien), 27°22'S 17°14'E  
 Iona, A, 16°50'S 12°20'E  
 Isabis 19, (Windhoek), 23°25'S 16°30'E  
 Jakkalsputs, Richtersveld, C, 28°39'S 16°55'E  
 Jansdraai, C, 25°53'S 20°49'E  
 Kakaokroon 487, Damaraland, 20°23'S 14°50'E  
 Kake, (Caprivi Strip), 18°05'S 21°31'E  
 Kam Rivier, (Rehoboth), 24°08'S 16°50'E  
 Kamanjab (Outjo), 19°37'S 14°47'E  
 Kameelsleep, C, 25°50'S 20°44'E  
 Kamkas 20, (Maltahöhe), 24°36'S 16°33'E  
 Kamombonde Oos 86, (Omaruru), 21°34'S 16°03'E  
 Kangas 371, (Rehoboth), 23°36'S 17°03'E  
 Kaoko Otavi, Kaokoland, 18°18'S 13°42'E  
 Kapako, Kavango, 18°01'S 21°23'E  
 Karasburg, (Karasburg), 28°01'S 18°45'E  
 Katima Molilo, (Caprivi Strip), 17°30'S 24°16'E  
 Keetmanshoop, (Keetmanshoop), 26°35'S 18°08'E  
 Keilberg 743, Otavi Highlands, (Grootfontein), 19°34'S 17°40'E  
 Keimasmund 88, (Karasburg), 28°47'S 19°11'E  
 Keres 39, (Windhoek), 22°38'S 16°55'E  
 Khorixas, formerly Welwitchia, Damaraland, 20°22'S 14°56'E  
 Khumib River, Skeleton Coast Park, 18°53'S 12°24'E  
 Kiriis 233, (Keetmanshoop), 26°24'S 19°40'E  
 Klein Aub 350, (Rehoboth), 23°48'S 16°39'E  
 Klein Karasburg, (Keetmanshoop), 27°17'S 18°05'E  
 Klein-Skrypan, C, 26°09'S 20°16'E  
 Klein Windhoek (Windhoek), 22°35'S 17°08'E  
 Koa Valley, ± 21 km SW of Goodhouse, C, 29°04'S 18°06'E  
 Kochena 308, (Keetmanshoop), 27°01'S 18°49'E  
 Koichab Pan, (Lüderitz), 26°18'S 15°36'E  
 Koichab River Bed, (Lüderitz), 26°15'S 16°05'E

- Kolmanskop, (Lüderitz), 26°43'S 15°15'E  
 Kongola, (Caprivi Strip), 17°45'S 23°25'E  
 Kos 28, (Windhoek), 23°16'S 16°08'E  
 Kranzberg 59, (Karibib), 21°58'S 15°39'E  
 Kub, (Mariental), 24°17'S 17°30'E  
 Kuboos, Richtersveld, C, 28°26'S 16°59'E  
 Kubub 15, (Lüderitz), 26°44'S 16°17'E  
 Kuibis, (Bethanien), 26°41'S 16°52'E  
 Kuibis North 168, (Bethanien), 26°28'S 16°48'E  
 Kuiseb Canyon, (Walvis Bay), 23°35'S 15°35'E  
 Kuiseb River Bed, near Gobabeb, (Walvis Bay), 23°35'S 15°03'E  
 Kuyper 666, Damaraland, 20°00'S 14°23'E  
 Kwang, C, 25°20'S 20°30'E  
 Leonardville (Gobabis), 23°31'S 18°50'E  
 Lichtenstein Mitte 366, (Windhoek), 22°48'S 16°58'E  
 Louwshoop 330, (Karasburg), 28°07'S 18°07'E  
 Lucira, A, 13°54'S 12°44'E  
 Lüderitzbucht, (Lüderitz), 26°35'S 15°10'E  
 Makalani 191, Grootberg Mts area, Damaraland, 19°45'S 14°14'E  
 Makambu, Kavango, 17°27'S 18°28'E  
 Makarikari Pans, now Makgadkgadi, B, 20°40'S 25°40'E  
 Maltahöhe, (Maltahöhe), 24°50'S 16°59'E  
 Maluila, A, 15°±30'S 13°±00'E  
 Mara 114, (Bethanien), 27°50'S 17°20'E  
 Marienfluss, Marien riv., Kaokoland, 17°08'S 12°17'E  
 Mariental, (Mariental), 24°38'S 17°58'E  
 Mata Mata, C, 25°53'S 20°01'E  
 Matchless Myn, (Windhoek), 22°42'S 17°19'E  
 Meob Bay Camp, (Walvis Bay), 24°42'S 14°47'E  
 Messum Crater area, Damaraland, 21°16'S 14°13'E  
 Mile 14, (Swakopmund), 22°28'S 14°28'E  
 Moçamedes, A, 15°10'S 12°10'E  
 Molepolole, B, 24°25'S 25°32'E  
 Molteblick or Moltkeblick, (Windhoek), 22°39'S 17°10'E  
 Mooriver, (Bethanien), 27°21'S 17°20'E  
 Moravet, C, 25°28'S 20°30'E  
 Möwebaai, 4 km N of, Skeleton Coast Park, 19°19'S 12°40'E  
 Möwebaai, 8 km NE of, Skeleton Coast Park, 19°17'S 12°42'E  
 Möwebaai, Möwe Point, Skeleton Coast Park, 19°20'S 12°43'E  
 Mukorob 14, Namaland, (Keetmanshoop), 25°32'S 18°09'E  
 Nabas, C, 28°18'S 17°20'E  
 Namuskluft 88, (Lüderitz), 27°53'S 16°50'E  
 Namutoni (Tsumeb), 18°48'S 16°56'E  
 Narib 4, (Maltahöhe), 24°04'S 16°03'E  
 Narudas Süd 268, (Karasburg), 27°22'S 18°51'E  
 Naukluft Mts (Maltahöhe), 24°10'S 16°10'E  
 Naus 27, (Bethanien), 25°48'S 16°30'E  
 Nauzerus 229, (Rehoboth), 23°51'S 16°19'E  
 Neisip 34, (Lüderitz), 26°17'S 16°35'E  
 Neuland 98, (Lüderitz), 26°11'S 16°19'E  
 Nochabeb 97, (Keetmanshoop), 27°24'S 18°30'E  
 Noordoewer, (Karasburg), 28°42'S 17°36'E  
 Nossob Rest Camp, C, 25°25'S 20°37'E  
 Nowantes, Kaokoland, 18°19'S 12°51'E  
 Nubuamis 37, (Windhoek), 22°28'S 17°04'E  
 Numaskloof, Brandberg Mts, Damaraland, 21°07'S 14°24'E  
 Numasplato, Damaraland, 21°08'S 14°30'E  
 Numasvallei, Damaraland, 21°07'S 14°24'E  
 Obib Berge (Lüderitz), 28°05'S 16°45'E  
 Obibwasser, (Lüderitz), 28°00'S 16°40'E  
 Ohopoho, Kaokoland, now called: Opuwa  
 Okahandja, (Okahandja), 21°58'S 16°56'E  
 Okamiparara 335, (Otjiwarongo), 20°25'S 17°28'E  
 Okaukuejo, Etosha Game Park, (Outjo), 19°11'S 15°53'E  
 Okawao, Etosha Game Park, (Outjo), 18°59'S 14°43'E  
 Okondeka, Etosha Game Park, (Outjo), 18°59'S 15°49'E  
 Okonjembo, Kaokoland, now Nowantes  
 Omaruru, (Omaruru), 21°26'S 15°56'E  
 Omaruru riv. mouth, (Swakopmund), 22°05'S 14°14'E

Omatjenguma, Kaokoland, 17°58'S 12°21'E  
 Ombazu, Kaokoland, 17°43'S 13°47'E  
 Ombombo-Ovambo, Kaokoland, 17°59'S 14°15'E  
 Ombu 130, (Omaruru), 21°41'S 15°41'E  
 Onanis 121, (Karibib), 22°48'S 15°42'E  
 Oncocua, A, 16°34'S 13°28'E  
 Ondarusu, near Kunene riv. Kaokoland, 17°25'S 13°55'E  
 Onderste Narries Farm, C, 28°34'S 19°50'E  
 Onguati 236, Damaraland, 19°46'S 14°38'E  
 Onguma 314, (Tsumeb), 18°44'S 17°03'E  
 Onverwag 412, Ugab riv. Bridge, Damaraland, 20°49'S 14°58'E  
 Onze Rust 192, (Mariental), 24°07'S 18°00'E  
 Opwu, formerly Ohopoho, Kaokoland, 18°04'S 13°51'E  
 Orongo, Owambo, 17°40'S 15°17'E  
 Ortmansbaum Farm 120, (Karasburg), 28°19'S 18°43'E  
 Orumana, Kaokoland, 18°15'S 13°34'E  
 Orupembe, also see Anabib waterhole, Kaokoland, 18°11'S 12°31'E  
 Oshakati, Owambo, 17°47'S 15°42'E  
 Otjewi, Kaokoland, 17°29'S 13°56'E  
 Otijjangasemo, Kaokoland, 17°27'S 13°15'E  
 Otjikoko-Süd 67, (Omaruru), 21°19'S 16°23'E  
 Otjikoto, (Tsumeb), 19°11'S 17°33'E  
 Otjinungwa, Kaokoland, 17°55'S 12°20'E  
 Otjitambi 25, (Outjo), 19°49'S 15°10'E  
 Otjitanda, Kaokoland, 17°38'S 12°46'E  
 Otjitundua, Kaokoland, 18°39'S 14°14'E  
 Otjituo, Hereroland, 19°40'S 18°36'E  
 Otjombae, Damaraland, 21°23'S 15°10'E  
 Outjo (Outjo), 20°07'S 16°09'E  
 Owambo Grens, Owambo, 18°30'S 16°45'E  
 Palmfontein farm, Grootberg Mts area, Damaraland, 19°40'S 14°14'E  
 Pastoril do Sul, Moçamedes, A, 15°±15'S 12°±30'E  
 Paulinenhof 72, (Windhoek), 22°40'S 17°17'E  
 Pella Mission, C, 28°59'S 19°10'E  
 Perdepan 516, (Gobabis), 23°40'S 19°00'E  
 Plaas 631, (Outjo), 19°23'S 14°44'E  
 Plateau 38, (Lüderitz), 26°40'S 16°30'E  
 Pofadder, C, 29°08'S 19°23'E  
 Pollentswa Riv. 5 miles from Nossob riv., B, 25°05'S 20°26'E  
 Popa Falls, Kavango, 18°07'S 21°33'E  
 Port Nolloth, C, 29°15'S 16°52'E  
 Porto Alexandre, A, 15°50'S 11°51'E  
 Portsmut 33, Hakos Mts, (Windhoek), 23°11'S 16°24'E  
 Praia Madalena, Moçamedes, A, 14°±30'S 12°±15'E  
 Purros, Kaokoland, 18°47'S 12°55'E  
 Quibis, see Kuibis  
 Ramboekas Pan, B, 24°47'S 20°15'E  
 Regenstein 32, (Windhoek), 22°44'S 17°00'E  
 Rehoboth, (Rehoboth), 23°19'S 17°05'E  
 Rehoboth, 5 km W of, (Rehoboth), 23°23'S 17°00'E  
 Remhoogte 227, (Maltahöhe), 23°58'S 16°17'E  
 Richthofen 126, (Windhoek), 22°34'S 17°45'E  
 Richtersveld, C, 28°15'S 17°10'E  
 Rietfontein, Hereroland, 21°54'S 20°55'E  
 Rocky point, Skeleton Coast Park, 19°01'S 12°29'E  
 Rooibank, (Walvis Bay), 23°12'S 14°41'E  
 Rooiberg 70, (Lüderitz), 27°09'S 16°47'E  
 Rooibrak, C, 26°03'S 20°27'E  
 Rookop, Namib Desert Park, (Walvis Bay), 22°59'S 14°40'E  
 Rosh Pinah, (Lüderitz), 27°58'S 16°44'E  
 Rössing Mts (Swakopmund), 22°30'S 14°36'E  
 Rostock 393, (Windhoek), 23°22'S 15°55'E  
 Ruacana, Owambo, 17°23'S 14°15'E  
 Rudenau Nord 6, (Okahandja), 22°05'S 16°42'E  
 Rundu, Kavango, 17°55'S 19°43'E  
 Sa da Bandeira, Huila, A, 14°54'S 13°35'E  
 Saddle Hill, (Lüderitz), 25°54'S 14°55'E  
 Saffier 148, (Maltahöhe), 25°27'S 16°06'E  
 Samahaling 277, (Keetmanshoop), 27°02'S 19°35'E

- Samangeigei, Kavango, 19°06'S 20°12'E  
 Sandamap 64, Damaraland, 21°56'S 15°16'E  
 Sandmund 270, (Karasburg), 27°17'S 18°57'E  
 Sangwali, (Caprivi Strip), 18°17'S 23°37'E  
 Sanitatas, Kaokoland, 18°17'S 12°40'E  
 Sarusas, Skeleton Coast Park, 18°50'S 12°30'E  
 Scheidthof 293, (Windhoek), 22°42'S 18°03'E  
 Schwarzkuppen 104, (Keetmanshoop), 27°23'S 18°20'E  
 Sesfontein, Kaokoland, 19°07'S 13°36'E  
 Sesriem 137, (Malta Höhe), 24°33'S 15°50'E  
 Seven Pans, C, 25°26'S 20°22'E  
 Simon De Wit Bridge, (Grootfontein), 19°34'S 18°07'E  
 Sissekab 54, (Grootfontein), 19°19'S 17°12'E  
 Skaap River, (Keetmanshoop), 26°37'S 18°07'E  
 Smithfield Beach, (Lüderitz), 26°35'S 15°09'E  
 Spencer Bay, (Lüderitz), 25°43'S 14°50'E  
 Spencer Bay Water, (Lüderitz), 25°47'S 14°54'E  
 Spitzkoppe Gross —, Farm 71 and Mts, Damaraland, 21°50'S 15°10'E  
 Spitzkoppe Klein —, Damaraland, 21°50'S 15°05'E  
 Springbok, C, 29°40'S 17°53'E  
 Springbokvlakte, C, 28°23'S 17°04'E  
 Sterkstroom 320, (Keetmanshoop), 25°43'S 19°19'E  
 Sukses Dam, Etosha Game Park (Outjo), 18°59'S 15°29'E  
 Sutherland, C, 32°24'S 20°40'E  
 S.W.A. Admin Borehole 6453, Bushmanland, 19°22'S 19°37'E  
 Swakop riv. bed, 20 km from Swakopmund, (Swakopmund), 22°39'S 14°44'E  
 Swakop riv. Canyon, (Swakopmund), 22°47'S 15°04'E  
 Swakop riv. mouth, (Swakopmund), 22°41'S 14°33'E  
 Swakopmund (Swakopmund), 22°41'S 14°32'E  
 Swartbaas West 276, (Keetmanshoop), 27°01'S 19°42'E  
 Swartberg, (Lüderitz), 27°08'S 15°26'E  
 Swartbooisdrift, Cunene riv., Kaokoland, 17°20'S 13°52'E  
 Swartmodder farm, C, 27°58'S 20°34'E  
 Swartpoort, Richtersveld near Ochta, C, 28°07'S 16°56'E  
 Tantus 30, (Windhoek), 23°08'S 16°15'E  
 Tinkas Mts, (Karibib), 22°43'S 15°29'E  
 Tinkas Plains, (Swakopmund), 22°50'S 15°25'E  
 Torra Bay, Skeleton Coast Park, (Swakopmund), 20°12'S 13°11'E  
 Torra Bay, 30 km SE of, (Swakopmund), 20°23'S 13°22'E  
 Toz Du Cunene, A, 17°16'S 11°48'E  
 Trekkopje 120, (Karibib), 22°19'S 15°07'E  
 Truidia 300, (Outjo), 19°32'S 15°20'E  
 Tses, (Keetmanshoop), 25°53'S 18°07'E  
 Tsirub 13, (Lüderitz), 26°52'S 16°02'E  
 Tsotsana, Kavango, 19°04'S 20°47'E  
 Tsuandes 107, (Outjo), 20°13'S 15°38'E  
 Tumas Mts, Namib Desert Park, (Walvis Bay), 23°07'S 15°30'E  
 Twee Rivieren, C, 26°30'S 20°35'E  
 Twilight, (Mariental), 24°17'S 17°50'E  
 Twilight 113, (Mariental), 24°12'S 17°56'E  
 Twyfelfontein 584, Damaraland, 20°35'S 14°22'E  
 Ugab Drift, Damaraland, 20°59'S 14°47'E  
 Ugab River Bridge, see Onverwag farm  
 Ugab riv. mouth, (Swakopmund), 21°12'S 13°36'E  
 Uis, Damaraland, 21°09'S 14°47'E  
 Uisib 427, (Grootfontein), 19°32'S 17°14'E  
 Uitsig 82, (Bethanien), 27°37'S 17°10'E  
 Unjab River, 3 km from river mouth, (Swakopmund), 20°09'S 13°13'E  
 Unjab river mouth, 10 km N of, (Swakopmund), 20°06'S 13°06'E  
 Unjabmond, (Swakopmund), 20°09'S 13°10'E  
 Upington, 8 km NW of, C, 28°31'S 21°05'E  
 Us 162, (Karasburg), 27°25'S 18°38'E  
 Vaalbank 319, (Mariental), 23°56'S 18°55'E  
 Valencia 42, Hakos Mts, (Windhoek), 23°11'S 16°30'E  
 Vanzylsrus, C, 26°52'S 22°04'E  
 Vioolsdrift, (Karasburg), 28°46'S 17°37'E  
 Visrivier, near Uis Mine, Damaraland, 21°09'S 14°47'E  
 Vrede 719, Damaraland, 20°23'S 14°14'E  
 Vredenhof 301, (Karasburg), 27°37'S 18°48'E

Waltersdorf 202, (Grootfontein), 19°54'S 18°21'E  
 Walvis Bay, (Walvis Bay), 22°56'S 14°29'E  
 Warmbad, (Karasburg), 28°27'S 18°44'E  
 Warmquelle (Kaokoland), 19°10'S 13°16'E  
 Wasservalle 382, (Windhoek), 22°55'S 16°25'E  
 Waterberg 416, (Otjiwarongo), 20°31'S 17°16'E  
 Welwitchia, (Outjo), see Khorixas  
 Wilhelmstal 8, (Karibib), 21°54'S 16°20'E  
 Williston, C, 31°21'S 20°55'E  
 Windhoek, (Windhoek), 22°34'S 17°06'E  
 Witmanshaar 25, (Bethanien), 25°48'S 16°42'E  
 Wlotzkas Baken (Swakopmund), 22°25'S 14°28'E  
 Zebrafontein, Damaraland, 20°39'S 15°14'E  
 Zebra Mountains, Kaokoland, 17°08'S 13°27'E  
 Zebra Pan, Namib Desert Park (Walvis Bay), 23°30'S 15°29'E  
 Zuidrivier, Namib Desert Park, (Walvis Bay), 23°32'S 15°00'E  
 Zwartbank, (Walvis Bay), 23°22'S 14°54'E

TABLE 1

*Distribution of Namibian scorpions according to habitat predilections.*

Habitat	Key description	Genera	Number of species	%
Underground or in caves	Endogeal EUEDAPHIC	<i>Lisposoma</i>	1	1,8
Burrows dug in soil	Fossorial HEMIEDAPHIC	<i>Opistophthalmus</i>	24	42,8
Shallow scrapes under rocks	Infralapidicolous HEMIEDAPHIC	<i>Lisposoma</i> <i>Buthosus</i> <i>Parabuthus</i> <i>Uroplectes</i> <i>Karasbergia</i>	21	37,8
Rock crevices and fissures	Infrasaxicolous and lithocasicolous HEMIEDAPHIC	<i>Hadogenes</i> <i>Uroplectes</i>	5	8,8
Under bark of trees and on vegetation ranging from shrubs to the largest trees	Supra- and infra-corticicolous EPIGEIC	<i>Buthosus</i> <i>Parabuthus</i> <i>Uroplectes</i>	5	8,8

TABLE 2

*Soil categories classified according to hardness ranges expressed as penetration force ranges (From Lamoral, 1978).*

Categories	Penetration force ranges in kg cm <sup>-2</sup>	Categories	Penetration force ranges in kg cm <sup>-2</sup>
I	0·02-0·1	XII	8·1- 9
II	0·11-0·5	XIII	9·1-10
III	0·51-1·0	XIV	10·1-12
IV	1·01-1·5	XV	12·1-14
V	1·51-2·0	XVI	14·1-16
VI	2·1 -3	XVII	16·1-18
VII	3·1 -4	XVIII	18·1-20
VIII	4·1 -5	XIX	20·1-22
IX	5·1 -6	XX	22·1-24
X	6·1 -7	XXI	24·1-26
XI	7·1 -8	XXII	26·1-28

## TABLES

TABLE 3

*Approximate areas of climatic regions of Namibia (After Barnard, 1965 and Köppen, 1931). Abbreviations: B = General dry zone in which evaporation exceeds precipitation, W = Desert, S = Semi-desert or steppes, h = average annual temperature above 18° C, k = average annual temperature below 18° C, g = warmest month in early summer, t' = warmest month in early autumn, l = all monthly averages between 22° and 10° C, w = summer rains, x' = scarce, intense rainstorms throughout the year, n = regular mist.*

Main type	Region	Climatic Code	Area in square kilometres	%
Cool Desert	Namib Coast Central Plateau Southern Plateau	BW kln BW kw BW kx'	57630 24700 8230	7 3 1
Warm Desert	Namaland Namib Interior Eastern Kaokoveld Orange basin	BW hw BW ht'w BW hgw BW hx'	189360 98800 32930 24700	23 12 4 3
Steppes	Damaraland Central Highlands North and North-East	BS hw BS kw BS hgw	164660 8230 214058	20 1 26

TABLE 4

*Orthobothriotaxies of type A (Buthidae) and C (Scorpionidae) (Abbreviations in Table 5).*

TYPE A: BUTHIDAE Total number of fundamental $\tau = 39$				TYPE C: SCORPIONIDAE Total number of fundamental $\tau = 48$	
Segments	Surface	Number of $\tau$	Codes	Number of $\tau$	Codes
FEMUR	internal dorsal external ventral	4 5 2 0 } 11	i <sub>1</sub> i <sub>2</sub> i <sub>3</sub> i <sub>4</sub> d <sub>1</sub> to d <sub>5</sub> e <sub>1</sub> e <sub>2</sub>	1 1 1 0 } 3	i d e
TIBIA	internal dorsal external ventral	1 5 7 0 } 13	i d <sub>1</sub> to d <sub>5</sub> eb <sub>1</sub> eb <sub>2</sub> esb <sub>1</sub> esb <sub>2</sub> em est et	1 2 13 3 } 19	i d <sub>1</sub> d <sub>2</sub> eb <sub>1</sub> to eb <sub>6</sub> esb <sub>1</sub> esb <sub>2</sub> em <sub>1</sub> em <sub>2</sub> est et <sub>1</sub> et <sub>2</sub> et <sub>3</sub> v <sub>1</sub> v <sub>2</sub> v <sub>3</sub>
HAND	dorsal external ventral	0 6 2 } 8	Eb <sub>1</sub> Eb <sub>2</sub> Eb <sub>3</sub> Esb Est Et V <sub>1</sub> V <sub>2</sub>	2 10 4 } 16	Db Dt Eb <sub>1</sub> Eb <sub>2</sub> Eb <sub>3</sub> Esb Est Et <sub>1</sub> to Et <sub>5</sub> V <sub>1</sub> to V <sub>4</sub>
FIXED FINGER	internal dorsal external	1 2 4 } 7	it db dt et est esb eb	2 4 4 } 10	ib it db dsb dst dt et esb est et

TABLE 5

*Key to trichobothrial codes used in Table 4 and elsewhere in the text and figures.*

Codes	Trichobothrial positions	Codes	Trichobothrial positions
i d e i d eb esb em est et v Db Dt Eb	internal dorsal external } FEMUR  internal dorsal external basal external suprabasal external median external subterminal external terminal ventral } TIBIA  dorsal basal dorsal terminal external basal } HAND	Esb Est Et V it db dsb dst dt eb esb est et	external suprabasal external subterminal external terminal ventral internal terminal dorsal basal dorsal suprabasal dorsal subterminal dorsal terminal external basal external suprabasal external subterminal external terminal } HAND (cont.)  } FIXED FINGER

TABLE 6  
*Character states separating species of Opistophthalmus from Namibia. 1-2 indicates variation in both sexes while 1/2 indicates sexual dimorphism with condition in ♂ shown first.*

CHARACTER STATES	CARAPACE	jenseni																						
		holmi	concentinus	pygmaeus sp.n.	fravescens	perrierorum sp.n.	wahlbergi	chrysistes	serifrons	adultus	schultzei	lorrane sp.n.	fritzsimoni	intercedens	gibbericaudata sp.n.	opinatus	cavimanaus	litioralis	carinatus	ugabensis	breviceauda	hackeri	gligas	
1. Anterior median suture: (1) present; (2) absent.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
2. Median eyes position: (1) antero-medial; (2) postero-medial; (3) distinctly posterior; (4) intermediate to (2) and (3).	3	3	4	4	4	2	2-4	2	2	2	2	4	2	2	4	1	1	1	1	1	1	1	1	1
3. Intercocular surface: (1) granular; (2) smooth or very nearly so.	2	2	1	2	2	1	1-2	1/2	1	2	1	1-2	2	1	2	2	2	1	1	1	1	1	1	1
4. Lateral surfaces: (1) granular; (2) smooth or very nearly so.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
5. Postero-median surface: (1) granular; (2) smooth or very nearly so.	1	1	1	1	2	1	1	1	1	1	1	2	1	2	1	2	1	1	1	1	1	1	1	1
6. Superciliary crests: (1) shallow; (2) well developed.	1	1	2	1	1	2	1	2	1	2	1	2	1	2	1	2	1	1	1	2	1	1	1	1
7. Median eyes size: (1) small; (2) large.	1	1	1	2	2	1	2	1	2	1	2	1	2	1	2	1	2	1	1	2	1	2	1	1
8. Median eyes, distance apart: (1) greater than eye diameter; (2) equal to or less than eye diameter.	1	1	1	2	2	1	1	2	1	1	2	1	1	2	1	1	2	1	1	2	1	2	1	1
9. Im/lhb ♂: (1) 1,20-1,40; (2) 1,41-1,60; (3) 1,61+.	3	2	2	2	2	2	1-3	1	1	1	<1	1	1	3	1	2	2	3	<1	<1	<1	<1	<1	<1
10. Im/lhb ♀: (1) 1,10-1,20; (2) 1,21-1,30; (3) 1,31+.	3	1	3	3	3	3	2-3	1	1	1	—	3	1	3	1	3	3	3	<1	2	<1	—	—	—
11. Upper surface handback: (1) smooth or very shallowly reticulated; (2) granular.	1	1	1	1	1	1	1	1	1	2	2	2	2	2	2	2	2	2	1	1	2	2	2	2
12. Finger keel: (1) predominantly costate; (2) predominantly granular; (3) indistinct.	1	1	1	1	1	1	1	1	1	2	2	2	2	2	2	2	2	2	1	1	1	1	1	1
13. Outer ventro-lateral keel: (1) predominantly costate; (2) predominantly granular.	1	2	2	1	2	2	1	2	1	2	2	1	2	1	1	1	1	1	1	1	1	3	3	3



TABLE 6 continued

CHARACTER STATES	gtigas	hackeri	breviceauda	ugabensis	cavimannus	opinatus	cotteezi	fritzimontsi	lornae	schultzii	adultus	setifrons	chrysistes	wahlbergi	perithorum	flavescens	pygmaeus	continuus	holmi	jenseni
CAUDAL SEGMENTS (cont.)																				
35. II, ventro-lateral keels: (1) absent to very obsolete; (2) costate. (See also 70)	2	2	2	2	1	2	1	1	1	1	1	1	1	2/1	2	1	2	1	1	1
36. III, dorsal keels: (1) absent to obsolete; (2) present and distinct.	2	2	2	2	2	2	1	2	2	2	2	2	2	2	2	1	1	2	1	1
37. III, ventral keels: (1) absent to obsolete; (2) costate; (3) costate granular.	2	2	2	2	2	2	1	1	1	1	1	1	1	4	2/1	2	1	1	3	1
38. II, ventro-lateral keels: (1) ibid.; (2) ibid.; (3) ibid.; (4) granular.	2	2	2	2	1	2	1	1	1	1	1	1	4	2/1	2	1	1	3	1	1
39. II, lateral profile of ventral surface: (1) sublinear; (2) shallowly convex; (3) noticeably convex.	2	2	2	2	1	2	1	1	1	1	1	1	1	4	2/1	2	1	1	3	1
40. III, ventral surface: (1) smooth; (2) granular; (3) punctate.	2	1	2	1	1	2	2	2	2	1	2	2	2	3	1	2	2	2	2	2
41. IV, dorsal keels: (1) absent to obsolete; (2) present and distinct.	1	1	1	1	1	2/1	1	1	3	2/1	1	2	1	1	1	1	1	3	2	2
42. IV, dorsal keels distal spine: (1) not enlarged; (2) slightly enlarged; (3) very distinctly enlarged and spiniform.	2	2	2	2	2	2	1	2	2	2	2	2	2	2	2	2	1	1	2	2
43. IV, ventral keels: (1) absent to very obsolete; (2) costate; (3) costate granular; (4) granular.	3	3	3	2	3	3	2	1-2	2	1	2	2	3	2	1	2	2	3	1	1
44. IV, ventro-lateral keels: (1) ibid.; (2) ibid.; (3) ibid.	3	2	3	3	3	1	3	1	1	3	4	1	1	4	4	1	3	1	4	1
45. IV, ventral surface in ♂: (1) smooth; (2) granular.	3	3	3	3	3	1	3	2	1	3	4	1	1-2	4	4	3	1	3	1	3
46. IV, ventral surface in ♀: (1) smooth; (2) granular.	1	1	2	2	1-2	1	2	1-2	1	2	2	2	1	2	1	1	1	2	2	2
47. IV, lateral profile of ventral surface: (1) sublinear; (2) shallowly convex; (3) noticeably convex or arcuate.	1	1	1	2	1-2	1	1	1	2	1-2	1	—	1	1	2	1	1	1	2	2
48. V, dorsal surface in ♂: (1) smooth; (2) granular.	2	2	2	1	1	1	1	2	2	3	2	2	2	2	2	2	2	2	2	2
49. V, dorsal surface in ♀: (1) smooth; (2) granular.	1	1	1	1	1	1	1	1	1	2	1	2	1	2	1	1	1	1	1	1

50.	V, dorsal keels; (1) absent; (2) partially developed; (3) distinct along entire length.	3	2	2/1	2	2	3	2/1	3	1	1	2	2	3	2	3	2	2	2	1	1
51.	V, ventral keel: (1) absent or indistinct from adjacent granules; (2) single row of granules.	2	2	2	2	2	2	2	2	2	1	2	2	1	1	1	1	1	1	1	1
52.	V, ventro-lateral keels: (1) granular; (2) spiniform or denticulate.	2	2	2	2	2	1	1	1	1	1	1	2	1	1	1	1	1	1	1	1
53.	V, ventro-lateral keels: (1) posteriorly convergent; (2) posteriorly divergent; (3) sub-parallel to each other.	2	2	2	2	2	2	2	2	1	3	2	2	3	3	2	2	1	1	1	1
54.	V, ventral intercarinal surface:	2	2	2	2	1	1	2	2	2	1	2	2	2	1	2	2	2	2	2	2
55.	V, ventral intercarinal surface: (1) sparsely granular; (2) granular. V, lateral profile of ventral surface: (1) shallowly concave; (2) shallowly convex; (3) sublinear.	2	2	2	2	2	2	2	2	3	1	2	3	2	2	3	1	3	1	3	1
<b>TELSON</b>																					
56.	Vesicle ventral surface: (1) smooth; (2) lightly to distinctly granular.	1	1	1	1	1	1	2	2	2	2	2	1	2	2	1	1	2	2	2	2
57.	Vesicle lateral surface: (1) smooth; (2) lightly to distinctly granular.	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	1	1	1	1
58.	Vesicle shape: (1) globose; (2) elongated.	1	1	2	2	2	2	1	1	1	2	2	2	2	1	1	1	2	2	2	2
<b>LEGS</b>																					
59.	Basitarsi I and II, posterior surface with: (1) row of 3-4 spine-like setae; (2) a few stiff setae; (3) many scattered setae; (4) comb-like row of stiff setae.	1	1	1	1	3	1	1	1	2	1	1	4	1	4	3	3	4	4	4	4
60.	Telotarsi III, ventral anterior row of spine-like setae: (1) absent; (2) present + number in brackets.	(1)	(2)	(3)	(2)	(2)	(3)	(2)	(2)	(1)	(1)	(1)	(1)	(1)	(1)	(1)	(1)	(2)	(2)	(1)	(1)
61.	Telotarsi III, ventral posterior row of spine-like setae: (1) absent; (2) present + number in brackets.	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)
62.	Telotarsi IV, ventral anterior row of spine-like setae: (1) absent; (2) present + number in brackets.	(7)	(7)	(5)	(6)	(5)	(6)	(5)	(6)	(5)	(5)	(5)	(5)	(5)	(5)	(5)	(5)	(3)	(3)	(4)	(4)
63.	Telotarsi IV, ventral posterior row of spine-like setae: (1) absent; (2) present + number in brackets.	(3)	(3)	(2)	(3)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)
64.	Telotarsi III and IV, ventral surface with: (1) no setae; (2) few setae; (3) many setae.	(7)	(7)	(5)	(6)	(5)	(6)	(5)	(6)	(5)	(5)	(5)	(5)	(5)	(5)	(5)	(5)	(3)	(3)	(4)	(4)
65.	Telotarsi lateral claws shape: (1) short and distinctly curved; (2) long and distally curved; (3) long and distally sublinear.	1	1	2	1	1	1	1	1	1	1	1	1	1	1	1	1	2	1	1	3

TABLE 6 continued

CHARACTER STATES	LEGS (cont.)	gligas	hackett	brevitacuda	ugabensis	carinatus	litoralis	cavimacrus	opinatus	coetzei	gibbericauda	intercedens	fritzsimonsi	lornae	schultzei	adultus	setifrons	chrysistes	wahlbergi	penithorrum	flavescens	pygmatus	continuus	holmi	jenseni
66. Telotarsi I-II, lateral claws length within each pair: (1) equal; (2) unequal.	1	1	1	1	2	1	1	2	1	2	1	1	2	1	2	2	1	2	2	1	2	1	2	2	2
67. Telotarsi III-IV, lateral claws length within each pair: (1) equal; (2) unequal.	1	1	1	1	2	1	1	2	1	1	1	1	1	1	1	1	1	2	1	2	1	2	1	1	1
68. Telotarsi median dorsal lobe, length in relation to lateral lobes: (1) equal or subequal; (2) distinctly shorter.	1	1	2	2	1	1	2	2	2	2	2	2	2	2	2	2	1	2	1	2	1	2	1	2	2
OTHER																									
69. Carapace, anterior furcated suture: (1) trifurcate, long and distinct; (2) bifurcate, short, obsolete to distinct; (3) absent.	3	3	3	2	2	2	2	2	3	2	2	2	3	1	3	3	3	3	3	3	3	3	3	3	3
70. Cauda II, ventral surface in ♂: (1) smooth; (2) granular (3) with transverse ridges; (4) punctate.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	2	3	2	3	2	1	1	4	1	2	2

TABLE 7  
*Characters subject to variations within distribution range of adults of  
 Opistophthalmus carinatus.*

Locality	Sex	Number of pectinal teeth	Transverse wrinkles on Sternite VII and cauda I; + = present; — = absent; ± = obsolete
Kaoko Otavi, Kaokoland	♂	18-20	+
Kaoko Otavi, Kaokoland	♂	19-20	+
South-western Angola	♂	23-23	+
Outjo	♂	24-23	+
Homob, Etosha Park	♂	21-22	+
Aus, Etosha Park	♂	21-22	+
Gobabeb, Namib Desert Park	♂	26-26	±
Windhoek	♂	28-27	+
Windhoek	♂	21-21	+
Windhoek	♂	23-22	+
Aus, Lüderitz district	♂	25-26	±
Rundu, Kavango	♂	26-26	±
Otjikoto, Tsumeb district	♂	25-25	±
Ghobab, SE of Rehoboth	♂	23-23	±
Gautshe Pan, Bushmanland	♂	26-27	±
Gobabis	♂	27-28	—
Keetmanshoop	♂	27-29	—
Mata Mata, Gemsbok Park	♂	29-30	—
Mata Mata, Gemsbok Park	♂	25-26	—
Kaoko Otavi, Kaokoland	♂	14-15	—
Sesfontein, Kaokoland	♂	17-17	—
Dunedin, Kaokoland	♂	12-12	—
South-western Angola	♂	17-17	—
Outjo	♂	18-18	—
Aus, Etosha Park	♂	18-18	—
Emeritus, Karibib district	♂	16-16	—
Gobabeb, Namib Desert Park	♂	19-19	±
Windhoek	♂	16-16	±
Windhoek	♂	17-17	±
Tsumeb	♂	14-15	—
Rehoboth	♂	18-18	—
Mata Mata, Gemsbok Park	♀	18-20	—

TABLE 8

*Characters subject to clinal variations within distribution range of adults of Opisthoththalmus opinatus. Morphometric data listed as means from specimens in list of material examined.*

Characters		Northern region	Intermediate region	Southern region (South of 26° of latitude)
Total body length in mm	♀ ♂	110 115	70 75	55 60
Carapace length in mm	♀ ♂	15,5 14,5	12,5 11,5	9,2 8,2
$\frac{lc}{x}$	♀ ♂	1,70 1,68	1,80 1,81	1,88 1,90
Interocular surface granulation	♀ ♂	Smooth occasionally with a few granules	lightly granular	granular
Number of pectinal teeth per pecten	♀ ♂	19 23	16 21	13 19
$\frac{whb}{lc}$	♀ ♂	0,80 0,70	0,85 0,78	0,85 0,79
$\frac{lmf}{lhb}$	♀ ♂	1,36 1,70	1,27 1,45	1,20 1,30
Hemispermatophore				
$\frac{ha \rightarrow w \text{ dist.} \times 100}{dcr \rightarrow w \text{ dist.}}$		36,0	32,0	30,0
$\frac{ha \rightarrow bsh \text{ dist.} \times 100}{ha \rightarrow w \text{ dist.}}$		45,0	36,0	30,0
$\frac{ha \rightarrow bsh \text{ dist.} \times 100}{dcr \rightarrow bsh \text{ dist.}}$		22,0	15,0	10,0

## INDEX TO GENERA AND SPECIES

New synonyms are in italics followed by the name recognised in this monograph. Species names are followed by their generic name in brackets. Previously synonymised species are listed in Lamoral & Reynders (1975).

	<i>Page</i>
<i>adustus</i> ( <i>Opistophthalmus</i> ) . . . . .	671
<i>aeratus</i> ( <i>Buthotus</i> ) = <i>conspersus</i> . . . . .	549
<i>alstoni</i> ( <i>Uroplectes</i> ) = <i>carinatus</i> . . . . .	618
<i>arenaceus</i> ( <i>Buthotus</i> ) . . . . .	541
<i>australis</i> ( <i>Parabuthus</i> ) = <i>laevifrons</i> . . . . .	586
<i>bergeri</i> ( <i>Parabuthus</i> ) = <i>granulatus</i> . . . . .	571
<i>bifossulatus</i> ( <i>Hadogenes</i> )—dubious sp. . . . .	661
<i>brachystylus</i> ( <i>Parabuthus</i> ) . . . . .	616
<i>bradfieldi</i> ( <i>Opistophthalmus</i> ) = <i>opinatus</i> . . . . .	733
<i>brevicauda</i> ( <i>Opistophthalmus</i> ) . . . . .	676
<i>brevimanus</i> ( <i>Parabuthus</i> ) . . . . .	561
<i>Buthotus</i> . . . . .	541
<i>carinatus</i> ( <i>Uroplectes</i> ) . . . . .	618
<i>carinatus</i> ( <i>Opistophthalmus</i> ) . . . . .	679
<i>cavimanus</i> ( <i>Opistophthalmus</i> ) . . . . .	688
<i>chrysites</i> ( <i>Opistophthalmus</i> ) . . . . .	692
<i>coetzeei</i> sp. n. ( <i>Opistophthalmus</i> ) . . . . .	694
<i>concinnus</i> ( <i>Opistophthalmus</i> ) . . . . .	698
<i>concolor</i> ( <i>Parabuthus</i> ) = <i>stridulus</i> . . . . .	606
<i>conspersus</i> ( <i>Buthotus</i> ) . . . . .	549
<i>cristatus</i> ( <i>Parabuthus</i> ) = <i>brevimanus</i> . . . . .	561
<i>elegans</i> ( <i>Lisposoma</i> ) . . . . .	662
<i>fitzsimonsi</i> ( <i>Opistophthalmus</i> ) . . . . .	702
<i>flavescens</i> ( <i>Opistophthalmus</i> ) . . . . .	704
<i>fulvipes</i> ( <i>Parabuthus</i> ) = <i>granulatus</i> . . . . .	571
<i>fuscus</i> ( <i>Parabuthus</i> ) = <i>granulatus</i> . . . . .	571
<i>gibbericauda</i> sp. n. ( <i>Opistophthalmus</i> ) . . . . .	707
<i>gaerdesi</i> ( <i>Opistophthalmus</i> ) = <i>brevicauda</i> . . . . .	676
<i>gariepensis</i> ( <i>Opistophthalmus</i> ) = <i>wahlbergi</i> . . . . .	756
<i>gigas</i> ( <i>Opistophthalmus</i> ) . . . . .	711
<i>gracilior</i> ( <i>Uroplectes</i> ) . . . . .	652
<i>gracilis</i> sp. n. ( <i>Parabuthus</i> ) . . . . .	566
<i>granulatus</i> ( <i>Parabuthus</i> ) . . . . .	571
<i>haackei</i> ( <i>Opistophthalmus</i> ) . . . . .	714
<i>Hadogenes</i> . . . . .	652
<i>histrio</i> ( <i>Opistophthalmus</i> ) = <i>carinatus</i> . . . . .	679
<i>holmi</i> ( <i>Opistophthalmus</i> ) . . . . .	716
<i>ibelli</i> ( <i>Parabuthus</i> ) = <i>laevifrons</i> . . . . .	586
<i>intercedens</i> ( <i>Opistophthalmus</i> ) . . . . .	719
<i>jenseni</i> ( <i>Opistophthalmus</i> ) . . . . .	723
<i>josehermana</i> sp. n. ( <i>Lisposoma</i> ) . . . . .	665
<i>kalaharicus</i> ( <i>Parabuthus</i> ) . . . . .	576
<i>Karasbergia</i> . . . . .	555
<i>karoicus</i> ( <i>Uroplectes</i> ) = <i>schlechteri</i> . . . . .	643
<i>kraepelini</i> ( <i>Parabuthus</i> ) . . . . .	579
<i>kuanyamarum</i> ( <i>Parabuthus</i> ) . . . . .	583
<i>laevicauda</i> ( <i>Opistophthalmus</i> ) = <i>schultzei</i> . . . . .	745
<i>laevifrons</i> ( <i>Parabuthus</i> ) . . . . .	586
<i>lawrencei</i> ( <i>Hadogenes</i> ) = <i>tityrus</i> . . . . .	657
<i>Lisposoma</i> . . . . .	661
<i>litoralis</i> ( <i>Opistophthalmus</i> ) . . . . .	726
<i>longiceps</i> ( <i>Opistophthalmus</i> ) = <i>adustus</i> . . . . .	671
<i>longimanus</i> ( <i>Uroplectes</i> ) . . . . .	629
<i>lornae</i> sp. n. ( <i>Opistophthalmus</i> ) . . . . .	729
<i>luciranus</i> ( <i>Opistophthalmus</i> ) = <i>ugabensis</i> . . . . .	753
<i>mediostriatus</i> ( <i>Uroplectes</i> ) = <i>carinatus</i> . . . . .	618
<i>methueni</i> ( <i>Karasbergia</i> ) . . . . .	555
<i>militum</i> ( <i>Parabuthus</i> ) = <i>stridulus</i> . . . . .	606

	<i>Page</i>
namibensis sp. n. ( <i>Parabuthus</i> ) . . . . .	591
nanus sp. n. ( <i>Parabuthus</i> ) . . . . .	594
<i>nigrocarinatus</i> ( <i>Opistophthalmus</i> ) = <i>setifrons</i> . . . . .	750
<i>nigrovesicalis</i> ( <i>Opistophthalmus</i> ) = <i>wahlbergi</i> . . . . .	756
<i>opinatus</i> ( <i>Opistophthalmus</i> ) . . . . .	733
<i>Opistophthalmus</i> . . . . .	668
<i>otjimbinguensis</i> ( <i>Uroplectes</i> ) . . . . .	632
<i>Parabuthus</i> . . . . .	560
<i>penritorum</i> sp. n. ( <i>Opistophthalmus</i> ) . . . . .	738
<i>phyllioides</i> ( <i>Hadogenes</i> ) . . . . .	653
<i>pilosus</i> ( <i>Uroplectes</i> ) . . . . .	635
<i>planimanus</i> ( <i>Uroplectes</i> ) . . . . .	638
<i>Protophthalmus</i> = <i>Opistophthalmus</i> . . . . .	668
<i>pygmaeus</i> sp. n. ( <i>Opistophthalmus</i> ) . . . . .	742
<i>raodus</i> ( <i>Parabuthus</i> ) . . . . .	598
<i>scabriceps</i> ( <i>Opistophthalmus</i> ) = <i>brevicauda</i> . . . . .	676
<i>scabifrons</i> ( <i>Opistophthalmus</i> ) = <i>opinatus</i> . . . . .	733
<i>schlechteri</i> ( <i>Parabuthus</i> ) . . . . .	602
<i>schlechteri</i> ( <i>Uroplectes</i> ) . . . . .	643
<i>schultzei</i> ( <i>Opistophthalmus</i> ) . . . . .	745
<i>setifrons</i> ( <i>Opistophthalmus</i> ) . . . . .	750
<i>setiventer</i> ( <i>Opistophthalmus</i> ) = <i>intercedens</i> . . . . .	719
<i>stridulus</i> ( <i>Parabuthus</i> ) . . . . .	606
<i>taenius</i> ( <i>Hadogenes</i> ) . . . . .	646
<i>teretipes</i> ( <i>Uroplectes</i> ) . . . . .	649
<i>tityrus</i> ( <i>Hadogenes</i> ) . . . . .	657
<i>tumidimanus</i> sp. n. ( <i>Uroplectes</i> ) . . . . .	753
<i>ugabensis</i> ( <i>Opistophthalmus</i> ) . . . . .	745
<i>undulatus</i> ( <i>Opistophthalmus</i> ) = <i>schultzei</i> . . . . .	618
<i>Uroplectes</i> . . . . .	610
<i>villosus</i> ( <i>Parabuthus</i> ) . . . . .	652
<i>vittatus</i> ( <i>Uroplectes</i> ) . . . . .	750
<i>vivianus</i> ( <i>Opistophthalmus</i> ) = <i>setifrons</i> . . . . .	756
<i>wahlbergi</i> ( <i>Opistophthalmus</i> ) . . . . .	762
<i>wernerii</i> ( <i>Opistophthalmus</i> )—dubious sp.	