

DET KGL. NORSKE VIDENSKABERS SELSKAB
MUSEET


GUNNERIA
28



Torleif Holthe

A QUANTITATIVE INVESTIGATION OF THE LEVEL-BOTTOM
MACROFAUNA OF TRONDHEIMSFJORDEN, NORWAY

TRONDHEIM 1977



A QUANTITATIVE INVESTIGATION OF THE LEVEL-BOTTOM
MACROFAUNA OF TRONDHEIMSFJORDEN, NORWAY

by

Torleif Holthe

University of Trondheim

The Royal Norwegian Society of Sciences and Letters, The Museum

ISBN 82-7126-147-9

ABSTRACT

Holthe, Torleif. 1977. A quantitative investigation of
the level-bottom macrofauna of Trondheimsfjorden, Norway.
Gunneria 28 : 1-20.¹⁾

As part of an investigation of Trondheimsfjorden as a recipient, a survey of the level-bottom macrofauna was undertaken during the period June 1972 to October 1974. The survey comprised 1,580 grab samples and covered 30 stations in the main fjord and the side fjords. A total of 310 macrobenthos taxa was identified. The results indicate that the benthos of the main fjord and most side fjords are not much affected by human activity. There are, however, some restricted parts of the fjord showing effects of heavy metal pollution, artificial changes in the sediment, or hyper-eutrophication.

*Torleif Holthe, University of Trondheim, The Royal Norwegian Society of Sciences and Letters, The Museum, Zoological Department,
N-7000 Trondheim*

1) Zoological Series 6.

CONTENTS

INTRODUCTION	7
MATERIAL AND METHODS	8
RESULTS	12
DISCUSSION	13
ACKNOWLEDGEMENTS	18
REFERENCES	19

Tables 2-33

Fig. 2-34

ERRATUM: p. 15

Ytterøya-Skogn Figs. 19-20
read: Figs. 19-22.

INTRODUCTION

The main tasks of the 1972 to 1974 investigation of Trondheims-fjorden as a recipient were to study the physical hydrography and the primary production of the fjord system, and to map sources of possible pollutants. It was, however, considered desireable also to gain information on the state of the biota in the various parts of the fjord. It was necessary to choose a limited group of organisms for this survey, that should as far as possible meet the following conditions: 1) The natural changes of the populations should not be too rapid. 2) The organisms should be stationary. 3) The habitat should be uniform. 4) Quantitative sampling should be possible. 5) Basic research results on synecology and the species' biology should be at hand. 6) It must be possible to find specialists who could identify the organisms and interpret the results.

The group of organisms that came nearest to fulfill these conditions was the level-bottom macrofauna.

Mapping of the sources of pollutants was undertaken by the Norwegian Institute for Water Research (NIVA 1976), investigations on the physical hydrography by The River and Harbour Laboratory at The Norwegian Institute of Technology (Jacobson 1976), investigations on plant nutrients, and primary production by The Royal Norwegian Society of Sciences and Letters, The Museum (Sakshaug 1976). None of these results are yet published, the quotations above refere to the reports to the investigation committee. A paper on the heavy metal pollution has been published (Lande 1977).

MATERIAL AND METHODS

The material at hand was sampled during six cruises in the period June 1972 to October 1974. A total of 30 stations was operated (Fig. 1). A survey of the cruises and stations is given in Table 1.




Fig. 1. Map of Trondheimsfjorden with the position of the sampling stations.

Table 1. Stations and cruises

Depth in meters	OF				BV			TH			SF		SD		AF			PT			YØ		VD		KV		SK		VB	
	1	2	3	4	1	2	3	1	2	3	1	2	1	2	1	2	3	1	2	3	1	2	1	2	2	3	4	1	2	1
June 1972	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
October 1972	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
June 1973	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
October 1973	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
June 1974	x	x	x	x				x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
October 1974	x	x	x				x			x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x

Sampling was performed by means of a 0.1 m^2 Petersen grab. all samples comprise 10 subsamples, each of 0.1 m^2 . At depths of 20 or 50 m the vessel was anchored at the sampling site. At greater depths the vessel was held in position by active maneuver. The material was sieved on board through sieves with a mesh width of 1 mm. Sorting into three main groups, viz. polychaetes, pelecypods, and other invertebrates, was also performed on board. The material was preserved in alcohol, and remains in the collections of The Royal Norwegian Society of Sciences and Letters, The Museum.

Identification was as far as possible carried out to the species level. A few higher taxa are generally left indetermined, these are: Polycladida, Nemertini, Amphipoda, Cumacea, and Holothuroidea.

Data processing was carried out by means of the program series "Benthos", which was originally devised to serve this investigation (Lundquist & Holthe 1974). The program was run at Regnesenteret, University of Trondheim (UNIVAC 1108).

For each station, cruise, and species, there are calculated mean (\bar{x}) and standard deviation (s).

For each station and cruise are calculated three indices of diversity, λ (Simpson 1949), H (Shannon & Weaver 1963), and d (Margalef 1957), and sampling efficiency by Engen's method, S_N (Strömgren et al. 1973).

$$\lambda = \frac{N^2 - \sum x_i^2}{N(N - 1)}$$

$$H \approx - \sum \frac{x_i}{N} \ln \frac{x_i}{N}$$

$$d = \frac{S - 1}{\ln N}$$

where N is the total number of individuals, S is the number of species, and x_i is the number of individuals of the i 'th species.

S_N is calculated from r_1 , the number of species (or higher indetermined taxa) with only one specimen present in the sample. With a level of significance of 0.05 the value of λ_m is taken from Engen's table if $r_1 \leq 10$, and if $r_1 > 10$, λ_m can be calculated as

$$\lambda_m = \frac{1.644 + \sqrt{1.644^2 + 4r_1}}{2}^2$$

Sample efficiency is then calculated by means of the expression:

$$1 - \frac{\lambda_m}{N} < S_N < 1$$

where N is the number of individuals.

For any pair of stations two coefficients of similarity are calculated. These are the quantitative coefficient of community (cc) and the quantitative Czkanowski's coefficient of similarity (Cz).

$$cc = \frac{c}{a + b - c} \cdot 100$$

where a is the number of species of the station A, b is the number of species of the station B, and c is the number of species common to both.

$$C_Z = \frac{2 \sum \min a_i b_i}{\sum a_i + \sum b_i}$$

where a_i and b_i are the numbers of individuals of the i 'th species at the station A and B, respectively.

RESULTS

A total of 310 species and higher indetermined taxa are listed in Table 2. Of the indetermined taxa only Amphipoda indet. and Holothuroidea indet. may represent considerable numbers of species, the latter group is also quantitatively dominant at some stations. The indisputable highest number of species is found among the Polychaeta (at least 143 spp), followed by the Pelecypoda with at least 62 spp.

The species lists and the quantitative results for the 30 stations are given in Table 3-32. The values of the coefficients of similarity are given in Table 33. A simplified presentation of the quantitative results is given in Figs. 2-31.

DISCUSSION

When extensive sampling is undertaken, as in the present investigation, the yield of information about the distributions of the individual species is considerable. A compilation of this and earlier results is yet made only for the polychaete fraction of the fauna (Holthe 1977). This compilation gives no reason to assume that the polychaete fauna has been qualitatively reduced since the turn of the century. Unfortunately there are no earlier quantitative results suited for comparison. Thus the main purpose of the present investigation is to serve as a quantitative baseline for future supervision of the fjord. The variations throughout the period of investigation are certainly caused by natural competition within the communities. No major changes in ecological stress caused by human activity has occurred during the period. It is, however, possible to discuss the state of benthic life in the various parts of the fjord:

Orkdalsfjorden (Stations OF 1-4, Tables 3-6, Figs. 2-5)

For all investigated depths Orkdalsfjorden shows low similarity with the other investigated areas, quantitatively as well as qualitatively. This is not caused by the occurrence of species which are not found elsewhere, but by a low number of species and low diversity. The dominant species are *Scoloplos armiger*, *Heteromastus filiformis*, *Chaetozone setosa*, and *Thyasira* sp. In depths from 20-100 m several species - especially polychaetes - occur in small numbers. In greater depths the number of species and also the number of individuals are very low. All over, the fauna seems to be poorer than it would be by normal competition between the species that have access to the fjord. This indicates the presence of a physical stress on the community.

The causes of this stress cannot be found in the hydrographical conditions nor in the nature of the sediments, as these are not different from the adjoining parts of Trondheimsfjorden. It is known that the bottom of Orkdalsfjorden is unstable; i.e. submarine landslides have occurred (Rosenquist 1960). But if this phenomenon should influence the benthic community structure, landslides must occur every other year and affect major parts of the fjord. The marine benthic

fauna will rapidly recolonize affected areas when the conditions again are favourable. (An example of rapid recolonization in a Swedish fjord is given by Rosenberg (1972)). The tables show that several benthic invertebrate species occur in Orkdalsfjorden, but that most of them do not succeed in establishing normal population densities. It is tempting to assume that the fauna of Orkdalsfjorden is influenced by heavy metal waste from the mining industry in the valley of the only river which discharge into this fjord. Measurements of the heavy metal content of littoral organisms (Lande 1977) and bottom sediment (Fiskum et al. 1974) support this assumption. Orkdalsfjorden is generally more affected by heavy metal pollution than other parts of Trondheimsfjorden (Lande 1977), and the degree of pollution of the sediment increases with the water depth (Fiskum et al. 1974).

Buvika (Stations BV 1-3, Tables 7-9, Figs. 6-8)

In this area all investigated depths (20-100 m) have a fauna rich in species and with high diversity. The variation apparent from one cruise to another at 20 m, is caused by the patchiness of the fauna, and does not describe a seasonal variation.

Trondheim (Stations TH 1-3, Tables 10-12, Figs. 9,11)

In the area north-west of the city (TH-2-3) it proved difficult to obtain samples. The sediment is mixed with coarse sand and stones, which prevent the grab from closing properly. Thus the results from these stations are not reliable as quantitative estimates.

East of the city, at a depth of 50 m, was found a community characterized by high densities of the polychaete *Melinna cristata* and a number of species that surpasses that of any other known level-bottom locality in Trondheimsfjorden. This station lies in the course of the current bringing waste from the city, but the fauna seems to be little affected by this condition.

Strindfjorden (Stations SF 1-2, Tables 13-14, Figs. 12-13)

The fauna of Strindfjorden is characterized by the ophiurids *Amphiura chiajei*, *A. filiformis*, and *Amphilepis norvegica*. These communities are typical for the deeper water (100-200 m), and are to be expected in unpolluted areas. The low number of individuals in October 1974 is probably caused by the difficulty of obtaining good samples in rough weather.

Stjördalsfjorden (Stations SD 1-2, Tables 15-16, Figs. 14-15)

The fauna inside the threshold to Stjördalsfjorden is rich in species, and the diversity is high. Character species are the polychaete *Maldane sarsi* and the pelecypods *Abra alba* and *A. nitida*. This fjord is influenced by sewage from the city of Trondheim (Jacobson 1976), and it is planned to supervise the oxygen conditions in Stjördalsfjorden to detect a possible pollution increase.

Åsenfjorden (Stations ÅF 1-3, Tables 17-19, Figs. 16-18)

These stations are sampled only twice, and the fauna seems to be rather poor in species and individuals. The diversity is not low.

Ytteröya - Skogn (Stations YÖ 1 and FT 1-3, Tables 20-30, Figs. 19-20)

While the fauna of Ytteröya (YÖ 1) in all respects appears natural, the fauna on the other side of the basin shows quantitative anomalies. There are many species present in the area, but the high number of individuals is made up by a few pelecypods, viz. *Thyasira* spp. (especially *T. sarsi*), *Abra alba*, and *A. nitida*. *T. sarsi* seems to be abundant in areas with much dead plant material, and the sediment at Stations FT 1-3 contains much wood fragments. This is not surprising,

as the regions' largest pulp and paper mill is located here. Even if this waste has little chemical effect, its altering of the mechanical properties of the sediment will give preference to certain pelecypod species, and thus influence the community structure.

Verdal (Stations VD 1-2, Tables 24-25, Figs. 23-24)

These stations are sampled only twice. The fauna is relatively rich in species and individuals. Character species are the polychaete *Maldane sarsi* and the pelecypods *Abra alba*, *A. nitida* and *Corbula gibba*.

Beitstadfjorden (Stations KV 2-4 and SK 1-2, Tables 26-30
Figs. 25-27 and 30-31)

In Beitstadfjorden two transects are studied: One from Kirknesvågen and out to a depth of 200 m (KV 2-4), and one outside the harbour of Steinkjer. The transect KV 2-4 can be compared to a similar transect in the main fjord (TH 1, SF 2, SF 1), and these transects show good accordance. The differences are not greater than it should be expected from natural causes when one transect is located in the outer part and another in the inner part of a fjord system. Outside Steinkjer the pelecypods *Thyasira* spp., *Abra alba*, and *A. nitida* occur in high numbers. This can be due to the mechanical properties of the sediment which is unusually tough.

Verrabotn (Stations VB 1-2, Tables 31-32, Figs. 28-29)

In this fjord it is difficult to find level-bottom suited for sampling. The results indicate a fauna with few species and individuals, the diversity is high.

Earlier investigations in Borgenfjorden (Holthe 1973, McClimans 1973, Strömgren 1973) have shown that the inner part of this land-locked basin is hypereutrophicated with following oxygen deficiency. This condition is caused by drainage from the surrounding agricultural areas and by the poor water exchange in the fjord. In Trondheimsfjorden probably only such land-locked basins are threatened by hypereutrophication.

For Trondheimsfjorden in general it must be concluded that the fauna is rich in species and individuals and controlled by natural competition; only locally - in basins with limited water exchange -- effects of human activity can be traced.

ACKNOWLEDGEMENTS

This work is part of an investigation of Trondheimsfjorden as a recipient of domestic, agricultural and industrial waste. The investigation was supported by the Department of Environment and the two regional administrations in the area: Sør-Trøndelag fylke and Nord-Trøndelag fylke.

I wish to thank the specialists who identified parts of the material: Cand.real. Jon-Arne Sneli (*Gastropoda*, *Polyplacophora*, and *Scaphopoda*), Cand.real. Eirik Lande (*Pelecypoda*), and Cand.real. Björn Gulliksen (*Tunicata*).

REFERENCES

- Fiskum, A., B. Aakhus, H. Borchsenius, A. Schelderup, L. Jørgensen, E. Hegseth & B. Os. 1974. *Orkdalsfjorden - forurensningskilder - skadefirkninger - mulige forbedringer.* University of Trondheim, Committee for Environmental Studies.
- Holthe, T. 1973. *Borgenfjordens Polychaeta, Echiurida, Sipunculida, Priapulida, Phoronida og Enteropneusta.* Unpublished thesis, University of Trondheim. 190 pp.
- Holthe, T. 1977. The polychaetous annelids of Trondheimsfjorden, Norway. *Gunneria* 29, in print.
- Jacobson, P. 1976. *Resipientundersøkelse av Trondheimsfjorden.* River and Harbour Laboratory, Norw. Inst. Techn., report.
- Lande, E. 1977. Heavy metal pollution in Trondheimsfjorden, Norway, and the recorded effects on the fauna and flora. *Environ. Pollut.* 12: 187-198.
- Lundquist, P. & T. Holthe. 1974. Brukerveiledning til fire datamaskinprogrammer for kvantitative makrobenthosundersøkelser. *K. norske Vidensk. Selsk. Mus. Rapport Zool. Ser. 1974-8*, 54 pp.
- McClimans, T. 1973. Physical oceanography of Borgenfjorden. *K. norske Vidensk. Selsk. Skr. 2:* 1-43.
- Margalef, R. 1957. La teoria de la informacion en ecologia. *Mems. R. Acad. Cienc. Artes Barcelona* 33: 373-449.
- NIVA, 1976. *Resipientundersøkelse av Trondheimsfjorden 1972-1975. Forurensningstilforsler.* Norsk institutt for vannforskning, report. 115 pp + data appendix.
- Rosenberg, R. 1972. Benthic faunal recovery in a Swedish fjord following the closure of a sulphite pulp mill. *Oikos* 23: 92-108.
- Rosenquist, I. 1960. Marine clays and quick clay slides in south and central Norway. *NGU nr. 212 g.*
- Sakshaug, E. 1976. *Resipientundersøkelsen av Trondheimsfjorden. Sluttrapport. Variasjon i planteneringsstoffer og klorofyll a.* Royal Norwegian Society of Sciences and Letters, The Museum, report. 18 pp.

- Shannon, C.E. & W. Weaver. 1963. *The mathematical theory of communication.* Univ. Illinois Press. 117 pp.
- Simpson, E.H. 1949. Measurement of diversity. *Nature* 163: 688.
- Strömgren, T. 1973. Zooplankton investigations in Borgenfjorden, 1967-1969. *K. norske Vidensk. Selsk. Mus. Miscea* 9, 37 pp.
- Strömgren, T., R. Lande & S. Engen, 1973. Intertidal distribution of the fauna on muddy beaches in the Borgenfjorden area. *Sarsia* 53: 49-70.

table 2

List of taxa encountered:

PORIFERA

Porifera sp. indet.

ANTHOZOA

Anthozoa spp. indet.
Funiculina quadrangularis (Pallas)
Paraedwardsia arenaria Carlgren

Pennatula phosphorea Linnaeus
Stylatula elegans (Koren and Danielsen)
Virgularia mirabilis (Müller)

TURBELLARIA

Polycladida spp. indet.

NEMERTINI

Nemertini spp. indet.

POLYCHAETA

Amage auricula Malmgren, 1865
Ampharete finmarchica (Sars, 1856)
Ampharetidae spp. indet.
Ampharete sp.
Amphicteis gunneri (Sars, 1835)
Amphitrite cirrata Müller, 1771
Amphitritinae spp. indet.
Anobothrus gracilis (Malmgren, 1867)
Antinoella sarsi (Kinberg in Malmgren, 1865)
Aphrodisia aculeata Linnaeus, 1758
Aphroditidae sp. indet.
Artacama proboscidea Malmgren, 1866
Asychis biceps (Sars, 1861)
Brada villosa (Rathke, 1843)
Capitella capitata (Fabricius, 1780)
Capitellidae sp. indet.
Ceratocephale loveni Malmgren, 1867
Chaetozone setosa Malmgren, 1867
Cirratulus cirratus (Müller, 1776)
Cirratulidae sp. indet.
Dasybranchus caducus (Grube, 1846)
Diplocirrus glaucus (Malmgren, 1867)
Diplocirrus longosetosa (Marenzeller, 1890)
Ditrupa arietina (Müller, 1776)
Driloneris filum (Claparede, 1868)
Eclipsippe vanelli (Fauvel, 1936)
Eteoninae sp. indet.
Eteone sp.
Euchone papillosa (Sars, 1851)
Euchone rubrocincta (Sars, 1861)
Euclymeninae spp. indet.
Eulalia sp.
Eumida sp.
Eunice pennata (Müller, 1776)
Flabelligeridae sp. indet.
Gattyana cirrosa (Pallas, 1776)
Genetyllis lutea (Malmgren, 1865)
Glycera alba (Müller, 1788)
Glycera capitata Ørsted, 1843
Glycera lapidum Quatrefages, 1865
Glycera rouxii Audouin and Milne-Edwards, 1833
Glycera sp.
Glyphanostomum pallescens (Theel, 1879)
Goniada maculata Ørsted, 1843

Goniada norvegica Ørsted, 1845
Harmothoe spp.
Hauchiella tribullata (McIntosh, 1869)
Hesionidae sp. indet.
Heteromastus filiformis (Claparède, 1864)
Heteroclymene robusta Arwidson 1906
Hyalinoecia tubicola (Müller, 1776)
Hydroides norvegica Gunnerus, 1768
Laetmonice filicornis Kinberg, 1855
Lanassa nordenskiöldi Malmgren, 1866
Lanassa venusta (Malm, 1874)
Laonice cirrata (Sars, 1851)
Laphania boeckii Malmgren, 1866
Leanira tetragona (Ørsted, 1845)
Leiochone borealis Arwidson, 1906
Lumbrineris fragilis (Müller, 1776)
Lumbrineris sp.
Lumbriclymene minor Arwidson, 1906
Lysilla loveni Malmgren, 1866
Maldanidae spp. indet.
Maldane sarsi Malmgren, 1865
Melinna cristata (Sars, 1851)
Microclymene tricirrata Arwidson, 1906
Myriochele sp.
Neoamphitrite affinis (Malmgren, 1866)
Neoamphitrite edwardsi (Quatrefages, 1865)
Neoamphitrite figulus (Dalliyell, 1853)
Neoamphitrite grayi (Malmgren, 1865)
Neoamphitrite sp.
Nephthys caeca (Fabricius, 1780)
Nephthys ciliata (Müller, 1776)
Nephthys hombergi Savigny, 1818
Nephthys incisa Malmgren, 1865
Nephthys paradoxa Malm, 1874
Nephthys sp.
Nereimyra punctata (Müller, 1776)
Nereis virens Sars, 1835
Nereis sp.
Nicromache lumbricalis (Fabricius, 1780)
Nothria conchylega (Sars, 1835)
Notomastus latericeus Sars, 1851
Onuphis quadricuspis Sars, 1872
Ophelina acuminata Ørsted, 1843
Ophelina norvegica Stør-Bovitz, 1946

<u>Ophelina</u> sp.		<u>Prionospio malmgreni</u>	Claparède, 1870
<u>Ophiromus flexuosus</u>	(Delle Chiaje, 1825)	<u>Pygospio elegans</u>	Claparède, 1863
<u>Owenia fusiformis</u>	Delle Chiaje, 1841	<u>Rhodine gracilior</u>	Tauber, 1879
<u>Panthalis oerstedi</u>	Kinberg, 1855	<u>Rhodine loveni</u>	Malmgren, 1866
<u>Paramphinoe jeffreysi</u>	(McIntosh, 1868)	<u>Rhodine</u> sp.	
<u>Paraonis gracilis</u>	(Tauber, 1879)	<u>Sabellides borealis</u>	Sars, 1856
<u>Pectinaria auricoma</u>	(Müller, 1776)	<u>Sabellides octocirrata</u>	Sars, 1835
<u>Pectinaria belgica</u>	(Pallas, 1766)	<u>Sabella penicillatus</u>	Linnaeus, 1767
<u>Pectinaria koreni</u>	(Malmgren, 1866)	<u>Sabellidae</u> spp. indet.	
<u>Pherusa plumosa</u>	(Müller, 1776)	<u>Samytha sexcirrata</u>	(Sars, 1856)
<u>Pholoe minuta</u>	(Fabricius, 1780)	<u>Scalibregma inflatum</u>	Linnaeus, 1767
<u>Phyllodocidae</u> spp. indet.		<u>Scoloplos armiger</u>	(Müller, 1776)
<u>Phyllodocia laminosa</u>	Savigny, 1818	<u>Sphaerodorum gracilis</u>	(Rathke, 1843)
<u>Phylo kupfferi</u>	(Ehlers, 1874)	<u>Spio filicornis</u>	(Müller, 1776)
<u>Phylo norvegicus</u>	(Sars, 1872)	<u>Spicchaetopterus typicus</u>	Sars, 1856
<u>Pista cristata</u>	(Müller, 1776)	<u>Spiophanes krøyeri</u>	Grube, 1860
<u>Polychaeta</u> spp. indet.		<u>Streblosoma bairdi</u>	(Malmgren, 1866)
<u>Polycirrus medusa</u>	Grube, 1850	<u>Streblosoma intestinalis</u>	Sars, 1872
<u>Polycirrus</u> sp.		<u>Syllidae</u> spp. indet.	
<u>Polydora</u> sp.		<u>Terebellidae</u> spp. indet.	
<u>Polymnia nebulosa</u>	(Montagu, 1818)	<u>Terebellides stroemi</u>	Sars, 1835
<u>Polynoidae</u> spp. indet.		<u>Tharyx marioni</u>	(Saint-Joseph, 1894)
<u>Polynoe kinbergi</u>	(Malmgren, 1865)	<u>Thelepus cincinnatus</u>	(Fabricius, 1780)
<u>Polyphysia crassa</u>	(Ørsted, 1843)	<u>Thelepiniae</u> spp. indet.	
<u>Praxillella gracilis</u>	(Sars, 1861)	<u>Trichobranchus roseus</u>	(Malm, 1874)
<u>Praxillella praetermissa</u>	(Malmgren, 1865)	<u>Trochochaeta multisetosa</u>	(Ørsted, 1844)
<u>Prionospio cirrifera</u>	Wirén, 1883	<u>Typosyllis cornuta</u>	(Rathke, 1843)

SIPUNCULIDA

<u>Onchnesoma squamatum</u>	(Koren and Danielsen, 1875)	<u>Phascolosoma</u> spp.
<u>Onchnesoma steenstrupi</u>	Koren and Danielsen, 1875	<u>Sipunculidae</u> spp. indet.
<u>Phascolion strombi</u>	(Montagu, 1804)	

CRUSTACEA

<u>Amphipoda</u> spp. indet.		<u>Geryon tridens</u> Krøyer, 1837
<u>Balanus balanus</u>	(Linnaeus)	<u>Gnathia oxyraea</u> (Lilljeborg)
<u>Calocharis macandreae</u>	Bell 1846	<u>Hyas coarctatus</u> Leach, 1815
<u>Carcinus maenas</u>	(Linnaeus, 1758)	<u>Macropipus depurator</u> (Linnaeus, 1758)
<u>Cumacea</u> spp. indet.		<u>Paguridae</u> spp. indet.
<u>Galathea</u> sp.		

PYCNOGONIDA

<u>Nymphon</u> sp.		<u>Pycnogonida</u> sp. indet.
--------------------	--	-------------------------------

CAUDOPOVEATA

<u>Chaetoderma nitidulum</u>	Lovén
------------------------------	-------

POLYPLACOPHORA

<u>Acanthochiton fascicularis</u>	(Linnaeus)	<u>Lepidopleurus alveolus</u> (Sars)
-----------------------------------	------------	--------------------------------------

GASTROPODA

<u>Actaeon tornatilis</u>	(Linnaeus, 1766)	<u>Cylichna cylindracea</u> (Pennant, 1777)
<u>Admete viridula</u>	(Fabricius, 1780)	<u>Eumila stenostoma</u> Jeffreys, 1858
<u>Alvania abyssicola</u>	(Forbes, 1853)	<u>Eumilella scillae</u> (Scacchi, 1836)
<u>Aporrhais pespelecani</u>	(Linnaeus, 1758)	<u>Iothia fulva</u> (Müller, 1776)
<u>Balcis</u> sp.		<u>Clelandella miliaris</u> (Brocchi, 1814)
<u>Buccinum humphreysianum</u>	Bennett, 1824	<u>Lepeta caeca</u> (Müller, 1776)
<u>Buccinum undatum</u>	(Linnaeus, 1758)	<u>Lunatia intermedia</u> (Philippi, 1836)
<u>Buccinum</u> sp.		<u>Lunatia pallida</u> (Broderip & Sowerby, 1829)
<u>Cylichna alba</u>	(Brown, 1827)	<u>Menestho divisa</u> (J.Adams, 1797)

Nassarius reticulatus (Linnaeus, 1758)
Natica clausa Broderip & Sowerby, 1829
Natica sp.
Neptunea despecta (Linnaeus, 1758)
Odostomia unidentata (Montagu, 1803)
Oenopota elegans (Møller, 1842)
Okenia pulchella (Morlet, 1880)
Philine quadrata (Wood, 1839)
Philine scabra (Müller, 1776)
Philine sp.

Retusa umbilicata (Montagu, 1803)
Scaphander lignarius (Linnaeus, 1766)
Scaphander punctostriatus (Mighels, 1841)
Skenea basistriata (Jeffreys, 1877)
Taranis moerchi (Malm, 1863)
Trichotropis borealis Broderip & Sowerby, 1829
Trophonopsis barvicensis (Johnston, 1841)
Turitella communis Rissö, 1826
Turridae sp. indet.
Velutina velutina (Müller, 1776)

PELECYPODA

Abra alba (Wood, 1802)
Abra nitida (Müller, 1776)
Acanthocardia echinata (Linnaeus, 1758)
Arctica islandica (Linnaeus, 1767)
Astarte elliptica (Brown, 1827)
Astarte montagui (Dillwyn, 1817)
Astarte sulcata (da Costa, 1778)
Astarte sp.
Bathyarca pectunculoides (Scacchi, 1834)
Cardiidae spp. indet.
Chlamys islandica (Müller, 1776)
Chlamys sulcata (Müller, 1776)
Corbula gibba (Olivi, 1792)
Cuspidaria rostrata (Spengler, 1793)
Cuspidaria sp.
Dosina lupinus (Linnaeus)
Ennucula tenuis (Montagu, 1818)
Gari fervensis (Gmelin, 1791)
Heteranomia squamula (Linnaeus, 1758)
Hiatella arctica (Linnaeus, 1767)
Kellia suborbicularis (Montagu, 1803)
Kellia sp.
Kelliella miliaris (Philippi, 1844)
Limatula sp.
Limatula sulcata (Brown, 1827)
Lucinoma borealis (Linnaeus, 1767)
Macoma calcarea (Gmelin, 1790)
Malletia obtusa Sars, 1869
Modiolus modiolus (Linnaeus, 1758)
Modiolula phaseolina (Philippi, 1844)
Monia squama (Gmelin, 1791)

Montacuta ferruginosa (Montagu, 1808)
Montacuta tenella Lovén, 1846
Montacuta sp.
Musculus niger (Gray, 1824)
Mya arenaria Linnaeus, 1758
Mya sp.
Mysella bidentata (Montagu, 1803)
Mysia undata (Pennant, 1777)
Mytilus edulis Linnaeus, 1758
Nuculana minuta (Müller, 1776)
Nuculana pernula (Müller, 1779)
Nucula nucleus (Linnaeus, 1758)
Nucula tumidula Malm, 1860
Nucula sp.
Paliolum furtivum (Lovén, 1846)
Paliolum vitreum (Gmelin, 1791)
Parvicardium minimum (Philippi, 1836)
Parvicardium ovale (Sowerby, 1840)
Pectinidae sp. indet.
Phaxas pellucidus (Pennant, 1777)
Pseudamussium septemradiatum (Müller, 1776)
Similipecten similis (Laskey, 1811)
Spisula elliptica (Brown, 1827)
Spisula sp.
Teredo sp.
Thracia myopsis (Beck) Møller, 1842
Thracia sp.
Thyasira spp.
Tropidomyra abbreviata (Forbes, 1843)
Venus casina Linnaeus, 1758
Yoldiella sp.

SCAPHOPODA

Dentalium entale Linnaeus
Dentalium occidentale Stimpson

Dentalium sp.
Entalina quinguangularis (Forbes)

PHORONIDÆ

Phoronis mülleri

Phoronis sp.

	ASTEROIDEA	
<u>Asteroidea</u> spp. indet.		
<u>Astropecten irregularis</u> (Pennant)		<u>Ctenodiscus crispatus</u> (Bruzzi)
		<u>Psilaster andromeda</u> (Müller and Trosch.)
	OPHIUROIDEA	
<u>Amphilepis norvegica</u> Ljungman		<u>Ophiura albida</u> Forbes
<u>Amphiura borealis</u> (G.O.Sars)		<u>Ophiura carnea</u> Sars
<u>Amphiura chiajei</u> Forbes		<u>Ophiura robusta</u> Ayres
<u>Amphiura filiformis</u> (Müller)		<u>Ophiura sarsi</u> Lütken
<u>Amphiura</u> sp.		<u>Ophiura texturata</u>
<u>Ophiopholis aculeata</u> (Müller)		Ophiura sp.
<u>Ophiura affinis</u> Lütken		Ophiuroidea spp. indet.
	ECHINOIDEA	
<u>Brisaster fragilis</u> (Düben and Koren)		<u>Echinocyamus pusillus</u> (Müller)
<u>Brissopsis lyrifera</u> (Forbes)		<u>Regularia</u> sp. indet.
<u>Echinocardium cordatum</u> (Pennant)		<u>Spatangus purpureus</u> Müller
<u>Echinocardium flavescentes</u> (Müller)		
	HOLOTHUROIDEA	
<u>Cucumaria elongata</u> Düben and Koren		Holothuroidea spp. indet.
<u>Echinocucumis hispida</u> (Barret)		
	TUNICATA	
<u>Ascidia conchilega</u> Müller, 1776		<u>Molgula kiaeri</u> Hartmeyer, 1901
<u>Ascidia</u> sp.		<u>Pelonaia corrugata</u> Forbes and Goodsir, 1841

Tables 3-32. Species lists and quantitative results for the stations.
For each taxon and cruise mean number of individuals per 0.1 m^2
(\bar{x}) and standard deviation (s) are given. There are nine degrees
of freedom. For each cruise the indices of diversity (λ , H and
 d) and sampling efficiency (S) are given.

Table 33. Values of the coefficients of similarity (cc and cz).

Table 3. Station OF 1, 20 m

	1972		1973		1974		Not sampled.	0.2	0.42	
	JUNE X	S	OCT. X	S	JUNE X	S	OCT. X	S		
<u>Paraedwardsia arenaria</u>					0.4	0.70		0.3	0.67	
<u>Chaetoderma nitidulum</u>		0.1	0.32	0.2	0.63	0.1	0.32			
Amphipoda indet.	2.3	5.17								
<u>Lunatia intermedia</u>								0.5	0.71	
<u>Nassarius reticulatus</u>	0.4	0.70						0.1	0.32	
<u>Natica</u> sp.										
Nemertini indet.		0.4	0.84	0.1	0.32	0.2	0.42		0.2	0.42
Ampharetidae indet								0.1	0.32	
<u>Anaitides</u> sp.	0.1	0.32	0.6	0.52	0.3	0.48		0.7	0.82	
Capitellidae indet.					0.1	0.32				
<u>Ceratocephale loveni</u>					0.1	0.32				
<u>Chaetozone setosa</u>		0.5	0.71	0.1	0.32			0.1	0.32	
<u>Cirratulus cirratus</u>		7.9	8.12	1.6	3.50	5.3	7.36			
<u>Diplocirrus glaucus</u>		0.5	0.97							
Eteone sp.	0.5	0.71								
Euclymeninae indet.			0.1	0.32						
<u>Glycera alba</u>	0.5	0.71			0.2	0.42	0.2	0.63	0.3	0.48
<u>Glycera lapidum</u>			0.6	0.84	0.1	0.32				
<u>Goniada maculata</u>	2.7	3.20	0.3	0.48	0.5	0.71	0.5	0.71	0.5	0.71
<u>Heteromastus filiformis</u>			0.7	0.82	0.2	0.42	0.3	0.48	2.3	3.09
<u>Lumbrineris fragilis</u>	0.1	0.32	0.1	0.32	0.2	0.42	0.2	0.42	0.5	0.71
<u>Maldane sarsi</u>			0.1	0.32	0.3	0.95				
<u>Myriochele</u> sp.	0.3	0.48			0.2	0.63				
<u>Nephtys caeca</u>	0.2	0.42								
<u>Nephtys ciliata</u>					0.2	0.63	0.9	0.99	2.3	2.11
<u>Nephtys hombergi</u>			0.3	0.48	0.6	0.84	0.1	0.32	0.2	0.42
<u>Nephtys paradoxa</u>					0.1	0.32				
<u>Neris</u> sp.							0.2	0.42		
<u>Owenia fusiformis</u>			0.1	0.32						
<u>Pectinaria auricoma</u>	0.4	0.52						0.2	0.42	
<u>Pectinaria koreni</u>	0.3	0.67	2.0	2.05	0.2	0.42	0.4	0.70		
<u>Pholoe minuta</u>	3.1	3.87	1.2	1.23	1.7	1.42	1.0	0.94	1.6	1.26
Phyllodocidae indet.						0.1	0.32		0.1	0.32
Polynoidae indet.						0.1	0.32			
<u>Pygospio elegans</u>	0.2	0.63								
<u>Scoloplos armiger</u>	41.8	38.06	35.9	16.60	30.8	14.59	30.0	11.65	69.2	25.51
<u>Sphaerodorum gracilis</u>									0.1	0.32
<u>Spio filicornis</u>	0.1	0.32			0.2	0.63				
<u>Mytilus edulis</u>	0.3	0.48	0.3	0.95			0.1	0.32		
<u>Parvicardium ovale</u>									0.1	0.32
<u>Parvicardium scabrum</u>					0.4	0.70	0.4	0.70		
<u>Thyasira</u> sp.					0.4	0.70	0.2	0.42	0.7	0.82
<u>Phoronis mülleri</u>	0.4	0.70								
Diversity:	X	0.39	0.50		0.35	0.43		0.38		
	H	1.01	1.23		1.00	1.05		1.12		
	d	2.55	2.88		3.53	2.83		5.33		
Sampling efficiency:		0.985 < S < 1	0.979 < S < 1		0.96 < S < 1	0.973 < S < 1		0.983 < S < 1		

Table 4. Station OF 2, 50 m

	1972		1973		1974	
	JUNE X	S	JUNE X	S	JUNE X	S
<u>Chaetoderma nitidulum</u>		0.3	0.67		0.67	
Amphipoda indet.	0.3	0.67				
<u>Carcinus maenas</u>			0.1	0.32		
<u>Macropipus depurator</u>			0.2	0.63		
Paguridae indet.			0.1	0.32		
<u>Buccinum</u> sp.			0.2	0.42		
<u>Lunatia intermedia</u>			0.2	0.42		
<u>Lunatia montagui</u>			0.2	0.42		
Nemertini indet.					0.1	0.32
<u>Ophiura</u> sp.			0.4	0.97		
<u>Anaitides</u> sp.	0.1	0.32	0.4	0.97	0.4	0.70
<u>Capitella capitata</u>						
<u>Chaetozone setosa</u>	0.2	0.42	1.1	1.10	1.6	1.78
<u>Cirratulus cirratus</u>			1.3	2.98		0.5
<u>Eteone</u> sp.	4.5	4.40				
<u>Glycera alba</u>						
<u>Glycera capitata</u>				0.4	0.70	0.6
<u>Glycera lapidum</u>	3.1	3.35	1.7	2.06		
<u>Goniada maculata</u>	0.3	0.48				
Hesionidae indet.						
<u>Heteromastus filiformis</u>	40.4	42.85	1.77	1.77	7.7	6.15
<u>Lumbrineris fragilis</u>			0.2	0.42	0.3	0.67
<u>Melinna cristata</u>						
<u>Myriochele</u> sp.	0.2	0.42	0.4	0.70	0.1	0.32
<u>Nephtys hombergi</u>	0.1	0.32	0.2	0.42		
<u>Nereis</u> sp.			0.1	0.32		
<u>Ophelina acuminata</u>					0.1	0.32
<u>Ophiodromus flexuosus</u>					0.6	0.70
<u>Paramphipnoma jeffreysi</u>			0.1	0.32		
<u>Pectinaria auricoma</u>					0.1	0.32
<u>Pectinaria koreni</u>			0.8	1.62	0.1	0.32
<u>Pholoe minuta</u>	0.8	1.03	0.9	0.88		
Phyllodocidae indet.					0.2	0.42
<u>Prionospio malmgreni</u>			0.1	0.32	0.3	0.67
<u>Scoloplos armiger</u>	60.1	30.76	24.7	14.92	10.7	3.53
<u>Macoma calcarea</u>						
<u>Mytilus edulis</u>			0.2	0.63		
<u>Thyasira</u> sp.	2.1	2.42	0.4	0.97	0.1	0.32
					0.32	
Diversity:	λ	0.58		0.52		0.63
	H				0.58	0.64
	d	1.10		1.47		1.26
		1.57		3.91		1.08
					1.20	1.53
					1.86	3.14
					1.88	2.58
Sampling efficiency:	0.994 < S < 1	0.969 < S < 1	0.958 < S < 1	0.985 < S < 1	0.962 < S < 1	0.985 < S < 1

Table 5. Station OF 3, 100 m

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{x}	s										
<u>Chaetoderma nitidulum</u>									0.1	0.32		
Amphipoda indet.	0.2	0.42	0.2	0.42	0.2	0.42	0.4	0.4	0.52	0.3	0.67	
<u>Carcinus maenas</u>			0.1	0.32								
<u>Brisaster fragilis</u>					0.1	0.32						
<u>Echinocardium cordatum</u>					0.2	0.42						
<u>Buccinum undatum</u>									0.1	0.32		
<u>Lunatia montagui</u>			0.3	0.67								
<u>Philine quadrata</u>			0.1	0.32								
Nemertini indet.			0.1	0.32	0.1	0.32	0.1	0.32			0.2	0.42
<u>Ophiura</u> sp.			0.1	0.32								
<u>Anaitides</u> sp.			0.1	0.32			0.1	0.32	0.8	0.79	0.5	0.97
<u>Chaetozone setosa</u>	2.2	2.25	0.6	0.84			1.8	1.55	1.8	1.75	2.0	2.58
<u>Cirratulus cirratus</u>									0.1	0.32		
Eteone sp.	0.5	0.53							1.1	1.10	0.5	0.53
<u>Glycera alba</u>												
<u>Glycera capitata</u>	0.2	0.63										
<u>Glycera lapidum</u>			0.2	0.63								
<u>Goniada maculata</u>			0.2	0.42			0.1	0.32				
<u>Goniada norvegica</u>									0.1	0.32		
Hesionidae indet.											0.1	0.32
<u>Heteromastus filiformis</u>	1.0	1.25	1.6	1.71	0.6	0.70	3.2	2.53	1.9	1.91	1.9	1.85
<u>Lumbrineris</u> sp.									0.1	0.32		
<u>Maldane sarsi</u>							0.6	1.26				
<u>Nephtys ciliata</u>	0.3	0.67	0.5	0.97	0.1	0.32	0.1	0.32	0.7	0.48	0.5	0.97
<u>Nephtys hombergi</u>	0.1	0.32			0.2	0.42	0.1	0.32	0.2	0.42		
<u>Nephtys incisa</u>			0.1	0.32					0.1	0.32		
<u>Nephtys paradoxa</u>	0.4	0.70	0.2	0.42	0.1	0.32	0.3	0.48			0.2	0.63
<u>Nereimyra punctata</u>	0.1	0.32	0.1	0.32	0.1	0.32					0.3	0.48
<u>Ophelina acuminata</u>									0.1	0.32		
<u>Paramphipnoma jeffreysi</u>	0.1	0.32	0.1	0.32								
<u>Pectinaria auricoma</u>					0.1	0.32			0.1	0.32		
<u>Pectinaria koreni</u>	0.1	0.32	0.1	0.32	0.2	0.42					0.2	0.42
<u>Pholoe minuta</u>	6.4	5.17	0.5	0.85			0.8	1.03	1.8	1.03	0.7	1.06
Phyllocoelidae indet.							0.2	0.63	0.2	0.42	0.1	0.32
Polynoidae indet.			0.1	0.32								
<u>Polyphysia crassa</u>							0.1	0.32				
<u>Praxillella praetermissa</u>	0.2	0.42	0.4	0.70								
<u>Scoloplos armiger</u>	5.5	5.04	9.2	5.07	11.9	9.63	15.2	6.37	31.7	11.20	9.9	6.28
Spionidae indet.			0.2	0.63					0.1	0.32		
<u>Montacuta tenella</u>					0.1	0.32						
<u>Thyasira</u> sp.	5.8	5.25	4.6	4.58	8.4	7.89	21.4	13.45	20.3	14.97	26.2	21.38
Diversity:	λ	0.80		0.72		0.58		0.65		0.62		0.58
	H	1.83		1.83		1.14		1.36		1.34		1.29
	d	2.57		4.16		2.40		2.46		2.65		2.30
Sampling efficiency:	0.960	$< S < 1$	0.914	$< S < 1$	0.937	$< S < 1$	0.969	$< S < 1$	0.977	$< S < 1$	0.982	$< S < 1$

Table 6. Station OF 4, 200 m

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s
<u>Stylatula elegans</u>					0.1	0.32						
<u>Chaetoderma nitidulum</u>					0.1	0.32						
Amphipoda indet.	0.5	1.27	0.1	0.32								
Nemertini indet.									0.6	0.52		
<u>Anaitides</u> sp.	0.1	0.32					0.1	0.32				
<u>Chaetozone setosa</u>	1.3	1.95	0.2	0.63	1.7	3.33	0.4	0.70	2.4	2.17		
<u>Diplocirrus glaucus</u>					0.1	0.32						
<u>Goniada maculata</u>					0.1	0.32						
<u>Harmothoe</u> sp.							0.1	0.32				
<u>Heteromastus filiformis</u>	0.4	0.84										
<u>Maldane sarsi</u>					0.1	0.32						
<u>Melinna cristata</u>					0.1	0.32			0.5	0.71		
<u>Nephtys ciliata</u>	0.3	0.67										
<u>Nephtys incisa</u>							0.1	0.32				
<u>Ophiodromus flexuosus</u>	0.1	0.32					0.1	0.32				
<u>Owenia fusiformis</u>					0.1	0.32						
Phyllodocidae indet.			0.1	0.32								
Polychaeta indet.					0.5	1.08						
<u>Scoloplos armiger</u>							0.1	0.32				
<u>Thyasira</u> sp.	0.4	1.26			0.4	0.97						
<hr/>												
Diversity:	λ	0.78		0.83		0.71		0.79		0.49		
	H	1.64		1.04		1.63		1.39		0.84		
	d	1.75		1.44		2.57		1.92		0.56		
Sampling efficiency:	0.797 < s < 1	0.575 < s < 1	0.576 < s < 1		0.150 < s < 1	0.914 < s < 1						

Not sampled.

Table 7. Station BV 1, 20 m

	JUNE		1972		OCT.		JUNE		1973		OCT.	
	\bar{X}	s	\bar{X}	s	\bar{X}	s	\bar{X}	s	\bar{X}	s	\bar{X}	s
<u><i>Stylatula elegans</i></u>							0.2		0.63			
<u><i>Chaetoderma nitidulum</i></u>	0.3	0.48	1.3		1.06		1.3		1.25	0.8		0.92
Amphipoda indet.	0.2	0.42							0.1	0.32		
Cumacea indet.									0.1	0.32		
<u><i>Brisaster fragilis</i></u>			0.1	0.32					0.1	0.32		
<u><i>Brissopsis lyrifera</i></u>			0.1	0.32					0.1	0.32		
<u><i>Echinocardium cordatum</i></u>							0.1	0.32				
<u><i>Spatangus purpureus</i></u>	0.1	0.32										
<u><i>Aporrhais pespelicanus</i></u>							0.1	0.32				
<u><i>Cylichna cylindracea</i></u>							0.1	0.32				
<u><i>Lunatia inermedia</i></u>	0.1	0.32										
<u><i>Lunatia montagui</i></u>							0.1	0.32				
<u><i>Philine quadrata</i></u>	0.1	0.32										
<u><i>Philine scabra</i></u>			0.1	0.32								
Holothuroidea indet.	26.2	15.67					4.3		2.79	1.0		1.49
Nemertini indet.			0.1	0.32	0.4		0.52	0.3		0.48		
<u><i>Amphiura chiajei</i></u>	5.0	3.23	7.6		4.09	16.8	3.19	2.9		2.23		
<u><i>Amphiura filiformis</i></u>	0.8	1.40				1.6	1.43	0.3		0.48		
<u><i>Amphiura</i> sp.</u>			0.6	0.84				0.2	0.42			
<u><i>Ophiura albida</i></u>	0.2	0.42	0.3	0.67								
Ampharetidae indet.						0.1	0.32					
<u><i>Artacama proboscidea</i></u>	0.1	0.32	0.3	0.48	0.7		0.67					
<u><i>Chaetozone setosa</i></u>	1.0	3.16										
<u><i>Cirratulus cirratus</i></u>	0.2	0.63										
<u><i>Diplocirrus glaucus</i></u>	0.3	0.48			0.1		0.32	0.1		0.32		
<u><i>Eteone</i> sp.</u>	0.1	0.32										
<u><i>Euchone papillosa</i></u>			0.1	0.32								
<u><i>Glycera alba</i></u>	1.2	1.14	0.9	0.74	0.9		0.99	0.9		0.99	0.9	0.99
<u><i>Glycera lapidum</i></u>			0.4	0.70								
<u><i>Glycera rouxii</i></u>	0.1	0.32										
<u><i>Goniada maculata</i></u>	2.4	2.37	1.1	0.99	3.8		1.81	0.5		1.27		
<u><i>Goniada norvegica</i></u>								0.3	0.67			
<u><i>Hauchiella tribullata</i></u>					0.1	0.32						
<u><i>Lanassa venusta</i></u>			0.2	0.42				0.5	0.71			
<u><i>Laonice cirrata</i></u>	0.8	1.23	0.6	0.70	1.2		0.92	0.8		0.92		
<u><i>Lumbrineris fragilis</i></u>	3.7	2.54	1.2	1.23	2.6		1.65	1.4		1.07		
<u><i>Lysilla loveni</i></u>	0.2	0.42	0.1	0.32				0.5	0.85			
<u><i>Maldane sarsi</i></u>	0.1	0.32	7.0	8.82	1.4		1.71	0.9		1.10		
<u><i>Melinna cristata</i></u>	0.4	0.84			0.6		1.58	0.1		0.32		
<u><i>Myriochele</i> sp.</u>	2.1	2.81	0.5	0.85	0.4		0.70	0.1		0.32		
<u><i>Nephtys ciliata</i></u>	0.3	0.67	0.4	0.84	0.4		0.70	0.7		0.67		
<u><i>Nephtys hombergi</i></u>	0.2	0.42			0.4		0.70	0.4		0.52		
<u><i>Nephtys incisa</i></u>	0.2	0.63	0.1	0.32								
Nereis sp.								0.1	0.32			

Table 7. cont.

Table 9. cont.

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s
<u>Owenia fusiformis</u>			0.5		1.58	0.1	0.32					
<u>Paramphinoe jeffreysi</u>	0.2	0.42	0.1		0.32							
<u>Pectinaria helgica</u>			0.1		0.32				0.1	0.32		
<u>Pholoe minuta</u>	3.4	9.05	0.2		0.63	0.1	0.32	0.3	0.48			
Phyllodocidae indet.								0.5	0.71			
<u>Pista cristata</u>			0.1		0.32							
<u>Polymnia nebulosa</u>								0.2	0.42			
Polynoidae indet.			0.2		0.42	0.1	0.32	0.3	0.67			
<u>Polyphysia crassa</u>	12.9	6.92	11.0		6.13	7.5	3.92	7.7	3.06			
<u>Praxillella praetermissa</u>			0.2		0.42	0.1	0.32					
<u>Scoloplos armiger</u>	1.5	2.84				0.2	0.42					
<u>Streblosoma bairdi</u>	3.2	1.62	2.9		3.25	2.6	1.51	2.5	2.22			
Syllidae indet.	0.1	0.32										
<u>Trichobranchus roseus</u>	0.2	0.42										
<u>Cuspidaria</u> sp.	0.1	0.32										
<u>Kelliella miliaris</u>			0.9		1.91							
<u>Montacuta tenella</u>						0.1	0.32	0.1	0.32			
<u>Mya</u> sp.	0.1	0.32				0.1	0.32					
<u>Mytilus edulis</u>			0.1		0.32							
<u>Parvicardium minimum</u>			0.3		0.48							
<u>Thyasira</u> sp.	10.4	9.61	27.1		13.60	23.4	13.36	20.9	8.63			
<u>Phoronis mülleri</u>			0.2		0.42							
<u>Onchnesoma steenstrupi</u>			0.2		0.42			0.1	0.32			
Diversity:	λ	0.88		0.80		0.76		0.79				
	H	2.51		2.24		2.13		2.09				
	d	5.03		5.96		5.60		4.78				
Sampling efficiency:		0.973 < s < 1		0.971 < s < 1		0.963 < s < 1		0.972 < s < 1				

Table 10. Station TH 1, 50 m

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{X}	s										
<u>Paraedwardsia arenaria</u>									0.1	0.32		
<u>Virgularia mirabilis</u>											0.1	0.32
<u>Ctenodiscus crispatus</u>											0.1	0.32
<u>Chaetoderma nitidulum</u>	1.0	1.25	0.6	0.84	0.9	0.99	0.1	0.32			0.2	0.42
Amphipoda indet.	0.8	1.40	2.4	1.65	0.3	0.67	1.3	0.82			0.8	1.03
Cumacea indet.	0.4	0.70	7.0	7.57	1.2	1.14	6.4	4.20	0.3	0.48	2.1	2.28
<u>Galathea</u> sp.			0.1	0.32								
<u>Gnathia oxyraea</u>					0.1	0.32						
<u>Hyas coarctatus</u>			0.2	0.42								
<u>Macropipus depurator</u>									0.1	0.32		
<u>Echinocardium cordatum</u>	0.2	0.42					0.1	0.32			0.2	0.42
<u>Aporrhais pespellicani</u>	0.4	0.70					0.3	0.48	0.2	0.63		
<u>Cylichna cylindracea</u>					0.3	0.95	0.1	0.32				
<u>Lunatia montagui</u>					0.1	0.32	0.1	0.32	0.1	0.32		
<u>Philine quadrata</u>	0.3	0.95					0.2	0.63	0.1	0.32		
<u>Philine scabra</u>			0.8	1.32	0.2	0.63						
<u>Philine</u> sp.	0.4	0.97										
<u>Scaphander lignarius</u>			0.1	0.32			0.1	0.32			0.1	0.32
<u>Scaphander punctostriatus</u>	0.1	0.32										
<u>Taranis moerchi</u>	0.2	0.63			0.1	0.32						
<u>Velutina velutina</u>									0.1	0.32		
<u>Echinocucumis hispida</u>					0.1	0.32						
<u>Holothuroidea</u> indet.			0.1	0.32	0.1	0.32	0.5	1.08	0.2	0.42	0.1	0.32
Nemertini indet.			1.3	0.82	0.9	0.57	1.4	0.84	0.4	0.70	1.0	0.67
<u>Amphilepis norvegica</u>					0.3	0.67			0.6	0.84		
<u>Amphiura chiajei</u>					0.2	0.42	0.2	0.42			0.2	0.42
<u>Amphiura filiformis</u>	0.1	0.32	0.3	0.48			0.1	0.32			0.4	0.52
Amphiura sp.			0.2	0.42	0.1	0.32						
<u>Ophiura albida</u>	0.7	0.67	0.3	0.48			1.2	0.79			0.1	0.32
Ophiuroidea indet.	0.5	0.85										
<u>Ophiura sarsi</u>			3.9	2.56	1.2	1.99					0.7	0.95
<u>Ophiura</u> sp.			0.9	1.10	1.5	1.96			0.2	0.42	0.5	0.53
<u>Ophiura texturata</u>			0.1	0.32								
<u>Ampharete finmarchica</u>			0.1	0.32			0.1	0.32				
Ampharetidae indet.	0.4	0.70	0.1	0.32			0.3	0.48			1.3	1.64
<u>Amphicteis gunneri</u>			0.1	0.32					0.1	0.32	0.1	0.32
Anaitides sp.	0.1	0.32	0.1	0.32			0.1	0.32	0.1	0.32	0.2	0.63
<u>Anobothrus gracilis</u>					0.4	0.70						
<u>Aphrodita aculeata</u>	0.2	0.42			0.1	0.32						
<u>Artacama proboscidea</u>							0.1	0.32	0.1	0.32		

Table 10. cont.

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s
<u>Asychis biceps</u>					0.2	0.42						
<u>Brada villosa</u>	0.3	0.48										
<u>Chaetozone setosa</u>	0.1	0.32			0.1	0.32			0.1	0.32	0.3	0.67
<u>Cirratulus cirratus</u>	0.2	0.63	0.3	0.67			0.3	0.67				
<u>Dasybranchus caducus</u>							0.1	0.32				
<u>Diplocirrus glaucus</u>	0.1	0.32	0.8	0.79	0.4	0.70	1.3	1.06	1.0	1.63	1.4	1.26
<u>Eclysippe vanelli</u>	0.1	0.32										
<u>Euchone papillosa</u>					0.1	0.32						
<u>Euchone rubrocincta</u>		0.5	1.27	0.2	0.63	0.1	0.32					
<u>Euclymeninae</u> indet.	0.2	0.42							0.1	0.32		
<u>Eulalia</u> sp.			0.1	0.32								
<u>Eunice pennata</u>					0.1	0.32					0.3	0.48
<u>Genethyllis lutea</u>	0.2	0.42										
<u>Glycera alba</u>							0.1	0.32	0.1	0.32	0.6	0.84
<u>Glycera rouxii</u>	0.5	0.71	0.2	0.42	1.2	0.63	0.9	1.10	0.9	0.74		
<u>Glycera</u> sp.			0.6	0.84								
<u>Goniada maculata</u>	0.6	0.70	0.5	0.71	0.7	0.95	1.0	1.05	0.5	0.71	0.4	0.70
<u>Harmothoe</u> sp.			0.2	0.42								
<u>Heteromastus filiformis</u>	0.7	0.82			0.7	1.49			0.2	0.42		
<u>Hyalinoecia tubicola</u>					0.1	0.32						
<u>Laetmonice filicornis</u>									0.1	0.32		
<u>Lanassa venusta</u>	0.1	0.32									0.1	0.32
<u>Laonice cirrata</u>	0.6	0.70	1.8	1.55			2.1	2.18	1.0	0.67	2.3	1.57
<u>Laphania boecki</u>	0.4	0.97			0.1	0.32	0.1	0.32	0.1	0.32		
<u>Leanira tetragona</u>			0.1	0.32								
<u>Lumbrineris fragilis</u>	0.6	0.52	0.4	0.70	0.6	0.70	0.3	0.48	0.6	0.84	0.3	0.48
<u>Maldanidae</u> indet.			0.1	0.32			0.2	0.63				
<u>Maldane sarsi</u>			0.4	0.97	0.3	0.67	1.2	1.62	0.3	0.95	0.6	0.84
<u>Melinna cristata</u>	35.1	17.65	6.5	4.95	68.5	19.28	3.1	2.18	8.5	4.22	15.0	14.24
<u>Myriochele</u> sp.	0.3	0.67	0.3	0.48	0.3	0.67	0.6	0.97	0.1	0.32	0.1	0.32
<u>Nephtys ciliata</u>	0.1	0.32	0.3	0.48			0.9	1.20	0.3	0.67		
<u>Nephtys hombergi</u>	0.8	1.14	0.1	0.32	0.1	0.32	0.1	0.32			0.3	0.48
<u>Nephtys paradoxa</u>			0.1	0.32							0.1	0.32
<u>Nereimyra punctata</u>			0.1	0.32			0.1	0.32	0.1	0.32		
<u>Nereis</u> sp.							0.1	0.32				
<u>Notomastus latericeus</u>	0.7	0.48	0.9	0.99	1.0	0.67	1.5	1.90	1.3	1.34	2.0	1.56
<u>Ophelina norvegica</u>			0.1	0.32								
<u>Ophiodromus flexuosus</u>	0.4	0.70	0.3	0.67	0.2	0.42	0.1	0.32			0.6	0.70
<u>Owenia fusiformis</u>	1.0	1.33	0.1	0.32	0.2	0.42	0.3	0.48	0.2	0.42		
<u>Paramphitone jeffreysi</u>	0.1	0.32										
<u>Paranis gracilis</u>					0.1	0.32						
<u>Pectinaria auricoma</u>	1.3	2.11	0.6	1.26	0.4	0.70	0.3	0.48	0.4	0.52	0.2	0.42
<u>Pectinaria koreni</u>	0.9	1.60	0.9	0.88	0.5	1.08	0.6	0.84	0.2	0.42	0.1	0.32
<u>Pholoe minuta</u>	2.6	2.07	5.3	2.87	0.9	0.99	6.2	4.32	0.5	0.71	2.7	2.36
<u>Phyllodocidae</u> indet.					0.1	0.32	0.1	0.32			0.2	0.42
<u>Pista cristata</u>	0.8	1.03	1.1	0.88	0.4	0.70	0.8	1.14	0.1	0.32	1.3	1.06
<u>Polycirrus medusa</u>			0.1	0.32								
<u>Polycirrus</u> sp.									0.3	0.95		
<u>Polynoidae</u> indet.	0.1	0.32			0.1	0.32	0.1	0.32	0.4	0.70	0.3	0.48
<u>Polyphysia crassa</u>	0.5	1.08	0.3	0.48	0.4	0.97	0.4	0.70			0.1	0.32
<u>Praxillella praetermissa</u>			0.2	0.42								
<u>Rhodine gracilior</u>											0.1	0.32
<u>Sabellides octocirrata</u>			0.3	0.48								
<u>Sabella penicilllus</u>			0.1	0.32	0.1	0.32						
<u>Scalibregma inflatum</u>			0.1	0.32								
<u>Scoloplos armiger</u>					0.1	0.32	0.1	0.32				

Table 10, cont.

	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s
<u>Sphaerodorum gracilis</u>			0.1	0.32	0.1	0.32	0.3	0.67	0.1	0.32		
Spionidae indet.			0.8	1.03	0.3	0.67	0.9	0.99		0.2	0.42	
<u>Streblosoma bairdi</u>	0.5	0.85	0.9	0.74	0.2	0.63	1.0	1.70	0.4	0.70	1.0	0.94
Syllidae indet.	0.2	0.42	0.3	0.67	1.2	1.14	0.2	0.42	0.4	0.52	0.2	0.63
Terebelliidae indet.	0.2	0.63					0.2	0.63				
<u>Terebellides stroemii</u>												
<u>Thelepus cincinnatus</u>	0.2	0.42	0.1	0.32	0.1	0.32	0.2	0.42	0.2	0.42	0.8	0.79
<u>Trichobranchus roseus</u>	0.2	0.42	0.3	0.67	0.1	0.32	1.4	1.17	0.2	0.42	0.6	0.70
<u>Abra alba</u>	0.3	0.67	0.2	0.42	2.2	2.57	6.8	3.33	0.8	6.39	3.6	1.65
<u>Abra nitida</u>	0.8	1.03	0.5	0.53	0.1	0.32	1.9	2.69	5.8	3.52	2.5	1.90
<u>Arctica islandica</u>					0.3	0.48						
<u>Astarte</u> sp.						0.1	0.32					
<u>Astarte sulcata</u>						0.2	0.42					
<u>Bathyarca pectunculoides</u>			0.1	0.32								
Cardiidae indet.	1.4	1.65									0.1	0.32
<u>Corbula gibba</u>												
<u>Cuspidaria</u> sp.	0.1	0.32			0.2	0.42					0.1	0.32
<u>Dosinia lupinus</u>									0.1	0.32		
<u>Ennucula tenuis</u>	0.1	0.32					0.1	0.32				
<u>Kellia</u> sp.	0.1	0.32										
<u>Kelliella miliaris</u>			0.6	0.97	0.2	0.42						
<u>Limatula</u> sp.	0.3	0.95									1.2	1.32
<u>Limatula sulcata</u>			4.1	3.21	1.5	1.58	0.7	1.06				
<u>Lucinoma borealis</u>									0.1	0.32		
<u>Macoma calcarea</u>	0.1	0.32			0.1	0.32	0.1	0.32	0.1	0.32	0.1	0.32
<u>Modiolula phaseolinus</u>					0.2	0.42						
<u>Monia squama</u>									0.1	0.32		
<u>Mysella bidentata</u>		0.1	0.32									
<u>Nuculana minuta</u>					1.3	1.25	0.4	0.52	0.2	0.42		
<u>Nuculana pernula</u>									0.4	0.52		
<u>Parvicardium minimum</u>			1.7	1.25	0.2	0.42	1.2	1.32	0.3	0.67	0.7	0.67
<u>Pseudamussium septemradiatum</u>										0.2	0.63	
<u>Thyasira</u> sp.	18.1	10.20	30.1	13.95	16.0	10.99	5.9	3.67	3.3	3.27	7.3	5.58
<u>Tropidomya abbreviata</u>					0.1	0.32						
<u>Yoldiella</u> sp.	0.7	0.95	2.2	2.15	6.1	5.15	0.5	0.85	0.6	0.97	1.1	1.52
<u>Phoronis mülleri</u>			0.1	0.32								
<u>Dentalium entale</u>			0.1	0.32					0.1	0.32		
<u>Dentalium occidentale</u>												
<u>Onchnesoma steenstrupi</u>	0.2	0.63	0.1	0.32	0.1	0.32					0.1	0.32
<u>Phascolion strombi</u>	0.8	0.92	0.1	0.32	0.3	0.48						
<u>Molgula kiaeri</u>							0.3	0.67				
Diversity:	λ	0.75		0.85		0.64		0.94		0.88		0.90
	H	2.29		2.81		1.93		3.32		2.76		3.00
	d	8.84		10.38		9.76		10.06		8.77		8.35
Sampling efficiency:	0.973 < s < 1	0.956 < s < 1	0.969 < s < 1	0.944 < s < 1	0.929 < s < 1	0.962 < s < 1						

Table 11. Station TH 2, 120 m

	1974	
	JUNE	
	\bar{x}	s
<u>Paraedwardsia arenaria</u>	0.1	0.32
<u>Stylatula elegans</u>	0.1	0.32
<u>Virgularia mirabilis</u>	0.1	0.32
<u>Chaetoderma nitidulum</u>	0.4	0.52
Amphipoda indet.	0.4	0.52
<u>Calocharis macandreae</u>	0.8	0.63
Nemertini indet.	0.3	0.48
<u>Amphilepis norvegica</u>	2.7	2.26
<u>Amphiura chiajei</u>	0.3	0.48
<u>Ophiura</u> sp.	0.6	0.70
<u>Asychis biceps</u>	0.1	0.32
<u>Chaetozone setosa</u>	0.5	0.85
Euclymeninae indet.	1.2	1.32
<u>Glycera alba</u>	0.1	0.32
<u>Lumbrineris fragilis</u>	0.3	0.48
<u>Melinna cristata</u>	0.3	0.48
<u>Nephtys incisa</u>	1.0	0.67
<u>Nephtys paradoxa</u>	0.2	0.42
<u>Onuphis quadricuspis</u>	0.1	0.32
<u>Phylo norvegicus</u>	0.1	0.32
<u>Pista cristata</u>	0.1	0.32
Polychaeta indet.	0.2	0.63
<u>Polyphysia crassa</u>	0.4	0.70
<u>Rhodine gracilior</u>	0.1	0.32
<u>Streblosoma bairdi</u>	0.1	0.32
<u>Thelepus cincinnatus</u>	2.6	2.95
<u>Limatula sulcata</u>	0.2	0.63
<u>Thyasira</u> sp.	0.1	0.32
<u>Onchnesoma steenstrupi</u>	0.1	0.32
	λ	0.91
Diversity:	H	2.74
	d	5.70
Sampling efficiency:	0.859 < s < 1	

Table 12. Station TH 3, 50 m

	1974 JUNE	
	\bar{x}	s
<u><i>Virgularia mirabilis</i></u>	0.1	0.32
<i>Amphipoda</i> indet.	0.2	0.63
<u><i>Calocharis macandreae</i></u>	0.1	0.32
<u><i>Brisaster fragilis</i></u>	0.1	0.32
<i>Holothuroidea</i> indet.	1.4	1.17
<i>Nemertini</i> indet.	0.3	0.67
<u><i>Amphilepis norvegica</i></u>	0.1	0.32
<u><i>Ophiura</i> sp.</u>	0.3	0.48
<u><i>Asychnis biceps</i></u>	0.6	1.07
<u><i>Dasybranchus caducus</i></u>	0.2	0.42
<u><i>Ditrupa arietina</i></u>	0.1	0.32
<u><i>Euchone rubrocincta</i></u>	0.1	0.32
<i>Euclymeninae</i> indet.	0.1	0.32
<u><i>Glycera rouxi</i></u>	0.3	0.48
<u><i>Leanira tetragona</i></u>	0.3	0.48
<u><i>Lumbrineris fragilis</i></u>	0.1	0.32
<u><i>Melinna cristata</i></u>	54.9	13.67
<u><i>Myriochele</i> sp.</u>	0.4	0.52
<u><i>Nephtys incisa</i></u>	0.1	0.32
<u><i>Nothria conchylega</i></u>	0.1	0.32
<u><i>Pectinaria auricoma</i></u>	0.1	0.32
<i>Syllidae</i> indet.	0.2	0.42
<u><i>Thelepus cincinnatus</i></u>	0.1	0.32
<u><i>Astarte sulcata</i></u>	1.0	0.82
<u><i>Kelliella miliaris</i></u>	0.1	0.32
<u><i>Parvicardium minimum</i></u>	0.1	0.32
<u><i>Thyasira</i> sp.</u>	1.5	3.41
<i>Porifera</i> indet.	0.1	0.32
<u><i>Dentalium entale</i></u>	0.1	0.32
	λ	0.24
Diversity:	H	0.76
	d	4.34
Sampling efficiency:	0.962 < s < 1	

Table 13. Station SF 1, 200 m

	1972				1973			
	JUNE		OCT.		JUNE		OCT.	
	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	
<i>Asterioidea</i> indet.					0.1	0.32		
<i>Psilaster andromeda</i>					0.1	0.32		
<i>Chaetoderma nitidulum</i>	0.2	0.42			0.1	0.32	0.1	0.32
Amphipoda indet.	1.1	2.33	0.5	1.27	1.5	2.07		
<i>Calocharis macandreae</i>	0.7	0.67	0.6	0.70	0.6	0.52		
<i>Philine scabra</i>					0.1	0.32		
Nemertini indet.		0.4		0.52	0.6	0.84	0.3	0.48
<i>Amphilepis norvegica</i>	13.2	2.49	10.3	7.20	10.7	5.87		
<i>Amphiura chiajei</i>			0.1	0.32				
<i>Ophiura</i> sp.					0.1	0.32		
Ampharetidae indet.					0.1	0.32		
<i>Asychis biceps</i>	0.9	1.52	1.8	1.14	1.5	1.18	1.0	1.63
<i>Chaetozone setosa</i>			0.1	0.32	0.2	0.42	0.1	0.32
<i>Dasybranchus caducus</i>	0.3	0.48	0.2	0.63	0.3	0.48	0.6	0.70
<i>Diplocirrus glaucus</i>			0.3	0.48	0.1	0.32		
<i>Diplocirrus longisetosa</i>			0.1	0.32				
Euclymeninae indet.							0.1	0.32
<i>Eunice pennata</i>	0.1	0.32						
<i>Heteromastus filiformis</i>	0.7	1.57	0.1	0.32	0.1	0.32	0.3	0.48
<i>Laonice cirrata</i>	0.2	0.42						
<i>Lumbrineris fragilis</i>	3.1	2.51	1.2	1.23	1.9	2.28	1.8	1.23
<i>Melinna cristata</i>					1.7	3.50	0.1	0.32
<i>Nephtys ciliata</i>	0.1	0.32						
<i>Nephtys incisa</i>	0.2	0.42			0.4	0.70	0.4	0.52
<i>Nephtys paradoxa</i>	0.1	0.32			0.1	0.32		
<i>Onuphis quadricuspis</i>			0.1	0.32				
<i>Ophelina norvegica</i>	0.3	0.48			0.3	0.48	0.1	0.32
<i>Paramphinome jeffreysi</i>	0.2	0.42	0.1	0.32				
<i>Panthalis cerstedi</i>	0.1	0.32	0.1	0.32				
<i>Pholoe minuta</i>					0.1	0.32		
<i>Phylo norvegicus</i>	0.5	0.71	1.0	0.82	0.8	1.03	0.6	0.52
<i>Pista cristata</i>	0.1	0.32						
<i>Polycirrus</i> sp.	0.3	0.48						
Polynoidae indet.	0.1	0.32						
<i>Polyphysia crassa</i>	0.3	0.95	0.5	0.71	0.3	0.48	0.5	0.85
<i>Rhodine loveni</i>	1.4	1.78	0.6	0.70	0.7	0.67	0.5	0.53
Sabellidae indet.			0.3	0.67				
<i>Scoloplos armiger</i>	0.1	0.32	0.1	0.32			0.1	0.32
<i>Terebellides stroemi</i>	0.2	0.42					0.1	0.32

Table 13. cont.

	1972				1973			
	JUNE		OCT.		JUNE		OCT.	
	\bar{X}	s	\bar{X}	s	\bar{X}	s	\bar{X}	s
<u>Abra alba</u>					0.1	0.32		
<u>Abra nitida</u>			0.2	0.42	0.1	0.32		
<u>Cuspidaria</u> sp.	0.1	0.32						
<u>Kelliella miliaris</u>			0.3	0.48	0.1	0.32	0.1	0.32
<u>Malletia obtusa</u>					0.1	0.32		
<u>Nucula</u> sp.	0.1	0.32						
<u>Nucula tumidula</u>			0.1	0.32	0.1	0.32		
<u>Thyasira</u> sp.	0.6	0.70	1.4	1.84	0.3	0.48		
<u>Tropidomya abbreviata</u>			0.2	0.42				
<u>Yoldiella</u> sp.	0.3	0.48	0.1	0.32	0.3	0.95		
<u>Dentalium occidentale</u>					0.1	0.32		
<u>Entalina quinquangularis</u>			0.2	0.42	0.6	0.97		
<u>Onchnesoma steenstrupi</u>	2.2	1.87	2.1	1.29	2.8	1.87	0.9	1.20
<u>Phascolosoma</u> sp.	2.0	2.21	0.1	0.32				
Diversity:	λ	0.78		0.78		0.81		0.90
	H	2.24		2.25		2.41		2.47
	d	5.09		5.14		5.89		3.91
Sampling efficiency:	0.946 < s < 1		0.922 < s < 1		0.911 < s < 1		0.805 < s < 1	

Table 14. Station SF 2, 100 m

Table 14. cont.

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s
<u>Nephtys incisa</u>					0.7	0.67					0.1	0.32
<u>Nephtys paradoxa</u>	0.3	0.48	0.7	0.95	0.4	0.52	0.1	0.32			0.1	0.32
<u>Nephtys</u> sp.											0.1	0.32
<u>Nereis</u> sp.					0.1	0.32						
<u>Onuphis quadricuspis</u>					0.4	0.70						
<u>Ophelina acuminata</u>	0.2	0.63										
<u>Ophelina norvegica</u>					0.3	0.67					0.4	0.52
<u>Ophiodromus flexuosus</u>	0.1	0.32									0.1	0.32
<u>Panthalis oerstedi</u>											0.1	0.32
<u>Paramphinome jeffreysi</u>	0.1	0.32			0.1	0.32						
<u>Pectinaria auricoma</u>	0.2	0.42	0.1	0.32	0.2	0.42					0.1	0.32
<u>Pholoe minuta</u>	0.4	0.84			0.1	0.32						
<u>Phylo norvegicus</u>			0.3	0.67	0.3	0.67					0.6	0.97
Polynoidae indet.							0.3	0.48			0.2	0.42
<u>Polyphysia crassa</u>	0.1	0.32	0.5	0.85	0.3	0.48	0.3	0.48			0.3	0.67
<u>Rhodine loveni</u>					0.3	0.67					0.1	0.32
<u>Scoloplos armiger</u>	0.5	1.27										
Spionidae indet.					0.1	0.32						
<u>Spiophanes krøyeri</u>	0.3	0.67									0.1	0.32
Thelepinæ indet.												
<u>Abra alba</u>											0.5	0.97
<u>Abra nitida</u>	1.0	1.25	6.0	3.68	4.5	2.76	0.8	1.32			7.5	8.92
<u>Astarte sulcata</u>				0.1	0.32		0.2	0.42				
<u>Bathyarca pectunculoides</u>	0.9	0.88	0.5	0.71	0.3	0.48						
Cardiidae indet.	0.8	0.79										
<u>Cuspidaria</u> sp.	0.7	0.67	0.1	0.32								
<u>Ennucula tenuis</u>				0.1	0.32							
<u>Kellia</u> sp.	0.5	0.71										
<u>Kelliella miliaris</u>			3.5	3.81	0.6	0.97					0.1	0.32
<u>Limatula</u> sp.	0.1	0.32										
<u>Limatula sulcata</u>			0.1	0.32	0.4	0.52	0.2	0.42			0.2	0.63
<u>Parvicardium minimum</u>			0.4	0.52							0.2	0.42
<u>Pseudamussium septemradiatum</u>							0.1	0.32				
<u>Thyasira</u> sp.	4.5	2.92	11.5	11.00	6.5	5.15	0.2	0.42			2.2	3.68
<u>Tropidomya abbreviata</u>			0.5	0.71								
<u>Yoldiella</u> sp.	1.6	1.65	2.8	1.81	1.9	1.73					0.3	0.48
<u>Phoronis</u> sp.	0.7	1.34										
<u>Dentalium entale</u>			0.1	0.32								
<u>Dentalium occidentale</u>					0.2	0.63						
<u>Entalina quinquangularis</u>					0.6	0.84						
<u>Onchesoma steenstrupi</u>	17.1	11.68	37.0	14.41	18.2	9.81	0.5	0.97			0.2	0.42
<hr/>												
Diversity:	λ	0.85		0.75		0.81		0.91			0.86	
	H	2.70		2.16		2.44		2.86			2.60	
	d	6.61		5.69		6.84		6.46			6.93	
Sampling efficiency:	0.970 < S < 1	0.977 < S < 1	0.964 < S < 1	0.804 < S < 1					0.895 < S < 1			

Table 15. Station SD 1. 20 m

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{X}	s										
<i>Anthozoa</i> indet.							0.1	0.32				
<u><i>Paraedwardsia arenaria</i></u>			0.1	0.32					0.1	0.32		
<u><i>Virgularia mirabilis</i></u>					0.1	0.32	0.3	0.48				
<i>Asteroidea</i> indet.			0.4	1.26	0.1	0.32						
<u><i>Ctenodiscus crispatus</i></u>	0.4	0.52	0.3	0.48	0.6	0.52	0.3	0.67	0.7	0.67	0.6	0.84
<u><i>Chaetoderma nitidulum</i></u>			0.5	0.71	1.3	1.89	1.2	1.40	0.4	0.97	0.1	0.32
<i>Amphipoda</i> indet.	1.4	1.43			0.1	0.32	0.2	0.42			0.1	0.32
<u><i>Calocharis macandreae</i></u>	0.1	0.32										
<i>Cumacea</i> indet.	0.1	0.32					0.1	0.32				
<u><i>Brisaster fragilis</i></u>	0.1	0.32	0.1	0.32								
<u><i>Brissopsis lyrifera</i></u>			0.1	0.32								
<u><i>Echinocardium cordatum</i></u>	0.2	0.42	0.1	0.32	0.9	0.74	0.1	0.32	0.3	0.67	0.3	0.48
<u><i>Echinocyamus pusillus</i></u>					0.1	0.32						
<u><i>Actaeon tornatilis</i></u>	0.1	0.32			0.2	0.42	0.1	0.32				
<u><i>Aporrhais pespellicani</i></u>	0.2	0.42	0.3	0.48					0.2	0.42		
<u><i>Cylichna alba</i></u>							0.1	0.32				
<u><i>Cylichna cylindracea</i></u>	0.2	0.42	0.2	0.42	1.0	0.82	0.2	0.42			0.2	0.42
<u><i>Lunatia intermedia</i></u>	0.1	0.32	0.1	0.32	0.1	0.32	0.1	0.32			0.1	0.32
<u><i>Philine scabra</i></u>					0.3	0.48	0.1	0.32				
<u><i>Turitella communis</i></u>	0.4	0.70			0.1	0.32	0.1	0.32				
<u><i>Cucumaria elongata</i></u>			0.1	0.32								
Holothuroidea indet.	0.9	1.45	0.4	0.70	1.2	1.14	2.9	2.85	1.3	0.95	0.3	0.48
Nemertini indet.			1.2	1.03	1.6	1.07	2.2	1.93	0.7	0.82	1.2	1.32
<u><i>Amphilepis norvegica</i></u>											0.3	0.48
<u><i>Amphiura chiajei</i></u>	3.2	1.99	6.4	2.37	4.0	3.02	3.3	2.26	2.9	1.73	5.9	2.60
<u><i>Amphiura filiformis</i></u>	3.0	2.21	1.6	0.97	4.5	2.32	3.2	2.15	1.4	1.17	2.6	1.96
<u><i>Ophiura affinis</i></u>			.					0.4	0.70			
<u><i>Ophiura albida</i></u>	3.0	1.63	1.6	1.07			0.3	0.95			0.2	0.42
<u><i>Ophiura robusta</i></u>					0.1	0.32						
<u><i>Ophiura sarsi</i></u>			0.1	0.32	0.3	0.67					0.1	0.32
<u><i>Ophiura</i> sp.</u>					0.8	1.32	0.3	0.67	0.7	1.16		
<u><i>Ophiura texturata</i></u>			0.2	0.42					0.3	0.48		
Ampharetidae indet.	0.1	0.32			0.1	0.32						
<u><i>Amphitrite cirrata</i></u>			0.1	0.32					0.2	0.42	0.1	0.32
Anaitides sp.	0.6	0.70	0.2	0.42			0.1	0.32			0.2	0.42
<u><i>Asychis biceps</i></u>									0.1	0.32		
Capitellidae indet.									0.1	0.32		
<u><i>Chaetozone setosa</i></u>							0.1	0.32				
<u><i>Diplocirrus glaucus</i></u>	3.2	2.39	0.1	0.32	3.6	2.91	2.2	1.40	1.3	2.16	0.5	0.71
<u><i>Eteone</i> sp.</u>			0.1	0.32					0.1	0.32		
<u><i>Euchone papillosa</i></u>							0.1	0.32				

Table 15. cont.

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s
<u>Gattyana cirrosa</u>			0.1		0.32							
<u>Glycera alba</u>	1.1	1.10	0.9	1.45	1.0	1.15	1.1	1.20	0.7	0.95	0.4	0.70
<u>Glycera rouxii</u>	0.3	0.95	0.8	1.03	1.2	1.40	1.7	1.77	0.7	0.67	1.4	0.97
<u>Goniada maculata</u>	2.3	2.31	0.7	0.48	3.3	2.71	2.9	1.10	1.5	1.72	0.7	0.82
<u>Heteromastus filiformis</u>	0.1	0.32										
<u>Hydroides norvegica</u>	0.1	0.32										
<u>Lanassa nordenskiöldi</u>									0.1	0.32		
<u>Lanassa venusta</u>	0.1	0.32										
<u>Laonice cinnata</u>	1.3	1.06	0.3	0.67	1.3	1.06	1.2	1.03	1.6	1.07	0.7	0.67
<u>Lumbrineris fragilis</u>	1.3	1.25	0.9	0.57	1.4	1.07	1.3	1.16	1.3	2.00	1.1	1.29
<u>Lysilla loveni</u>									0.1	0.32		
<u>Maldane sarsi</u>	11.8	6.43	12.7	10.45	27.3	27.12	25.9	24.46	17.5	10.38	5.5	3.37
<u>Melinna cristata</u>									0.6	0.70		
<u>Myriochela sp.</u>	11.0	6.48	5.0	5.98	4.3	3.86	7.7	8.99	1.6	0.84	1.2	1.40
<u>Neoamphitrite grayi</u>										0.1	0.32	
<u>Nephtys ciliata</u>					0.1	0.32	0.2	0.42	0.1	0.32	0.1	0.32
<u>Nephtys hombergi</u>	0.4	0.52							0.1	0.32		
<u>Nephtys incisa</u>									0.1	0.32	0.1	0.32
<u>Nephtys sp.</u>			0.3	0.48								
<u>Nereis virens</u>								0.1	0.32			
<u>Ophelina acuminata</u>	0.1	0.32						0.1	0.32	0.1	0.32	
<u>Ophelina sp.</u>					0.1	0.32						
<u>Ophiadromus flexuosus</u>	0.2	0.42			0.1	0.32	0.2	0.42	0.1	0.32		
<u>Owenia fusiformis</u>	0.6	0.84			0.1	0.32	0.4	1.26				
<u>Pectinaria auricoma</u>	0.6	0.97			1.4	1.51	1.0	0.67	0.6	0.70	0.1	0.32
<u>Pectinaria koreni</u>	1.1	1.20	0.5	0.53	1.7	1.64	1.9	1.52	1.0	0.47		
<u>Pholoe minuta</u>								0.2	0.42	0.1	0.32	
<u>Pista cristata</u>									0.1	0.32	0.1	0.32
<u>Polycirrus sp.</u>	0.1	0.32							0.1	0.32		
<u>Polynmnia nebulosa</u>					0.1	0.32						
<u>Polynoidae indet.</u>	0.1	0.32	0.1	0.32	0.2	0.42	0.3	0.48	0.1	0.32		
<u>Polyphysia crassa</u>	0.1	0.32			0.5	0.53	0.6	0.97				
<u>Praxillella praetermissa</u>	0.2	0.63	1.0	1.05	0.6	0.70	1.8	3.68	0.1	0.32	0.1	0.32
<u>Sabellina penincillatus</u>								0.1	0.32			
<u>Scalibregma inflatum</u>	0.1	0.32						0.2	0.63			
<u>Scoloplos armiger</u>	0.2	0.42	0.1	0.32	0.2	0.63	0.1	0.32				
<u>Sphaerodororum gracilis</u>	0.1	0.32										
<u>Spi filicornis</u>	0.1	0.32										
<u>Spionidae indet.</u>			0.2	0.42	0.2	0.42	0.4	0.52	0.9	0.99	0.6	0.52
<u>Spiophanes krøyeri</u>	1.2	1.03										
<u>Streblosoma bairdi</u>	0.2	0.63			0.2	0.42				0.1	0.32	
<u>Terebellidae indet.</u>									0.1	0.32		
<u>Terebellides stroemii</u>	0.1	0.32	0.2	0.42	0.1	0.32			0.8	0.92	0.1	0.32
<u>Trichobranchus roseus</u>	0.3	0.67	0.1	0.32	0.7	0.82	1.0	0.82	1.4	1.51	0.8	1.23
<u>Trochochaeta multisetosa</u>					0.1	0.32						
<u>Abra alba</u>	0.1	0.32			39.9	27.22	24.3	30.83	2.7	4.06	1.0	1.49
<u>Abra nitida</u>	1.1	2.47	0.1	0.32	10.9	7.25	8.0	9.85	0.9	1.91	0.7	1.64
<u>Acanthocardia echinata</u>			0.1	0.32			0.3	0.48			0.3	0.67
<u>Arctica islandica</u>	1.3	0.82	1.0	0.82	2.3	1.16	0.7	0.67	0.1	0.32	0.6	0.70
<u>Cardiidae indet.</u>	0.9	1.10										
<u>Corbula gibba</u>	2.6	2.95	0.4	0.52	1.4	1.65	1.6	2.17	0.5	0.53	2.8	2.39
<u>Cuspidaria sp.</u>										0.1	0.32	
<u>Ennucula tenuis</u>	0.1	0.32			0.2	0.63	0.3	0.67			0.2	0.42
<u>Hiatella arctica</u>	0.2	0.42	0.1	0.32								
<u>Macoma calcarea</u>	0.4	0.97							0.1	0.32		

Table 15. cont.

	1972				1973				1974				
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.		
	\bar{x}	s											
<u>Musculus niger</u>	0.4	0.70	0.4	0.70	0.2	0.42					0.2	0.42	
<u>Mya arenaria</u>										0.2	0.42	0.6	0.70
<u>Mya sp.</u>	0.5	0.97	0.9	0.99									
<u>Mysella bidentata</u>	0.2	0.63			0.3	0.67	0.3	0.67					
<u>Mytilus edulis</u>										0.2	0.63		
<u>Nuculana pernula</u>							0.1	0.32					
<u>Parvicardium minimum</u>					2.1	1.66	0.8	0.79	0.5	0.71	0.2	0.42	
<u>Parvicardium scabrum</u>			0.1	0.32			0.1	0.32					
<u>Phaxas pellucidus</u>													
<u>Pseudamussium septemradiatum</u>										0.1	0.32		
<u>Spisula elliptica</u>										0.1	0.32		
<u>Thracia myopsis</u>					0.4	0.70							
<u>Thracia sp.</u>	0.1	0.32			0.4	0.52	0.2	0.42	0.3	0.67	0.4	0.52	
<u>Thyasira sp.</u>	2.6	3.06	1.0	1.25	14.1	8.09	6.7	7.29	4.6	7.38	3.3	6.29	
<u>Venus casina</u>										0.1	0.32		
<u>Phoronis mulleri</u>			0.2	0.42									
<u>Dentalium entale</u>			0.1	0.32						0.1	0.32	0.1	0.32
<u>Onchnesoma steenstrupi</u>	0.1	0.32					0.1	0.32					
<u>Phascolion strombi</u>	0.2	0.42	0.1	0.32	0.1	0.32	0.1	0.32	0.1	0.32			
<u>Ascidia sp.</u>	0.1	0.32											
<u>Pelonaia corrugata</u>	0.2	0.63											
Diversity:	λ	0.92		0.87		0.86		0.88		0.87		0.93	
	H	3.15		2.77		2.62		2.79		2.91		3.07	
	d	9.73		8.25		7.87		8.40		8.45		7.79	
Sampling efficiency:		0.952 < s < 1		0.933 < s < 1		0.982 < s < 1		0.977 < s < 1		0.948 < s < 1		0.935 < s < 1	

Table 16. Station SD 2, 50 m

	1972		1973		1974							
	JUNE X	S	OCT. X	S	JUNE X	S	OCT. X	S	JUNE X	S	OCT. X	S
<i>Actinaria</i> indet.			0.1	0.32								
<i>Virgularia mirabilis</i>	0.1	0.32			0.2	0.42						
<i>Ctenodiscus crispatus</i>	1.3	0.95	2.4	1.78	1.5	1.78	1.9	1.29	1.7	1.42	1.6	1.26
<i>Chaetoderma nitidulum</i>	0.4	0.52	0.1	0.32	0.2	0.42	0.9	1.29	0.1	0.32	0.2	0.42
Amphipoda indet.	0.1	0.32			0.1	0.32	0.1	0.32			0.1	0.32
Cumacea indet.	0.7	1.25	0.1	0.32	0.3	0.67	0.3	0.67				
<i>Brisaster fragilis</i>	0.1	0.32					0.3	0.48				
<i>Brisopsis lyrifera</i>	0.2	0.63	0.3	0.48	0.1	0.32						
<i>Buccinum</i> sp.			0.1	0.32								
<i>Buccinum undatum</i>	0.1	0.32										
<i>Cylichna cylindracea</i>	0.2	0.63	0.3	0.48								
<i>Lunatia intermedia</i>							0.1	0.32				
<i>Lunatia pallida</i>									0.6	0.70		
<i>Natica</i> sp.					0.1	0.32						
<i>Philine scabra</i>	0.2	0.63	0.2	0.63	0.1	0.32						
Holothuroidea indet.	0.5	1.27			0.5	0.85	0.4	0.70	0.2	0.42	0.5	0.71
Nemertini indet.			1.2	0.92	0.7	0.82	1.4	1.07	0.4	0.52	0.8	0.79
<i>Amphilepis norvegica</i>					0.5	1.27						
<i>Amphiura chiajei</i>	0.3	0.48	0.3	0.67	0.7	0.95	0.7	1.57	1.4	1.78	0.6	0.70
<i>Amphiura filiformis</i>	0.2	0.42	0.1	0.32			0.4	0.70	0.9	0.74	0.4	0.70
<i>Ophiura albida</i>							0.1	0.32				
<i>Ophiura sarsi</i>			0.1	0.32			0.1	0.32	0.1	0.32		
<i>Ophiura</i> sp.									0.1	0.32		
Ampharetidae indet.							0.1	0.32				
<i>Anaitides</i> sp.	0.5	1.27					0.1	0.32				
<i>Anobothrus gracilis</i>			0.1	0.32								
<i>Artacama proboscidea</i>									0.1	0.32		
Capitellidae indet.									0.1	0.32		
<i>Ceratocephale loveni</i>	0.3	0.95							0.2	0.42		
<i>Chaetozone setosa</i>							0.1	0.32				
Cirratulidae indet.							0.1	0.32				
<i>Diplocirrus glaucus</i>	9.0	6.78	6.2	3.05	3.1	2.08	3.3	3.27	1.2	1.23	0.7	1.06
<i>Euchone papillosa</i>							0.1	0.32				
Euclymeninae indet.											0.1	0.32
<i>Glycera alba</i>	0.2	0.63			0.4	0.84	1.6	1.35	1.4	1.43		
<i>Glycera rouxi</i>	2.1	1.52	4.0	1.94	2.3	1.57	1.1	1.20	2.0	1.05	1.9	1.79
<i>Goniada maculata</i>	2.7	2.00	3.5	1.72	4.4	1.58	3.5	2.51	2.2	1.69	0.6	1.35
<i>Heteromastus filiformis</i>	1.7	1.83			0.1	0.32	0.3	0.67	0.7	0.67		
<i>Lanassa venusta</i>							0.1	0.32				
<i>Laonice cirtata</i>	0.1	0.32	0.3	0.48	0.1	0.32	0.5	0.71	0.4	0.52		
<i>Lumbrineris fragilis</i>	0.4	0.70	0.1	0.32	0.2	0.42	0.8	1.03	0.3	0.48	0.2	0.42
<i>Maldane sarsi</i>	15.7	22.12	24.8	25.78	12.1	9.39	25.0	24.67	22.1	15.65	3.7	3.23

Table 16. cont.

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{X}	s	\bar{X}	s	\bar{X}	s	\bar{X}	s	\bar{X}	s	\bar{X}	s
<u>Myriochele</u> sp.	12.2	7.21	14.5	14.33	10.7	6.85	5.4	4.53	2.5	1.90	0.9	1.60
<u>Neoamphitrite affinis</u>						0.5	1.58					
<u>Neoamphitrite grayi</u>						0.5	1.27					
<u>Nephtys ciliata</u>	0.1	0.32			0.1	0.32						
<u>Nephtys</u> sp.		0.1	0.32									
<u>Nereimyra punctata</u>								0.1	0.32			
<u>Ophiodromus flexuosus</u>	0.1	0.32	0.3	0.67	0.1	0.32	0.2	0.42	0.1	0.32	0.2	0.42
<u>Owenia fusiformis</u>	0.5	0.71	0.2	0.63	2.6	2.46	0.4	0.70	0.3	0.48	0.3	0.95
<u>Pectinaria auricoma</u>	0.1	0.32	0.1	0.32	0.4	0.52	0.6	0.70	1.2	0.92	0.3	0.67
<u>Pectinaria koreni</u>	1.0	1.56	0.3	0.48	0.1	0.32	0.3	0.48	0.2	0.63	0.1	0.32
<u>Pholoe minuta</u>	1.4	1.17	1.6	0.97	0.8	0.63	0.8	1.62			0.5	0.71
Phyllodocidae indet.						0.1	0.32	0.1	0.32			
Polynoidae indet.	0.2	0.42	0.2	0.42		0.6	0.84			0.4	0.52	
<u>Polyphsia crassa</u>	0.8	0.63	2.4	0.97	0.7	0.82	0.5	0.53	0.3	0.48	0.1	0.32
<u>Praxillella gracilis</u>	0.1	0.32	0.4	0.52		0.2	0.42	0.6	0.70			
<u>Praxillella praetermissa</u>	7.9	2.51	10.0	3.89	4.7	2.79	3.8	2.10	4.3	2.41	2.3	1.95
<u>Rhodine gracilior</u>	0.1	0.32				0.1	0.32					
<u>Scalibregma inflatum</u>									0.1	0.32		
<u>Scoloplos armiger</u>								0.1	0.32			
Spionidae indet.						0.6	1.35	0.4	0.84	0.1	0.32	
<u>Spiophanes krøyeri</u>	0.1	0.32	0.2	0.42								
<u>Streblosoma bairdi</u>								0.1	0.32			
Terebellidae indet.		0.1	0.32	0.1	0.32							
<u>Terebellides stroemi</u>						0.1	0.32					
<u>Trichobranchus roseus</u>	0.1	0.32						0.1	0.32			
<u>Abra alba</u>	5.5	3.75	7.5	3.34	50.6	24.18	72.5	33.75	57.7	18.35	27.8	8.51
<u>Abra nitida</u>	16.0	6.00	5.2	4.57	8.0	3.43	8.8	5.22	10.7	6.48	4.6	5.87
<u>Arctica islandica</u>					0.3	0.67	0.2	0.63				
<u>Corbula gibba</u>	0.5	0.53			0.2	0.42	0.6	0.84	0.3	0.48		
<u>Ennucula tenuis</u>	1.2	1.55			2.4	1.78	1.6	1.35	2.2	1.23	1.3	2.06
<u>Macoma calcarea</u>		0.1	0.32									
<u>Montacuta</u> sp.		0.3	0.67									
<u>Montacuta tenella</u>					0.2	0.63						
<u>Mya arenaria</u>					0.1	0.32	0.2	0.63				
<u>Mysella bidentata</u>	0.1	0.32	0.2	0.63	0.2	0.63	0.1	0.32				
<u>Nuculana pernula</u>	1.0	1.33	0.3	0.48	0.5	0.53	0.4	0.52	0.8	0.63	0.6	0.97
<u>Parvicardium minimum</u>					0.2	0.42	0.1	0.32			0.1	0.32
<u>Parvicardium ovale</u>				0.1	0.32							
<u>Thyasira</u> sp.	6.8	3.94	3.8	2.90	9.9	4.41	8.8	7.35	4.6	4.84	5.2	6.20
<u>Phoronis mülleri</u>	0.4	1.26	0.3	0.67								
<u>Onchnesoma steenstrupi</u>	0.6	0.52	0.1	0.32					0.2	0.42		
<u>Polycladia</u> indet.			0.1	0.32								
Diversity:	λ	0.90		0.87		0.79		0.73		0.74		0.73
	H	2.69		2.50		2.23		2.09		2.02		2.08
	d	6.57		6.15		5.64		6.97		5.62		4.73
Sampling efficiency:	0.978 < s < 1	0.975 < s < 1	0.985 < s < 1	0.985 < s < 1	0.984 < s < 1	0.975 < s < 1						

Table 18. Station AF 2, 20 m

	1974			
	JUNE		OCT.	
	\bar{X}	s	\bar{X}	s
<u>Ctenodiscus crispatus</u>			0.2	0.42
<u>Chaetoderma nitidulum</u>	0.4	0.70		
Amphipoda indet.			0.2	0.42
<u>Macropipus depurator</u>			0.2	0.42
<u>Lunatia intermedia</u>			0.1	0.32
<u>Cyllichna cylindracea</u>			0.2	0.42
Nemertini indet.	0.3	0.67	0.1	0.32
<u>Amphiura chiajei</u>	0.4	0.97		
<u>Amphiura filiformis</u>	0.3	0.95	0.8	1.14
<u>Amphiura</u> sp.			0.4	0.70
<u>Ophiura affinis</u>			0.1	0.32
<u>Ophiura albida</u>			0.3	0.67
<u>Ophiura</u> sp.			0.3	0.67
<u>Ophiura texturata</u>	0.1	0.32	0.2	0.42
<u>Anaitides</u> sp.			0.1	0.32
<u>Aphrodisita aculeata</u>	0.1	0.32		
<u>Chaetozone setosa</u>			0.1	0.32
<u>Diplocirrus glaucus</u>	0.2	0.42		
<u>Euchone papillosa</u>			0.5	1.08
<u>Glycera alba</u>	0.1	0.32	0.4	0.70
<u>Glycera capitata</u>	0.1	0.32		
<u>Glycera rouxi</u>	0.2	0.42		
<u>Heteromastus filiformis</u>	0.2	0.63	0.3	0.95
<u>Lanassa venusta</u>			0.1	0.32
<u>Maldane sarsi</u>	11.9	21.96	25.6	57.10
<u>Myriochela</u> sp.	2.9	2.60	0.9	1.29
<u>Nephtys ciliata</u>			0.1	0.32
<u>Ophelina acuminata</u>			0.1	0.32
<u>Ophiodromus flexuosus</u>			0.2	0.42
<u>Owenia fusiformis</u>	0.4	0.97		
<u>Paramphipnoma jeffreysi</u>			0.3	0.95
<u>Pectinaria auricoma</u>	0.3	0.48		
<u>Pectinaria koreni</u>	0.2	0.42	0.2	0.42
<u>Pholoe minuta</u>	0.2	0.63	0.2	0.42
Polynoidae indet.			0.1	0.32
<u>Scalibregma inflatum</u>			0.2	0.63
<u>Streblosoma bairdi</u>	0.2	0.42		
Syllidae indet.	0.3	0.48		
<u>Terebellides stroemi</u>			0.1	0.32
<u>Trichobranchus roseus</u>	0.1	0.32		
<u>Abra nitida</u>	0.2	0.42	0.3	0.67
<u>Arctica islandica</u>			0.1	0.32
<u>Astarte elliptica</u>	0.1	0.32	0.1	0.32
<u>Astarte montagui</u>	2.3	1.89	1.0	1.41
<u>Corbula gibba</u>	1.2	1.14	2.4	1.71
<u>Ennucula tenuis</u>	0.1	0.32	0.1	0.32
<u>Mya arenaria</u>			0.2	0.63
<u>Nuculana pernula</u>	0.4	0.84		
<u>Thyasira</u> sp.	1.5	2.17	3.2	4.80
<u>Phoronis mülleri</u>	0.1	0.32		
<u>Phascolion strombi</u>	0.1	0.32		
Sipunculida indet.	0.2	0.42		
Diversity:	λ	0.75	0.57	
	H	2.12	1.67	
	d	5.25	5.68	
Sampling efficiency:	0.936 < S < 1		0.952 < S < 1	

Table 19. Station AF 3, 50 m

	1974			
	JUNE		OCT.	
	\bar{x}	s	\bar{x}	s
<i>Chaetoderma nitidulum</i>	0.2	0.42		
<i>Cumacea indet.</i>	0.6	0.70	0.2	0.42
<i>Nemertini indet.</i>	0.2	0.42		
<i>Ophiura affinis</i>			0.2	0.63
<i>Ophiura sarsi</i>			0.1	0.32
<i>Ophiura texturata</i>	0.2	0.63		
<i>Ampharetidae indet.</i>			0.1	0.32
<i>Anaitides sp.</i>	0.6	0.52		
<i>Artacama proboscidea</i>			0.2	0.63
<i>Chaetozone setosa</i>	1.6	1.71	3.7	3.43
<i>Diplocirrus glaucus</i>	1.1	1.29	0.6	0.70
<i>Euchone papillosa</i>	3.7	2.50	1.9	3.11
<i>Glycera alba</i>	1.2	1.23		
<i>Glycera rouxi</i>	0.2	0.42	0.1	0.32
<i>Goniada maculata</i>	0.5	0.97	0.3	0.48
<i>Heteromastus filiformis</i>	9.2	8.78	8.2	8.46
<i>Lanassa venusta</i>	0.1	0.32		
<i>Laonice cirtata</i>	0.2	0.42		
<i>Lumbrineris fragilis</i>	0.1	0.32		
<i>Maldane sarsi</i>	0.1	0.32	0.8	1.32
<i>Melinna cristata</i>	0.1	0.32		
<i>Myriochele sp.</i>	2.0	1.56	0.9	1.29
<i>Neoamphitrite grayi</i>			0.1	0.32
<i>Neoamphitrite sp.</i>	0.1	0.32		
<i>Nephysa ciliata</i>			0.3	0.67
<i>Nephysa hombergi</i>	0.2	0.42		
<i>Nereimyra punctata</i>	0.1	0.32	0.1	0.32
<i>Nereis sp.</i>			0.1	0.32
<i>Ophelina acuminata</i>			0.4	1.26
<i>Ophelina norvegica</i>	0.2	0.42		
<i>Ophiodromus flexuosus</i>	0.3	0.67	0.7	0.82
<i>Paramphinome jeffreysi</i>	0.8	1.03	1.0	1.49
<i>Pectinaria Koreni</i>	0.3	0.48		
<i>Pholoe minuta</i>	0.3	0.48	0.4	0.52
<i>Phyllodocidae indet.</i>	0.1	0.32		
<i>Polyphysia crassa</i>	0.1	0.32	0.3	0.67
<i>Sabellides octocirrata</i>	0.4	0.70		
<i>Scalibregma inflatum</i>	0.6	1.35	7.1	10.31
<i>Scoploplos armiger</i>	0.4	0.70	0.5	0.97
<i>Spionidae indet.</i>	0.1	0.32	0.1	0.32
<i>Syllidae indet.</i>	1.1	1.52	0.5	0.85
<i>Terebellidae indet.</i>	0.1	0.32		
<i>Terebellides stroemi</i>	0.4	0.70	0.5	0.97
<i>Trichobranchus roseus</i>	0.1	0.32		
<i>Abra alba</i>	0.8	1.23		
<i>Abra nitida</i>	1.8	1.23	1.2	0.92
<i>Astarte montagui</i>	0.2	0.42	0.6	0.70
<i>Corbula gibba</i>	1.5	1.90	2.5	2.27
<i>Lucinoma borealis</i>	0.1	0.32		
<i>Mya arenaria</i>	0.1	0.32		
<i>Palliolium furtivum</i>			0.1	0.32
<i>Palliolium vitreum</i>	0.2	0.63		
<i>Thyasira sp.</i>	10.4	9.12	6.0	6.13
	λ	0.88		0.89
Diversity:	H	2.72		2.59
	d	7.10		5.18
Sampling efficiency:		0.952 < S < 1		0.962 < S < 1

Table 20. Station PT 1, 20 m

	1972				1973				1974				
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.		
	\bar{x}	s											
<u>Virgularia mirabilis</u>	0.6	0.70	0.1	0.32	0.1	0.32	0.2	0.42	0.3	0.48			
<u>Astropecten irregularis</u>			0.1	0.32					0.1	0.32	0.1	0.32	
<u>Ctenodiscus crispatus</u>	0.2	0.42			0.1	0.32					0.3	0.48	
<u>Chaetoderma nitidulum</u>	0.2	0.42	0.3	0.48	1.0	1.15	0.6	1.07	0.9	1.10	1.7	1.49	
<u>Amphipoda</u> inde.	1.2	1.75	0.6	0.84	0.4	0.97	0.3	0.48			1.1	2.18	
<u>Balanus balanus</u>					0.1	0.32							
<u>Calocharis macandreae</u>											0.1	0.32	
<u>Cumacea</u> indet.				0.1	0.32	0.4	0.52	0.1	0.32	0.1	0.32	0.1	0.32
<u>Brisaster fragilis</u>	0.1	0.32			0.1	0.32	0.1	0.32					
<u>Echinocardium cordatum</u>					0.2	0.63	0.3	0.48	0.1	0.32	0.4	0.70	
<u>Aporrhais pespelicanii</u>	0.1	0.32	0.2	0.42	0.1	0.32							
<u>Buccinum</u> sp.				0.1	0.32								
<u>Okenia pulchella</u>				0.1	0.32								
<u>Philine quadrata</u>				0.1	0.32								
<u>Philine scabra</u>						0.9	0.88						
<u>Philine</u> sp.	0.1	0.32											
<u>Turitella communis</u>	0.8	1.03	0.5	1.58	0.3	0.48	0.2	0.42	0.5	0.71			
<u>Cucumaria elongata</u>				0.9	0.74				0.9	1.29			
<u>Holothuroidea</u> indet.	0.4	0.70	0.2	0.63	3.1	2.02	3.3	1.77	4.1	4.20	4.5	3.03	
<u>Nemertini</u> indet.				0.4	0.52	1.4	0.84	1.5	0.85	1.4	1.17	2.2	1.32
<u>Amphiura chiajei</u>	3.1	2.69	2.4	1.90	3.3	1.89	0.2	0.63	1.2	0.92	3.3	2.36	
<u>Amphiura filiformis</u>	2.0	1.94			2.8	1.75	0.5	1.27	1.4	1.07	0.5	1.27	
<u>Amphiura</u> sp.			0.1	0.32									
<u>Ophiura affinis</u>						0.2	0.42				0.3	0.67	
<u>Ophiura albida</u>	3.0	2.49	0.3	0.48	0.9	0.88	0.3	0.67	0.5	0.85	0.7	0.95	
<u>Ophiura sarsi</u>					0.2	0.42	0.3	0.48			0.1	0.32	
<u>Ophiura</u> sp.				0.4	0.70	0.5	0.53	0.7	0.67	0.1	0.32	0.5	0.71
<u>Ophiura texturata</u>				0.9	1.20	0.2	0.63	0.5	0.71	0.7	1.06	0.9	0.74
<u>Ampharetidae</u> indet.									0.1	0.32			
<u>Anaitides</u> sp.	0.3	0.67	2.9	1.45	1.6	1.07	1.9	1.37	0.9	0.99	1.8	1.40	
<u>Asychnis biceps</u>	4.1	2.08	2.6	3.31	3.8	2.10	4.1	4.20	3.1	1.60	3.3	1.34	
<u>Ceratocephale loveni</u>	0.1	0.32				0.1	0.32						
<u>Chaetozone setosa</u>										0.2	0.42		
<u>Cirratulus cirratus</u>				0.2	0.63		0.3	0.67			0.2	0.63	
<u>Diplocirrus glaucus</u>	3.6	1.71	0.2	0.63	1.9	3.38	2.8	1.81	7.9	3.96	4.0	2.11	
<u>Eteoninae</u> indet.						0.1	0.32						
<u>Eteone</u> sp.	0.1	0.32											
<u>Euchone papillosa</u>						0.2	0.42	0.2	0.42	0.2	0.42		
<u>Euchone rubrocincta</u>				0.1	0.32								
<u>Euclymeninae</u> indet.										0.1	0.32		
<u>Eumida</u> sp.	0.1	0.32											
<u>Glycera alba</u>	1.2	0.42	1.0	1.05	0.8	1.03	0.1	0.32	0.5	0.53	0.6	0.97	
<u>Glycera rouxi</u>					0.1	0.32	0.1	0.32			0.2	0.42	
<u>Goniada maculata</u>	0.7	0.95	1.1	1.20	0.6	1.35	1.3	1.49	0.6	0.84	1.9	1.45	

Table 20. cont.

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s
<u>Goniada norvegica</u>					0.1	0.32						
<u>Heteromastus filiformis</u>	0.3	0.48	0.3	0.67	0.8	1.23	2.7	2.16	0.9	1.20	0.5	0.71
<u>Lanassa nordenskiøldi</u>	0.1	0.32	0.1	0.32								
<u>Lanassa venusta</u>					0.1	0.32			0.4	0.70	0.2	0.63
<u>Laonice cinctata</u>	0.5	0.53	1.0	1.25	1.0	1.25	0.8	0.63	0.9	0.99	1.1	0.88
<u>Lumbrineris fragilis</u>	1.4	1.26	2.2	1.75	0.6	0.52	0.6	0.70	0.2	0.42	0.8	0.79
<u>Maldanidae</u> indet.	0.5	0.71										
<u>Maldane sarsi</u>	1.4	3.10	0.2	0.42	3.1	4.65	8.6	4.43	3.1	5.04	2.2	3.65
<u>Melinna cristata</u>			0.1	0.32					0.1	0.32		
<u>Myriochele</u> sp.			0.3	0.67	1.4	1.84	2.4	2.41	1.8	1.48	1.9	1.79
<u>Neoamphitrite affinis</u>					0.1	0.32						
<u>Neoamphitrite grayi</u>			0.1	0.32	0.1	0.32						
<u>Nephtys caeca</u>							0.2	0.42				
<u>Nephtys ciliata</u>	1.3	1.06	1.7	1.70	1.6	1.07	0.9	0.99	1.3	1.16	0.9	0.88
<u>Nephtys hombergi</u>							0.1	0.32	0.1	0.32		
<u>Nephtys paradoxa</u>			0.3	0.48	0.1	0.32				0.2	0.42	
<u>Nephtys</u> sp.					0.2	0.42						
<u>Nicomache lumbicalis</u>	1.6	2.22	2.3	2.36	0.4	0.70	0.4	0.70	0.9	1.20	1.5	1.58
<u>Ophelina acuminata</u>							0.1	0.32			0.4	0.70
<u>Ophiodromus flexuosus</u>	0.4	0.84	0.6	0.52	0.5	0.71	0.9	0.88	1.2	1.03	0.7	0.57
<u>Owenia fusiformis</u>			0.1	0.32								
<u>Paraonis gracilis</u>	0.1	0.32			0.2	0.42	0.2	0.42	0.1	0.32	0.1	0.32
<u>Pectinaria auricoma</u>					0.8	1.32	1.0	1.25			0.1	0.32
<u>Pectinaria koreni</u>	7.1	2.42	10.4	8.15	5.1	2.02	4.0	3.20	5.3	2.50	6.5	4.88
<u>Pholoe minuta</u>	2.0	1.76	4.2	2.30	6.0	4.45	3.8	2.25	3.1	1.29	8.8	3.16
<u>Phyllodocidae</u> indet.									0.1	0.32	0.2	0.42
<u>Polydora</u> sp.			0.1	0.32			0.1	0.32				
<u>Polynoidae</u> indet.	0.2	0.42			0.2	0.42	0.2	0.42	0.2	0.42	0.2	0.42
<u>Polypysia crassa</u>									0.1	0.32	0.1	0.32
<u>Praxillella gracilis</u>	0.1	0.32									0.1	0.32
<u>Rhodine gracilior</u>	1.0	0.67	0.8	0.63			0.1	0.32	0.1	0.32	0.4	0.70
<u>Scalibregma inflatum</u>									0.1	0.32	0.3	0.48
<u>Scoloplos armiger</u>	0.8	1.14	5.8	2.44	5.9	4.28	1.1	1.29	2.4	1.07	2.2	1.62
<u>Spio filiformis</u>											0.1	0.32
<u>Spionidae</u> indet.	2.6	3.06							0.1	0.32		
<u>Streblosoma bairdi</u>	0.2	0.42			0.2	0.42	0.2	0.42	0.1	0.32	0.1	0.32
<u>Syllidae</u> indet.	0.4	0.70	0.3	0.67	0.4	0.52	0.5	0.71	0.2	0.42	0.1	0.32
<u>Terebellidae</u> indet.					0.1	0.32						
<u>Terebellides stroemi</u>	0.1	0.32	0.2	0.42	0.2	0.42	0.2	0.42	2.6	1.65	5.7	2.11
<u>Tharyx marioni</u>			0.1	0.32								
<u>Thelepus cincinnatus</u>					0.1	0.32	0.1	0.32				
<u>Trichobranchus roseus</u>	0.5	0.85	0.6	0.84	0.6	0.70	0.5	0.85	0.7	0.95	1.0	1.05
<u>Trochochaeta multisetosa</u>							0.1	0.32				
<u>Abra alba</u>	1.0	0.94	0.2	0.42	92.8	23.92	39.5	17.74	7.1	3.96	4.4	2.76
<u>Abra nitida</u>	1.3	1.16	1.1	1.10	141.6	62.64	78.8	29.34	9.4	5.76	21.5	11.58
<u>Acanthocardia echinata</u>				0.1	0.32	0.3	0.48		0.1	0.32	0.2	0.42
<u>Arctica islandica</u>	0.6	0.84			0.4	0.70	0.1	0.32	0.2	0.42	0.3	0.48
<u>Corbula gibba</u>	1.2	1.32	0.2	0.42	9.4	6.43	3.0	3.02	5.0	3.74	55.0	20.39
<u>Ennucula tenuis</u>					0.1	0.32	0.1	0.32	0.2	0.42		
<u>Gari ferven</u>											0.1	0.32
<u>Hiatella arctica</u>			0.2	0.63	0.2	0.63						0.2
<u>Kellia suborbicularis</u>												0.42
<u>Kelliella miliaris</u>					0.1	0.32						
<u>Lucinoma borealis</u>							0.1	0.32				
<u>Macoma calcarea</u>	0.2	0.42			0.2	0.42	0.2	0.42	0.8	1.03	0.8	0.79

Table 20. cont.

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{x}	s										
<u>Montacuta ferruginosa</u>					0.1		0.32		0.2		0.42	
<u>Montacuta</u> sp.											0.4	0.52
<u>Musculus niger</u>	0.4		0.52									
<u>Mya arenaria</u>					0.2		0.42		0.1		0.32	
<u>Mya</u> sp.					0.4		0.70		0.1		0.32	
<u>Mysella bidentata</u>	0.2		0.42		1.7		1.77		0.2		0.42	
<u>Mysia undata</u>									0.1		0.32	
<u>Nucula</u> sp.	0.1		0.32								0.1	
<u>Thracia</u> sp.									0.3		0.48	1.4
<u>Thyasira</u> sp.	1.8		1.14		2.1		3.75		6.7		5.68	42.2
<u>Venus casina</u>									21.05		23.1	
											4.91	41.6
											0.5	8.98
												0.71
<u>Phoronis müllerii</u>	0.3		0.48		0.3		0.48		0.2		0.42	
<u>Dentalium entale</u>					0.5		0.71				0.4	
<u>Onchnesoma steenstrupi</u>	0.1		0.32						0.1		0.32	
<u>Phascolion strombi</u>	0.9		1.37		0.2		0.42		0.2		0.42	
									0.2		0.4	
											0.52	
	λ		0.95		0.93		0.70		0.79		0.91	
Diversity:	H		3.38		3.15		1.87		2.19		3.06	
	d		8.30		8.78		8.21		8.47		8.55	
Sampling efficiency:		0.964 < s < 1		0.957 < s < 1		0.996 < s < 1		0.987 < s < 1		0.976 < s < 1		0.989 < s < 1

Table 21. Station FT 2, 50 m

	1972				1973				1973			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{x}	s										
<u>Stylatula elegans</u>									0.1	0.32	0.1	0.32
<u>Virgularia mirabilis</u>	1.0	1.25	0.9	0.57	0.8	1.23	0.2	0.42	1.6	1.35	0.3	0.67
<u>Ctenodiscus crispatus</u>	0.7	0.67	1.2	0.92	0.3	0.48	0.2	0.42	0.4	0.52		
<u>Chaetoderma nitidulum</u>			0.4	1.26	0.3	0.48	0.6	0.70	0.3	0.95	0.2	0.42
<u>Lepidopleurus alveolus</u>							0.6	1.35				
Amphipoda indet.	1.1	1.45	1.7	2.00	1.3	1.95	0.8	1.03	0.2	0.42	0.3	0.48
<u>Calocharis macandreae</u>									0.1	0.32		
Cumacea indet.	1.8	1.75	1.9	2.13	2.0	1.76	1.0	1.33	0.2	0.42	0.7	0.67
<u>Galathea</u> sp.					0.1	0.32						
<u>Gnathia oxyraea</u>			0.1	0.32					0.1	0.32		
<u>Brisaster fragilis</u>							0.1	0.32				
<u>Brisopsis lyrifera</u>									0.1	0.32		
<u>Echinocardium cordatum</u>							0.1	0.32				
<u>Echinocyanus pusillus</u>						0.1	0.32					
<u>Buccinum undatum</u>	0.1	0.32										
<u>Eulima stenostoma</u>					0.3	0.67						
<u>Iothia fulva</u>							0.1	0.32				
<u>Lepeta caeca</u>							0.1	0.32				
<u>Lunatia pallida</u>	0.2	0.42	0.3	0.48			0.2	0.42			0.3	0.48
<u>Neptunea despecta</u>			0.1	0.32								
<u>Philine scabra</u>					0.1	0.32						
<u>Turitella communis</u>			0.2	0.63								
Holothuridea indet.	0.2	0.63	0.3	0.48	0.3	0.67	1.1	1.45	0.8	0.92	0.9	1.60
Nemertini indet.				1.1	0.88	1.1	0.74	1.7	1.49	1.3	1.34	1.4
<u>Amphilepis norvegica</u>									0.2	0.42		
<u>Amphiura chiajei</u>	0.5	0.71	0.6	0.97	0.4	0.97	0.8	1.03	0.4	0.52		
<u>Amphiura filiformis</u>	0.1	0.32			0.3	0.48	0.5	0.85	0.3	0.48		
<u>Ophiopholis aculeata</u>						0.4	1.26					
<u>Ophiura affinis</u>							0.1	0.32				
<u>Ophiura albida</u>	0.3	0.67					0.6	0.70	0.7	0.95		
<u>Ophiura carnea</u>							0.8	1.48				
Ophiuroidea indet.	0.1	0.32										
<u>Ophiura robusta</u>						0.3	0.95					
<u>Ophiura sarsi</u>			0.3	0.48	0.1	0.32	0.6	0.52	0.2	0.42		
<u>Ophiura</u> sp.					1.3	1.34	0.5	0.53	0.1	0.32		
<u>Ophiura texturata</u>								0.2	0.63			
Ampharetidae indet.							0.1	0.32				
<u>Anaitides</u> sp.	0.6	0.70	0.6	1.07	0.1	0.32	0.2	0.42	0.3	0.67	1.9	1.66
<u>Asy whole biceps</u>	0.1	0.32	2.0	2.16	0.6	0.70	0.8	1.23	0.4	0.70	0.1	0.32
<u>Chaetozone setosa</u>	0.1	0.32	0.4	0.52			0.1	0.32				
<u>Cirratulus cirratus</u>	1.4	1.90					0.1	0.32			0.2	0.63
Cirratulidae indet.					0.1	0.32						
<u>Dasybranchus caducus</u>	0.4	0.52	0.4	0.52					0.4	0.70		

Table 21. cont.

	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{X}	s										
<u>Diplocirrus glaucus</u>			4.6	2.46	1.1	0.99	6.2	4.80	6.1	6.77	7.6	2.99
<u>Driloneris filum</u>			0.9	0.74			0.2	0.42	0.2	0.42	0.3	0.67
<u>Eteone sp.</u>			0.1	0.32								
<u>Euclymeninae</u> indet.					0.3	0.48					0.5	0.97
<u>Eunice pennata</u>							1.8	4.05				
<u>Glycera alba</u>	0.2	0.42	1.0	1.15	0.3	0.48	0.4	0.70	0.7	1.06	0.4	0.84
<u>Glycera rouxii</u>	0.1	0.32	0.2	0.42	0.4	0.70	0.2	0.42			0.6	0.52
<u>Goniada maculata</u>	0.7	0.67	0.6	0.52	0.1	0.32	0.4	0.52	0.4	0.52	1.4	1.43
<u>Hesionidae</u> indet.											0.1	0.32
<u>Heteromastus filiformis</u>	67.1	29.32	23.8	25.42	7.1	9.79	3.7	4.64	6.7	5.12	17.8	10.12
<u>Laonice cirrata</u>	0.2	0.42	0.2	0.42	0.1	0.32	0.4	0.52	0.1	0.32	0.1	0.32
<u>Lumbrineris fragilis</u>	0.6	0.84	0.1	0.32	0.3	0.48	0.8	1.14	0.7	0.48	0.7	0.95
<u>Maldanidae</u> indet.							0.2	0.63				
<u>Maldane sarsi</u>	4.6	3.20	18.1	11.47	1.7	1.34	8.0	5.62	5.2	6.25	3.2	3.99
<u>Melinna cristata</u>	0.2	0.63			0.1	0.32	1.1	1.60				
<u>Microclymene tricirrata</u>			0.1	0.32								
<u>Myriochele</u> sp.	1.1	0.88	3.6	2.32	0.3	0.48	1.2	1.32	0.4	0.84	0.5	0.85
<u>Nephtys ciliata</u>	0.9	0.99	0.5	0.53	0.3	0.48	0.1	0.32	0.7	0.82	0.3	0.48
<u>Nephtys paradox</u>	0.1	0.32			0.1	0.32						
<u>Nereis virens</u>			0.1	0.32								
<u>Notomastus latericeus</u>									0.3	0.67		
<u>Ophelina acuminata</u>	0.2	0.63	0.1	0.32					0.1	0.32		
<u>Ophiodromus flexuosus</u>	0.5	0.71	0.2	0.42	0.5	0.97	0.2	0.42	0.2	0.42	0.1	0.32
<u>Owenia fusiformis</u>	1.9	1.66	9.4	9.44	0.8	1.32	0.6	1.07	0.4	0.97	1.6	1.43
<u>Paraphinome jeffreysi</u>	0.1	0.32					0.4	0.70				
<u>Paraonis gracilis</u>							0.1	0.32				
<u>Pectinaria auricoma</u>	0.4	0.70			1.1	0.99	1.1	1.10	0.8	0.79	1.1	1.29
<u>Pectinaria belgica</u>											0.1	0.32
<u>Pectinaria koreni</u>	2.9	1.52	6.0	3.80	4.4	2.37	2.2	2.66	2.3	1.89	0.4	0.52
<u>Pholoe minuta</u>	3.0	1.89	4.7	2.95	0.8	0.79	2.2	2.44	1.6	1.26	3.7	2.31
<u>Phyllodocidae</u> indet.							0.2	0.42	0.1	0.32		
<u>Polynoidae</u> indet.							0.4	0.52	0.2	0.42	0.5	0.71
<u>Polyneoe kinbergi</u>	0.1	0.32										
<u>Polyphysia crassa</u>			0.1	0.32	0.1	0.32	0.6	1.07	0.6	0.70	0.3	0.67
<u>Praxillella gracilis</u>					0.3	0.67	0.1	0.32				
<u>Praxillella praetermissa</u>							0.4	0.70	0.5	0.71		
<u>Prionospio cirrifera</u>			1.0	1.33								
<u>Rhodine loveni</u>			0.1	0.32								
<u>Sabellidae</u> indet.											0.1	0.32
<u>Scalibregma inflatum</u>							0.2	0.42	0.7	0.82	0.6	0.84
<u>Scoloplos armiger</u>			0.3	0.48	0.1	0.32			0.1	0.32		
<u>Spionidae</u> indet.							0.1	0.32				
<u>Streblosoma intestinalis</u>												
<u>Syllidae</u> indet.	0.2	0.42	0.4	0.52	0.1	0.32	1.3	1.16	0.4	0.52	0.5	0.71
<u>Terebellides stroemii</u>	0.1	0.32					0.4	0.70	0.3	0.67	0.3	0.67
<u>Trichobranchus roseus</u>	0.1	0.32			0.1	0.32	0.2	0.42	0.1	0.32		
<u>Trochochaeta multisetosa</u>					0.1	0.32						
<u>Abra alba</u>	36.9	8.56	21.1	9.47	404.2	80.21	201.6	94.87	131.9	20.11	66.5	13.83
<u>Abra nitida</u>	63.5	12.87	33.0	13.57	64.9	36.87	19.4	13.47	20.9	9.95	13.9	4.63
<u>Arctica islandica</u>	0.5	0.71			0.2	0.42					0.2	0.42
<u>Astarte sulcata</u>							0.2	0.63				
<u>Bathyarca pectunculoides</u>					0.1	0.32						
<u>Cardiidae</u> indet.	0.2	0.42										
<u>Corbula gibba</u>	0.3	0.67			0.4	0.70			0.1	0.32	0.5	0.71
<u>Ennucula tenuis</u>	0.4	0.70	2.1	1.60	1.6	1.07	0.4	0.70	0.5	0.71	0.7	1.25
<u>Kelliella miliaris</u>					0.7	1.06						
<u>Macoma calcarea</u>	0.1	0.32	0.2	0.63			0.1	0.32	0.1	0.32		
<u>Modiolus modiolus</u>					0.1	0.32						
<u>Montacuta ferruginosa</u>									0.1	0.32		

Table 21. cont.

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s
<u>Montacuta</u> sp.											0.2	0.42
<u>Musculus</u> niger	0.1	0.32			0.1	0.32						
<u>Mya</u> arenaria					0.1	0.32			0.1	0.32	0.1	0.32
<u>Mya</u> sp.	0.2	0.63	0.1	0.32								
<u>Mysella</u> bidentata	0.5	0.71	0.1	0.32	0.2	0.42	0.1	0.32				
<u>Nuculana</u> minuta							0.3	0.95			0.1	0.32
<u>Nuculana</u> pernula					0.1	0.32						
<u>Parvicardium</u> minimum					1.5	2.80	1.7	1.16	0.1	0.32	0.2	0.42
<u>Thracia</u> sp.					0.1	0.32						
<u>Thyasira</u> sp.	113.8	24.20	194.5	49.23	230.8	55.75	46.0	29.84	61.9	30.06	107.5	32.97
<u>Yoldiella</u> sp.					0.1	0.32	0.1	0.32			0.2	0.42
<u>Phoronis</u> mülleri	0.3	0.67										
<u>Dentalium</u> entale							0.3	0.67	0.1	0.32		
<u>Onchnesoma</u> steenstrupi	0.4	0.70					0.3	0.48				
<u>Phascolion</u> strombi			0.1	0.32	0.1	0.32	0.5	0.85				
Sipunculida indet.	0.1	0.32										
Diversity:	λ	0.76	0.65		0.59	0.58	0.66	0.71				
	H	1.81	1.77		1.21	1.61	1.65	1.78				
	d	6.34	6.03		6.40	8.92	6.76	6.04				
Sampling efficiency:	0.993 < s < 1	0.994 < s < 1	0.996 < s < 1	0.992 < s < 1	0.991 < s < 1	0.993 < s < 1						

Table 22. Station FT 3, 100 m

	1972		1973		1974	
	JUNE X	S	JUNE X	S	JUNE X	S
<u>Paraedwardsia arenaria</u>		0.5	0.71	0.4	0.70	
<u>Pennatula phosphorea</u>					0.1	0.32
<u>Virgularia mirabilis</u>				0.1	0.32	
0.3	0.67	0.1	0.32			
<u>Ctenodiscus crispatus</u>					0.1	0.32
<u>Psilaster andromeda</u>		0.1	0.32			
<u>Chaetoderma nitidulum</u>	1.2	1.40	1.0	1.05	0.7	0.82
0.8	1.05	0.7	0.82	0.8	0.79	2.2
Amphipoda indet.	1.4	1.26	1.8	1.55	0.4	0.70
<u>Calocaris macandreae</u>				0.1	0.32	0.8
Cumacea indet.	0.2	0.42	0.9	1.52	0.5	0.97
0.6	1.0	0.6	0.97	0.6	0.97	1.0
<u>Brisaster fragilis</u>	0.4	0.70			0.1	0.32
<u>Brissopsis lyrifera</u>			0.1	0.32		
<u>Balcis</u> sp.					0.1	0.32
<u>Cylichna cylindracea</u>			0.1	0.32	0.1	0.32
<u>Eulimella scillae</u>	0.1	0.32				
<u>Natica clausa</u>			0.1	0.32	0.1	0.32
<u>Odostomia unidentata</u>					0.1	0.32
<u>Philine quadrata</u>			0.1	0.32		
<u>Philine scabra</u>			0.2	0.42	0.1	0.32
<u>Trichotropis borealis</u>	0.1	0.32				
<u>Trophonopsis barvicensis</u>			0.1	0.32		
Holothuroidea indet.	0.3	0.67	0.8	1.03	0.6	0.84
0.2	0.42	0.2	0.42	0.2	0.42	0.2
Nemertini indet.			1.1	1.29	0.8	0.79
0.6	1.1	0.8	1.29	1.1	0.79	1.6
<u>Amphilepis norvegica</u>	6.2	3.26	5.5	2.37	2.3	2.36
<u>Amphiura chiajei</u>	0.8	1.14	0.6	0.84	0.4	0.52
<u>Amphiura filiformis</u>					2.4	1.71
<u>Amphiura</u> sp.	0.5	1.58			0.1	0.32
<u>Ophiura albida</u>			0.3	0.67	0.3	0.48
<u>Ophiura carnea</u>					0.3	0.48
<u>Ophiura sarsi</u>	0.7	0.82	0.1	0.32		
<u>Ophiura</u> sp.	0.2	0.63	1.5	1.43		
Ampharetidae indet.	0.9	0.99			0.2	0.63
<u>Amphicteis gunneri</u>			0.2	0.63		
<u>Anaitides</u> sp.	0.4	0.52			0.1	0.32
<u>Aphrodita aculeata</u>					0.1	0.32
Aphroditidae indet.	0.1	0.32				
<u>Asychnis biceps</u>	1.1	1.52	3.6	3.47	4.3	2.95
<u>Brada villosa</u>	0.1	0.32	0.3	0.95	0.1	0.32
Capitellidae indet.						0.1
<u>Ceratocephale loveni</u>	0.1	0.32	0.5	1.27	0.1	0.32
<u>Chaetozone setosa</u>	2.0	3.06	4.3	5.08	0.4	0.70
<u>Dasybranchus caducus</u>	0.1	0.32			5.9	5.09
<u>Diplocirrus glaucus</u>	6.1	5.32	5.7	4.97	0.8	1.03
<u>Driloneris filum</u>					6.2	2.94
<u>Euchone papillosa</u>			0.1	0.32	55.7	23.97
0.1	0.32				1.1	37.3
1.60	1.2				1.2	25.53

Table 22. cont.

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{X}	s	\bar{X}	s	\bar{X}	s	\bar{X}	s	\bar{X}	s	\bar{X}	s
<i>Euclymeninae</i> indet.	0.1	0.32	0.4	0.52	0.1	0.32	0.1	0.32			0.1	0.32
<i>Flabelligeridae</i> indet.									0.7	1.06		
<i>Glycera rouxii</i>	0.5	1.08	0.1	0.32	0.2	0.42	0.2	0.42	0.3	0.48	0.2	0.42
<i>Goniada maculata</i>	0.1	0.32	0.2	0.42							0.2	0.42
<i>Harmothoe</i> sp.			0.2	0.42								
<i>Heteromastus filiformis</i>	0.1	0.32	0.2	0.42			0.3	0.67	4.0	2.58	0.8	0.92
<i>Heteroclymene robusta</i>									0.1	0.32		
<i>Laetmonice filicornis</i>							0.1	0.32				
<i>Laonice cirrata</i>	0.2	0.42	0.8	1.23							1.0	1.05
<i>Leanira tetragona</i>			0.2	0.42								
<i>Lumbrineris fragilis</i>	1.7	1.25	2.8	1.75	0.3	0.48	2.6	1.90	1.9	1.60	2.2	1.69
<i>Maldanidae</i> indet.	0.2	0.63							0.1	0.32		
<i>Maldane sarsi</i>			0.3	0.67					0.3	0.67	1.1	1.85
<i>Melinna cristata</i>	9.9	10.70	12.8	11.97	7.7	10.71	1.8	1.62	0.5	1.27	2.9	1.29
<i>Myriochele</i> sp.	0.1	0.32	0.1	0.32	0.2	0.42	0.4	0.70	0.2	0.42	0.4	0.52
<i>Nephtys ciliata</i>					0.3	0.48			0.1	0.32		
<i>Nephtys hombergi</i>											0.1	0.32
<i>Nephtys incisa</i>											0.5	0.53
<i>Nephtys paradoxa</i>	0.1	0.32			0.1	0.32	0.2	0.63	0.4	0.52	0.1	0.32
<i>Ophelina acuminata</i>	0.7	1.06										
<i>Ophelina norvegica</i>			0.3	0.48			0.1	0.32	0.1	0.32	0.3	0.67
<i>Ophiodromus flexuosus</i>							0.1	0.32	0.1	0.32	0.2	0.42
<i>Owenia fusiformis</i>					0.2	0.42			0.1	0.32		
<i>Panthalis oerstedi</i>									0.1	0.32	0.3	0.67
<i>Paramphinome jeffreysi</i>	0.7	0.95	0.1	0.32			0.6	1.26	0.9	1.29	1.2	1.87
<i>Pectinaria auricoma</i>					0.1	0.32	0.1	0.32	0.8	0.79	0.3	0.67
<i>Pectinaria belgica</i>							0.1	0.32				
<i>Pectinaria koreni</i>	0.1	0.32			0.2	0.42						
<i>Pholoe minuta</i>	0.5	0.97	0.3	0.48	0.1	0.32	0.1	0.32	1.0	1.25	0.8	1.14
<i>Phylo norvegicus</i>			0.3	0.67	0.1	0.32	0.3	0.67	0.1	0.32	0.5	0.71
<i>Pista cristata</i>											0.1	0.32
<i>Polynoidae</i> indet.							0.1	0.32			0.2	0.42
<i>Polyphysia crassa</i>			0.1	0.32	0.3	0.67	0.3	0.48	0.4	* 0.52	0.1	0.32
<i>Rhodine loveni</i>	0.3	0.67			0.4	0.70	0.2	0.42	0.2	0.42	0.6	0.70
<i>Sabellidae</i> indet.	0.1	0.32										
<i>Sabella penicilllus</i>					0.1	0.32						
<i>Scalibregma inflatum</i>	0.1	0.32							0.1	0.32		
<i>Scoloplos armiger</i>	0.5	0.97										
<i>Sphaerodorum gracilis</i>											0.1	0.32
<i>Spiochaetopterus typicus</i>											0.1	0.32
<i>Spionidae</i> indet.			0.1	0.32			0.2	0.42	0.6	0.84		
<i>Streblosoma intestinalis</i>			0.9	0.99			6.6	3.57	10.0	9.08		
<i>Syllidae</i> indet.	0.1	0.32					0.2	0.42	1.0	1.05	0.4	0.70
<i>Terebellidae</i> indet.	0.1	0.32			0.3	0.67						
<i>Terebellides stroemii</i>	0.3	0.95	0.4	0.84			0.4	0.70	0.8	0.79	0.2	0.42
<i>Tharyx marioni</i>			0.1	0.32								
<i>Thelepiniae</i> indet.			1.0	1.05							2.1	2.02
<i>Trichobranchus roseus</i>	0.1	0.32							0.1	0.32		
<i>Trochochaeta multisetosa</i>											0.1	0.32
<i>Abra alba</i>	1.5	2.51	0.5	0.85	5.1	4.18	2.9	2.81	4.6	5.64	0.7	1.06
<i>Abra nitida</i>	30.1	31.19	57.6	31.82	14.5	7.66	24.4	5.74	56.9	24.52	34.1	19.91
<i>Arctica islandica</i>	0.4	0.97										
<i>Astarte</i> sp.	0.8	1.32			0.1	0.32						
<i>Astarte sulcata</i>			0.8	1.23	0.9	1.37					0.1	0.32
<i>Bathyarca pectunculoides</i>	0.3	0.67	1.1	1.20	0.8	1.62	0.1	0.32				

Table 22. cont.

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s
<u>Cardiidae</u> indet.	0.1	0.32										
<u>Chlamys islandica</u>			0.1	0.32								
<u>Corbula gibba</u>											0.1	0.32
<u>Cuspidaria rostrata</u>			0.1	0.32								
<u>Cuspidaria</u> sp.	0.1	0.32			0.1	0.32					0.1	0.32
<u>Ennucula tenuis</u>			0.2	0.42								
<u>Kelliella miliaris</u>			1.9	2.77	1.3	1.95	0.1	0.32	0.1	0.32	0.9	1.10
<u>Limatula</u> sp.	0.9	1.29										
<u>Limatula sulcata</u>			1.2	1.32	0.9	1.10	0.8	0.79	0.1	0.32	0.3	0.67
<u>Modiolula phaseolinus</u>	0.1	0.32			0.5	0.97						
<u>Montacuta tenella</u>					0.1	0.32						
<u>Mysella bidentata</u>			0.1	0.32								
<u>Nuculana pernula</u>			0.1	0.32								
<u>Nucula</u> sp.	0.1	0.32										
<u>Nucula tumidula</u>			0.2	0.63								
<u>Palliolum vitreum</u>			0.1	0.32								
<u>Parvicardium minimum</u>			3.1	3.00	1.1	1.45	0.7	0.67	0.6	0.84	0.1	0.32
<u>Pseudamussium septemradiatum</u>			0.2	0.42					0.1	0.32		
<u>Thyasira</u> sp.	17.8	14.89	25.5	13.01	20.8	15.56	16.6	5.72	9.6	7.07	22.1	10.33
<u>Tropidomya abbreviata</u>					0.2	0.42			0.1	0.32		
<u>Yoldiella</u> sp.	3.8	4.08	9.2	6.48	8.2	6.84	4.1	2.02	1.5	1.18	0.6	0.70
<u>Phoronis mülleri</u>	0.2	0.42										
<u>Dentalium entale</u>			0.4	0.97			0.4	0.70	0.3	0.67		
<u>Dentalium occidentale</u>					0.2	0.63	0.2	0.63				
<u>Dentalium</u> sp.			0.1	0.32								
<u>Onchnesoma squamatum</u>			0.1	0.32					0.1	0.32		
<u>Onchnesoma steenstrupi</u>	21.1	13.25	14.1	11.80	10.1	16.38	8.2	5.85	10.3	6.38	7.5	9.38
<u>Polycladida</u> indet.	0.1	0.32										
	λ	0.86		0.84		0.88		0.89		0.83		0.84
Diversity:	H	2.54		2.55		2.65		2.73		2.36		2.34
	d	8.07		8.89		7.96		7.54		7.62		6.96
Sampling efficiency:	0.973 < s < 1		0.982 < s < 1		0.969 < s < 1		0.976 < s < 1		0.986 < s < 1		0.993 < s < 1	

Table 23. Station YØ 1, 50 m

	1972		1973		1974							
	JUNE X	S	OCT. X	S	JUNE X	S	OCT. X	S	JUNE X	S	OCT. X	S
<u>Ctenodiscus crispatus</u>	0.2	0.42	0.3	0.48								
<u>Chaetoderma nitidulum</u>	1.5	1.72	0.8	0.92	2.0	1.83	0.8	0.92				
Amphipoda indet.	1.6	1.35	2.9	1.85	0.5	0.97	1.3	1.57				
<u>Calocharis macandreae</u>					0.5	0.97						
Cumacea indet.	2.8	2.15	2.7	1.83	4.4	2.76	0.8	1.75				
<u>Gnathia oxyraea</u>					0.1	0.32						
<u>Brisaster fragilis</u>	0.1	0.32			0.1	0.32						
<u>Admete viridula</u>							0.1	0.32				
<u>Cylichna alba</u>	0.2	0.42										
<u>Philine quadrata</u>			0.1	0.32	0.1	0.32						
<u>Philine scabra</u>					0.2	0.42						
<u>Retusa umbilicata</u>					0.1	0.32						
<u>Taranis moerchi</u>	0.1	0.32										
<u>Trophonopsis barvicensis</u>	0.1	0.32										
Turridae indet.					0.1	0.32						
Holothuroidea indet.	0.1	0.32	0.1	0.32	0.1	0.32	1.3	1.25				
Nemertini indet.			1.5	1.08	1.7	1.06	1.7	0.82				
<u>Amphiura chiajei</u>	6.6	1.65	6.4	3.31	6.9	2.38	2.3	1.42				
<u>Amphiura filiformis</u>	0.8	1.87	1.1	1.52			1.5	0.53				
<u>Amphiura</u> sp.			0.2	0.63	0.2	0.63						
<u>Ophiura affinis</u>							0.1	0.32				
<u>Ophiura albida</u>	0.7	0.82	0.1	0.32								
<u>Ophiura sarsi</u>	0.2	0.42	3.0	2.11			0.1	0.32				
<u>Ophiura</u> sp.			0.4	0.70	2.0	1.63	0.8	0.79				
Ampharetidae indet.	0.2	0.42					0.3	0.67				
<u>Ampharetidae</u> sp.					0.1	0.32						
<u>Amphicteis gunneri</u>	0.1	0.32										
<u>Anaitides</u> sp.			0.2	0.42								
<u>Anobothrus gracilis</u>							0.2	0.42				
<u>Aphrodita aculeata</u>					0.1	0.32						
<u>Asychis biceps</u>	0.3	0.67	1.1	1.45	0.8	0.63	0.5	0.71				
<u>Ceratocephale loveni</u>	0.1	0.32			0.2	0.63						
<u>Chaetozone setosa</u>	0.8	0.79	4.4	3.31	0.8	1.32	1.8	2.25				
<u>Cirratulus cirratus</u>	0.2	0.63	0.3	0.48	0.1	0.32	0.1	0.32				
<u>Dasybranchus caducus</u>	0.1	0.32	0.2	0.42	0.1	0.32						
<u>Diplocirrus glaucus</u>	0.2	0.42	0.9	0.74	0.3	0.67	0.2	0.42				
<u>Driloneris filum</u>			0.2	0.42	0.2	0.63	0.1	0.32				
<u>Euchone rubrocincta</u>	0.1	0.32										
Euclymeninae indet.			0.2	0.42								
<u>Glycera alba</u>			0.1	0.32	0.1	0.32	0.1	0.32				
<u>Glycera rouxi</u>	0.1	0.32			0.2	0.63	0.2	0.42				
<u>Glyphanostomum pallescens</u>			0.1	0.32								
<u>Goniada maculata</u>					0.2	0.42						
<u>Heteromastus filiformis</u>			0.2	0.63	0.1	0.32	0.2	0.42				
<u>Laonice cirrata</u>			1.2	1.03			0.4	0.70				

Table 23. cont.

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s
<u>Lumbrineris fragilis</u>	0.9	0.88	1.1	1.37	0.6	0.70	1.1	1.37				
<u>Lumbriclymene minor</u>	0.1	0.32										
Maldanidae indet.									0.2	0.42		
<u>Maldane sarsi</u>	0.2	0.63			0.1	0.32						
<u>Melinna cristata</u>	0.8	1.03	0.5	0.85	0.1	0.32	4.4	5.93				
<u>Myriochela sp.</u>	0.1	0.32	0.6	0.70	0.3	0.67						
<u>Nephrys ciliata</u>	0.1	0.32							0.1	0.32		
<u>Nephys paradoxica</u>					0.4	0.70						
<u>Notomastus latericeus</u>	0.1	0.32	0.3	0.48	0.4	1.26	0.2	0.63				
<u>Ophelina norvegica</u>									0.1	0.32		
<u>Ophelina</u> sp.					0.1	0.32						
<u>Ophiocnemus flexuosus</u>			0.2	0.42	0.1	0.32	0.2	0.42				
<u>Owenia fusiformis</u>			0.2	0.42			1.8	3.16				
<u>Paramphinoe jeffreysi</u>	1.2	1.23	1.6	1.17	0.8	1.75	0.1	0.32				
<u>Pectinaria auricoma</u>							0.1	0.32				
<u>Pectinaria koreni</u>	0.1	0.32			0.3	0.48	0.6	1.26				
<u>Pholoe minuta</u>	3.2	2.10	2.0	2.36			2.2	1.87				
Phyllodocidae indet.							0.1	0.32				
<u>Phyllodocinae</u> indet.	0.1	0.32					0.1	0.32				
<u>Phylo kupfferi</u>												
<u>Phylo norvegicus</u>			0.3	0.67	0.3	0.67			0.1	0.32		
<u>Polycirrus</u> sp.												
Polynoidae Indet.			0.1	0.32	0.5	0.71	0.2	0.42				
<u>Polypysia crassa</u>	1.4	1.43	0.4	0.70								
<u>Rhodine loveni</u>					0.7	0.67						
<u>Sabellides borealis</u>			0.5	1.08								
Sabellidae indet.	0.4	0.97	0.1	0.32								
<u>Sabella penicillatus</u>			0.2	0.63			0.7	1.97				
<u>Scoloplos armiger</u>	0.1	0.32										
<u>Sphaerodorum gracilis</u>	0.1	0.32					0.2	0.42				
<u>Streblosoma bairdi</u>			0.4	0.70			0.2	0.63				
Syllidae indet.	0.3	0.48	0.2	0.63	0.2	0.42	0.2	0.42				
Terebellidae indet.	0.2	0.42			0.4	0.70						
<u>Terebellides stroemii</u>	0.1	0.32	0.4	0.70			0.1	0.32				
<u>Thelodus cincinnatus</u>			0.1	0.32								
<u>Trichobranchus roseus</u>							0.2	0.42				
<u>Abra alba</u>	7.1	4.33	8.7	5.96	30.6	18.64	16.1	10.80				
<u>Abra nitida</u>	5.8	3.22			4.4	3.41	2.5	2.27				
<u>Astarte sulcata</u>							0.4	0.70				
Cardiidae indet.	0.1	0.32										
<u>Chlamys suciata</u>	0.1	0.32										
<u>Cymidaria</u> sp.					0.1	0.32						
<u>Ennucula tenuis</u>	0.1	0.32										
<u>Heteranomia squamula</u>					0.1	0.32						
<u>Hiatella arctica</u>	0.1	0.32	0.1	0.32								
<u>Kellellia miliaris</u>			4.6	4.27	4.8	3.68						
<u>Limatula sulcata</u>			0.1	0.32	0.5	0.71	0.2	0.42				
<u>Macoma calcarea</u>			0.1	0.32								
<u>Modiolus modiolus</u>			0.1	0.32								
<u>Montacuta</u> sp.	0.1	0.32										
<u>Mysella bidentata</u>	0.1	0.32	0.2	0.42	0.3	0.48						
<u>Palliolium vitreum</u>	0.1	0.32										
<u>Parvicardium minimum</u>			1.2	0.42	0.5	0.53	4.9	3.81				
<u>Thyasira</u> sp.	6.9	4.70	9.0	5.01	15.3	3.47	1.5	1.35				
<u>Yoldiella</u> sp.	0.1	0.32	1.2	1.03	1.9	1.45						
<u>Phoronis müllerii</u>							0.1	0.32				
<u>Nymphon</u> sp.			0.1	0.32								
Pycnocionidae indet.			0.1	0.32								
<u>Dentalium entale</u>							0.1	0.32				
<u>Dentalium occidentale</u>							0.1	0.32	0.1	0.32		
<u>Dentalium</u> sp.			0.1	0.32								
<u>Onchonesoma steenstrupi</u>	6.9	4.91	7.0	5.58	16.6	5.82	0.5	0.97				
<u>Phascolion strombi</u>	0.5	0.71	0.3	0.48	0.1	0.32						
<u>Phascolosoma</u> sp.	0.1	0.32										
<u>Ascidia conchilega</u>	0.1	0.32										
Polycladida indet.							0.1	0.32				
	λ	0.92		0.94		0.85		0.89				
Diversity:	H	2.90		3.15		2.50		2.92				
	d	8.72		8.69		7.93		8.57				
Sampling efficiency:		0.931 < s < 1		0.968 < s < 1		0.972 < s < 1		0.952 < s < 1				

Table 24. Station VD 1, 20 m

	JUNE		OCT.	
	\bar{x}	s	\bar{x}	s
<u>Virgularia mirabilis</u>	0.5	0.71	2.1	1.73
<u>Ctenodiscus crispatus</u>	0.3	0.48	0.1	0.32
<u>Chaetoderma nitidulum</u>	0.5	0.85	0.1	0.32
Amphipoda indet.	0.3	0.48	0.9	0.74
Cumacea indet.	0.1	0.32		
Holothuroidea indet.			0.1	0.32
Nemertini indet.	2.0	0.94	1.4	0.97
<u>Amphiura chiajei</u>	5.0	2.75	11.5	3.10
<u>Amphiura filiformis</u>	0.3	0.48	1.5	1.51
<u>Amphiura</u> sp.			1.1	2.51
<u>Ophiura albida</u>	2.3	1.64	4.3	2.79
<u>Ophiura sarsi</u>	0.3	0.67		
<u>Ophiura</u> sp.			1.0	1.56
<u>Ophiura texturata</u>	0.2	0.42		
<u>Amphitrite cirrata</u>	0.2	0.42	0.2	0.63
<u>Anaitides</u> sp.	0.1	0.32	0.6	0.70
<u>Anobothrus gracilis</u>			0.1	0.32
<u>Aphrodisia aculeata</u>			0.1	0.32
<u>Artacama proboscidea</u>			0.2	0.42
<u>Chaetozone setosa</u>	0.3	0.67		
<u>Cirratulus cirratus</u>	0.1	0.32	0.2	0.42
<u>Dasybranchus caducus</u>			0.1	0.32
<u>Diplocirrus glaucus</u>	1.8	2.35	0.1	0.32
Euclymeninae indet.			0.7	0.95
<u>Gattyana cirrosa</u>			0.1	0.32
<u>Glycera alba</u>	0.9	0.88	0.4	0.52
<u>Glycera rouxii</u>	0.6	0.70	0.1	0.32
<u>Goniada maculata</u>	1.4	0.97	1.0	1.05
<u>Heteromastus filiformis</u>	1.2	0.92	0.1	0.32
<u>Lanassa nordenskiöldi</u>			0.5	0.71
<u>Lanassa venusta</u>			0.4	0.70
<u>Laonice cirrata</u>	0.8	1.03	0.7	0.67
<u>Lumbrineris fragilis</u>	3.5	2.22	2.5	1.51
<u>Maldane sarsi</u>	7.0	6.06	3.3	4.72
<u>Melinna cristata</u>			0.1	0.32
<u>Myriochele</u> sp.	0.9	0.88	0.6	0.84
<u>Neoamphitrite figulus</u>			0.4	0.97
<u>Nephtys ciliata</u>	0.6	1.07	0.6	1.07
<u>Nephtys paradoxa</u>	0.2	0.42	0.2	0.42
<u>Nereis virens</u>			0.1	0.32
<u>Ophelina acuminata</u>	0.3	0.67	0.2	0.42
<u>Ophiodromus flexuosus</u>	0.6	0.52	1.0	1.05
<u>Paranonis gracilis</u>	0.5	1.27		
<u>Pectinaria koreni</u>	0.2	0.42	0.1	0.32
<u>Pholoe minuta</u>	0.3	0.48	0.3	0.48
Polynoidae indet.			0.1	0.32

Table 24. cont.

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{X}	s	\bar{X}	s	\bar{X}	s	\bar{X}	s	\bar{X}	s	\bar{X}	s
<u>Polyphysia crassa</u>	0.1	0.32	0.1	0.32								
<u>Praxillella gracilis</u>	1.0	1.05	0.5	0.85								
<u>Rhodine gracilior</u>	0.1	0.32										
Sabellidae indet.				0.4			0.52					
<u>Scalibregma inflatum</u>	0.1	0.32	0.3	0.48								
<u>Scoloplos armiger</u>	0.1	0.32										
Spionidae indet.	1.7	1.16	10.2	5.31								
<u>Streblosoma bairdi</u>	0.8	1.23	0.5	0.71								
Syllidae indet.	0.1	0.32	0.4	0.70								
<u>Terebellides stroemii</u>	0.2	0.42	0.2	0.42								
<u>Trichobranchus roseus</u>	0.2	0.63	1.1	1.29								
<u>Trochochaeta multisetosa</u>	0.2	0.42										
 <u>Abra alba</u>	58.8	22.91	5.2	3.49								
<u>Abra nitida</u>	9.5	5.93	6.0	3.74								
<u>Corbula gibba</u>	4.8	2.25	49.8	20.83								
<u>Hiatella arctica</u>	0.3	0.48	1.4	1.51								
<u>Montacuta</u> sp.			0.1	0.32								
<u>Montacuta tenella</u>			0.1	0.32								
<u>Musculus niger</u>			0.1	0.32								
<u>Mya arenaria</u>			0.1	0.32								
 <u>Mysella bidentata</u>			0.2	0.42								
<u>Nuculana minuta</u>	0.1	0.32										
<u>Parvicardium minimum</u>	0.2	0.42	0.9	1.10								
<u>Thracia</u> sp.			0.2	0.42								
<u>Thyasira</u> sp.	0.8	1.03	1.9	1.52								
 <u>Phoronis mülleri</u>	0.1	0.32	0.1	0.32								
 <u>Dentalium entale</u>	0.2	0.42	0.5	0.71								
 <u>Onchnesoma steenstrupi</u>	0.6	1.26										
<u>Phascolion strombi</u>	0.1	0.32										
 Diversity:	λ	0.71		0.80								
	H	2.18		2.51								
	d	7.39		8.76								
Sampling efficiency:	0.984 < S < 1				0.976 < S < 1							

Table 25. Station VD 2, 50 m

	1974			
	JUNE		OCT.	
	\bar{X}	s	\bar{X}	s
<u>Pennatula phosphorea</u>	0.3	0.67		
<u>Virgularia mirabilis</u>	0.9	0.88	0.4	0.84
<u>Ctenodiscus crispatus</u>	0.5	0.53		
<u>Chaetoderma nitidulum</u>	1.7	2.41	0.1	0.32
Amphipoda indet.	1.4	2.07	7.1	3.54
Cumacea indet.	0.2	0.42	0.7	0.67
Holothuroidea indet.	0.2	0.42		
Nemertini indet.	1.9	1.20	0.9	0.99
<u>Amphilepis norvegica</u>	0.4	0.84		
<u>Amphiura chiajei</u>	11.2	3.29	8.1	4.75
<u>Amphiura filiformis</u>			1.0	1.15
<u>Ophiura sarsi</u>	1.0	1.15	0.2	0.63
<u>Ophiura</u> sp.	0.7	0.95	0.2	0.42
Ampharetidae indet.			0.2	0.42
<u>Anaitides</u> sp.			0.1	0.32
<u>Anobothrus gracilis</u>	0.1	0.32		
<u>Aphrodita aculeata</u>	0.1	0.32		
<u>Asychis biceps</u>	0.1	0.32		
<u>Brada villosa</u>			0.1	0.32
Capitellidae indet.	0.5	0.71		
<u>Ceratocephale loveni</u>	0.5	0.71	0.5	0.71
<u>Chaetozone setosa</u>	0.3	0.48		
<u>Diplocirrus glaucus</u>	0.7	0.95	1.8	1.14
<u>Driloneris filum</u>	1.8	1.14	1.4	1.17
<u>Glycera rouxi</u>			0.6	0.84
<u>Glycera</u> sp.	0.1	0.32		
<u>Goniada maculata</u>	0.2	0.42		
<u>Harmothoe</u> sp.			0.4	0.52
<u>Heteromastus filiformis</u>	1.9	1.37	8.2	4.61
<u>Hydroides norvegica</u>			0.1	0.32
<u>Laonice cirtata</u>			0.1	0.32
<u>Lumbrineris fragilis</u>				
<u>Maldane sarsi</u>	81.7	31.77	90.8	26.89
<u>Myriochele</u> sp.	1.6	1.71	2.2	1.62
<u>Neoamphitrite grayi</u>	0.8	2.53	0.1	0.32
<u>Nephtys ciliata</u>	0.1	0.32		
<u>Nephtys paradoxa</u>	0.4	0.70	0.3	0.48
<u>Ophelina norvegica</u>	0.2	0.42		
<u>Ophiodromus flexuosus</u>	0.8	0.79	1.4	1.35
<u>Owenia fusiformis</u>	0.1	0.32	2.8	1.99
<u>Paramphinoe jeffreysi</u>	0.3	0.67	0.1	0.32
<u>Pectinaria auricoma</u>	0.1	0.32		
<u>Pectinaria koreni</u>	0.4	0.52	0.3	0.48
<u>Pholoe minuta</u>	1.2	1.40	3.0	2.26
Phyllodocidae indet.	0.2	0.42	0.1	0.32
<u>Phylo norvegicus</u>	0.3	0.67		

Table 25. cont.

	1974		1973		1974	
	JUNE X	OCT. S	JUNE X	S	JUNE X	OCT. S
<u>Polyphysia crassa</u>	0.1	0.32				
<u>Praxillella gracilis</u>	0.2	0.42				
<u>Praxillella praetermissa</u>	0.1	0.32				
<u>Rhodine sp.</u>	0.1	0.32				
<u>Sabellidae</u> indet.	0.4	0.52				
<u>Scalibregma inflatum</u>		0.3	0.48			
<u>Sphaerodorum gracilis</u>	0.1	0.32				
<u>Streblosoma bairdi</u>	1.3	1.34				
<u>Syllidae</u> indet.	0.3	0.48				
<u>Terebellides stroemi</u>		0.1	0.32			
<u>Trochochaeta multisetosa</u>	0.2	0.42				
<u>Abra alba</u>	20.8	15.42	73.3	34.68		
<u>Abra nitida</u>	16.5	8.42	16.3	3.80		
<u>Corbula gibba</u>		0.1	0.32			
<u>Hiatella arctica</u>		0.1	0.32			
<u>Limatula sulcata</u>		0.1	0.32			
<u>Mya arenaria</u>		0.1	0.32			
<u>Mytilus edulis</u>	0.1	0.32				
<u>Parvicardium minimum</u>		0.3	0.48			
<u>Thyasira sp.</u>	1.8	1.87	0.4	0.52		
<u>Yoldiella sp.</u>		0.2	0.42			
<u>Dentalium entale</u>		0.1	0.32			
<u>Phascolion strombi</u>	0.1	0.32				
	λ	0.70	0.72			
Diversity:	H	1.94	1.77			
	d	6.93	5.31			
Sampling efficiency:	0.987 <S<1	0.991 <S<1				

Table 26. Station KV 2, 50 m

	1972				1973			
	JUNE		OCT.		JUNE		OCT.	
	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s
<u>Chaetoderma nitidulum</u>	0.5	0.71	0.3	0.67	0.2	0.42	0.9	1.20
<u>Acanthochiton fascicularis</u>							0.1	0.32
Amphipoda indet.	0.3	0.48	3.7	1.57	0.9	0.99	0.4	0.52
Cumacea indet.	0.7	1.25	3.5	1.35	0.2	0.42	0.5	0.53
Regularia indet.	0.1	0.32						
<u>Admete viridula</u>	0.1	0.32						
<u>Balcis</u> sp.							0.1	0.32
<u>Cylichna alba</u>	0.1	0.32						
<u>Eulimella scillae</u>					0.1	0.32		
<u>Clelandella miliaris</u>							0.2	0.63
<u>Lunatia pallida</u>	0.1	0.32						
<u>Philine quadrata</u>			0.1	0.32				
Holothuroidea indet.					0.2	0.42	0.1	0.32
Nemertini indet.			0.4	0.70	0.4	0.52	0.7	0.82
<u>Amphiura chiajei</u>	0.1	0.32	0.3	0.48	0.3	0.48	0.1	0.32
<u>Amphiura filiformis</u>	0.1	0.32	1.5	2.76			0.4	0.70
<u>Amphiura</u> sp.			0.5	0.71	0.1	0.32		
<u>Ophiura affinis</u>							0.3	0.67
<u>Ophiura albida</u>	0.2	0.42						
Ophiuroidea indet.	0.1	0.32						
<u>Ophiura sarsi</u>			0.4	0.70	0.3	0.67		
<u>Ophiura</u> sp.							0.1	0.32
Ampharetidae indet.					0.2	0.42	0.4	0.70
<u>Amphicteis gunneri</u>	0.2	0.63	0.2	0.42	0.1	0.32		
<u>Anaitides</u> sp.	0.1	0.32	0.1	0.32	0.4	0.70	0.1	0.32
<u>Aphrodita aculeata</u>			0.1	0.32				
<u>Asychis biceps</u>	23.2	10.84	25.9	11.31	32.0	7.51	11.4	9.67
<u>Brada villosa</u>	0.1	0.32						
<u>Chaetozone setosa</u>	0.3	0.67	2.8	1.81	0.3	0.48	0.3	0.48
<u>Diplocirrus glaucus</u>	0.4	0.52	1.4	0.84	0.5	0.71	0.6	0.70
<u>Driloneris filum</u>					0.1	0.32		
<u>Eclysippe vanelli</u>			0.1	0.32				
Euclymeninae indet.					0.3	0.67	0.4	0.52
<u>Glycera alba</u>							0.2	0.63
<u>Glycera lapidum</u>	0.1	0.32						
<u>Glycera rouxii</u>	0.1	0.32			0.3	0.67	0.1	0.32
<u>Glycera</u> sp.			0.6	0.70				
<u>Goniada maculata</u>	0.1	0.32	0.1	0.32				
<u>heteromastus filiformis</u>	0.2	0.63	1.3	1.34	0.1	0.32	0.2	0.42
<u>Laetmonice filicornis</u>			0.1	0.32				
<u>Laonice cirrata</u>	0.4	0.70	1.2	1.14			0.2	0.42
<u>Lumbrineris fragilis</u>	0.3	0.48	1.2	1.32	1.8	0.92	0.7	0.67
Maldanidae indet.							0.2	0.63
<u>Maldane sarsi</u>					0.2	0.63		

Table 26. cont.

	1972				1973			
	JUNE		OCT.		JUNE		OCT.	
	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s
<u>Melinna cristata</u>	3.2	3.29	1.5	1.35	8.0	4.11	10.4	7.34
<u>Microclymene tricirrata</u>			0.2	0.42	0.1	0.32		
<u>Myriochele</u> sp.	0.5	1.58	0.4	0.70	0.3	0.48	0.9	0.99
<u>Nephtys ciliata</u>	0.1	0.32	0.1	0.32			0.1	0.32
<u>Nephtys paradoxa</u>			0.1	0.32				
<u>Nephtys</u> sp.					0.1	0.32		
<u>Nothria conchylega</u>			0.2	0.42			0.1	0.32
<u>Notomastus latericeus</u>	0.5	0.97	1.6	0.97	0.2	0.42	1.0	0.67
<u>Ophelina acuminata</u>							0.1	0.32
<u>Ophelina norvegica</u>			0.2	0.42				
<u>Ophiodromus flexuosus</u>					0.2	0.42	0.2	0.63
<u>Owenia fusiformis</u>							0.3	0.95
<u>Paramphinome jeffreysi</u>			2.1	2.73	0.1	0.32	0.2	0.63
<u>Paraonis gracilis</u>					0.1	0.32		
<u>Pectinaria koreni</u>	0.1	0.32	0.2	0.63	0.2	0.63	0.2	0.42
<u>Pholoe minuta</u>			0.4	0.84	0.1	0.32	0.4	0.84
Phyllodocidae indet.					0.1	0.32	0.1	0.32
<u>Phylo norvegicus</u>			0.1	0.32				
Polynoidae indet.	0.2	0.42	0.2	0.42			0.3	0.67
<u>Praxillella praetermissa</u>	0.1	0.32	0.6	0.84				
<u>Rhodine gracilior</u>					0.2	0.42		
<u>Rhodine loveni</u>			0.3	0.67			0.1	0.32
Sabellidae indet.					0.1	0.32		
<u>Sabella penicilllus</u>	0.6	1.07	0.4	0.52			0.2	0.42
<u>Samytha sexcirtata</u>			0.1	0.32				
<u>Sphaerodororum gracilis</u>			0.1	0.32				
<u>Spiophanes krøyeri</u>			0.1	0.32				
Syllidae indet.					0.2	0.42	0.2	0.42
<u>Terebellides stroemi</u>	0.2	0.42	0.4	0.52			0.4	0.52
<u>Thelepus cincinnatus</u>			0.2	0.42			0.3	0.67
<u>Trichobranchus roseus</u>							0.5	1.08
<u>Typosyllis cornuta</u>	0.2	0.42						
<u>Abra alba</u>	0.2	0.42	0.2	0.63	0.2	0.42	0.5	0.97
<u>Abra nitida</u>	6.7	4.32	10.0	5.75	14.4	4.03	3.7	4.55
<u>Bathycarca pectunculoides</u>	0.2	0.42	0.2	0.42	0.1	0.32		
Cardiidae indet.	0.8	0.79						
<u>Ennucula tenuis</u>					0.1	0.32		
<u>Hiatella arctica</u>	0.1	0.32						
<u>Kelliella miliaris</u>			2.8	2.57	0.9	1.60	0.1	0.32
<u>Limatula</u> sp.	2.7	1.70						
<u>Limatula sulcata</u>			1.0	0.82	1.2	1.14	1.8	2.39
<u>Modiolula phaseolinus</u>	0.2	0.42	0.7	1.25	0.8	0.92		
<u>Nuculana minuta</u>	0.1	0.32	0.1	0.32	0.1	0.32		
<u>Nucula tumidula</u>			0.1	0.32				
<u>Palliolium vitreum</u>	0.1	0.32						
<u>Parvicardium minimum</u>			2.1	1.10	0.7	1.06	0.9	1.37
<u>Similipecten similis</u>							0.1	0.32
<u>Thyasira</u> sp.	3.1	2.64	12.9	5.99	14.8	5.71	3.2	2.39
<u>Yoldiella</u> sp.	3.1	1.52	7.1	1.91	8.4	4.25	2.2	1.87
<u>Dentalium entale</u>							0.4	0.97
<u>Dentalium</u> sp.			0.2	0.63				
<u>Onchnesoma squamatum</u>					0.1	0.32		
<u>Onchnesoma steenstrupi</u>	0.1	0.32	3.7	2.06	1.3	1.16		
<u>Phascolion strombi</u>	0.2	0.42			0.1	0.32		
Diversity:	λ	0.76		0.89		0.81		0.88
	H	2.22		2.81		2.20		2.80
	d	7.21		8.01		7.03		8.27
Sampling efficiency:	0.946	$\leq S \leq 1$	0.978	$\leq S \leq 1$	0.973	$\leq S \leq 1$	0.955	$\leq S \leq 1$

Table 27. Station KV 3, 100 m

	1972				1973			
	JUNE		OCT.		JUNE		OCT.	
	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s
<u><i>Stylatula elegans</i></u>								
<u><i>Chaetoderma nitidulum</i></u>	0.5	0.71	0.3	0.67	0.8	0.92	0.4	0.52
<u><i>Amphipoda</i> indet.</u>	0.1	0.32	2.4	2.91	0.1	0.32		
<u><i>Calocharis macandreae</i></u>	0.3	0.67	0.2	0.63				
<u><i>Cumacea</i> indet.</u>	0.2	0.42	0.1	0.32	0.1	0.32		
<u><i>Geryon tridens</i></u>			0.2	0.42	0.1	0.32	0.1	0.32
<u><i>Iothia fulva</i></u>					0.1	0.32		
<u><i>Philine quadrata</i></u>			0.2	0.42				
<u><i>Retusa umbilicata</i></u>			0.1	0.32				
<u><i>Skenea basistriata</i></u>					0.1	0.32		
<u><i>Nemertini</i> indet.</u>			0.8	1.03	0.5	0.53	0.6	0.84
<u><i>Amphilepis norvegica</i></u>	3.0	2.05	4.1	3.00	5.5	3.75		
<u><i>Amphiura chiajei</i></u>			0.2	0.42				
<u><i>Amphiura filiformis</i></u>						0.6	1.26	
<u><i>Amphiura</i> sp.</u>	0.1	0.32			0.2	0.42		
<u><i>Ophiura</i> sp.</u>								
<u><i>Ampharetidae</i> indet.</u>						0.1	0.32	
<u><i>Amphicteis gunneri</i></u>	0.2	0.42			0.1	0.32		
<u><i>Asychis biceps</i></u>	0.6	0.84	0.6	0.84	1.2	1.23	0.6	0.52
<u><i>Brada villosa</i></u>	0.4	0.70	0.1	0.32	0.1	0.32		
<u><i>Dasybranchus caducus</i></u>			0.2	0.42	0.5	0.53	0.5	0.71
<u><i>Diplocirrus glaucus</i></u>					0.5	0.71	0.1	0.32
<u><i>Driloneris filum</i></u>					0.1	0.32		
<u><i>Eclipsippe vanelli</i></u>			0.1	0.32				
<u><i>Euclymeninae</i> indet.</u>	1.3	1.16			2.2	2.39	0.4	0.52
<u><i>Eunice pennata</i></u>	0.1	0.32	0.3	0.48	0.1	0.32		
<u><i>Goniada maculata</i></u>						0.2	0.42	
<u><i>Heteromastus filiformis</i></u>					0.2	0.63	0.1	0.32
<u><i>Leiochone borealis</i></u>			2.8	3.16				
<u><i>Lumbrineris fragilis</i></u>	0.4	0.70	0.7	0.67	0.3	0.48	0.2	0.42
<u><i>Maldane sarsi</i></u>	0.6	1.35			0.2	0.63	0.2	0.63
<u><i>Melinna cristata</i></u>	0.3	0.67			0.9	1.66	0.1	0.32
<u><i>Myriochele</i> sp.</u>	0.3	0.48	3.6	7.35	0.8	2.20	0.2	0.42
<u><i>Nephtys ciliata</i></u>			0.1	0.32				
<u><i>Nephtys incisa</i></u>	0.2	0.42				0.1	0.32	
<u><i>Nephtys paradoxa</i></u>						0.3	0.67	
<u><i>Nephtys</i> sp.</u>					0.2	0.42		
<u><i>Ophelina norvegica</i></u>			0.1	0.32				
<u><i>Paramphinome jeffreysi</i></u>	0.6	0.70	0.3	0.48	0.9	0.74	0.2	0.42
<u><i>Pholoe minuta</i></u>	0.3	0.67	0.1	0.32	0.4	0.52		
<u><i>Phyllodocidae</i> indet.</u>					0.1	0.32		
<u><i>Phylo norvegicus</i></u>			0.9	0.99	0.7	0.82	0.4	0.52
<u><i>Polynoidae</i> indet.</u>							0.1	0.32
<u><i>Polyphysia crassa</i></u>					0.1	0.32		
<u><i>Rhodine loveni</i></u>	1.9	1.60	0.9	0.99			1.5	1.51
<u><i>Rhodine</i> sp.</u>					0.8	0.79		

Table 27. cont.

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s
Sabellidae indet.									0.1	0.32		
<u>Scoloplos armiger</u>	0.7	0.67			0.1	0.32						
Spionidae indet.												
<u>Streblosoma bairdi</u>	0.1	0.32			0.2	0.42			0.2	0.42		
<u>Steblosoma intestinalis</u>												
<u>Terebellides stroemii</u>	0.1	0.32			0.2	0.42						
<u>Thelepus cincinnatus</u>												
Thelepinae indet.							0.4	0.70				
<u>Abra nitida</u>	1.8	1.99	3.9		2.81	1.8	1.62	0.7	0.95			
<u>Astarte sulcata</u>							0.3	0.48				
<u>Bathyarca pectunculoides</u>	0.2	0.42	0.1		0.32	0.7	0.95					
<u>Chlamys sulcata</u>							0.1	0.32				
<u>Cuspidaria</u> sp.					0.1	0.32			0.1	0.32		
<u>Kelliella miliaris</u>					1.1	2.42	1.0	1.33	0.1	0.32		
<u>Limatula</u> sp.	0.1	0.32										
<u>Nucula nucleus</u>									0.5	0.71		
<u>Nucula</u> sp.	0.4	0.52										
<u>Nucula tumidula</u>					0.6	0.70	0.4	0.52				
<u>Palliolum vitreum</u>	0.1	0.32					0.7	1.25				
Pectinidae indet.							0.1	0.32	0.1	0.32		
<u>Pseudamussium septemradiatum</u>							0.1	0.32				
<u>Thyasira</u> sp.	0.4	0.84	1.9		1.60	1.4	2.07	0.5	0.71			
<u>Tropidomya abbreviata</u>									0.1	0.32		
<u>Yoldiella</u> sp.	2.1	2.23	2.2		2.53	3.9	3.38	0.5	1.27			
<u>Phoronis mülleri</u>					0.1	0.32						
<u>Entalina guinguangularis</u>							0.1	0.32				
<u>Onchnesoma squatum</u>							0.2	0.42				
<u>Onchnesoma steenstrupi</u>	19.5	10.81	21.4		11.91	25.0	10.31	2.9	3.18			
Polycladia indet.							0.1	0.32				
Diversity:	λ	0.70			0.80		0.77		0.92			
	H	2.04			2.34		2.37		2.94			
	d	4.91			5.61		7.14		6.39			
Sampling efficiency:	0.962 < s < 1	0.965 < s < 1			0.956 < s < 1		0.850 < s < 1					

Table 28. Station KV 4, 208 m

	1972				1973				
	JUNE		OCT.		JUNE		OCT.		
	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	
<u>Stylatula elegans</u>	0.1	0.32	0.2	0.42	0.4	0.97			
<u>Virgularia mirabilis</u>							0.1	0.32	
Amphipoda indet.	3.9	3.18	2.1	3.14	0.4	0.52	1.1	1.45	
<u>Calocharis macandrense</u>	0.2	0.42			0.1	0.32	0.1	0.32	
Cumacea indet.			0.1	0.32	0.2	0.63			
<u>Geryon tridens</u>	0.1	0.32	0.2	0.42					
Nemertini indet.			0.2	0.42	0.6	0.84	0.7	1.06	
<u>Amphilepis norvegica</u>	2.0	1.76	3.3	2.62	4.2	1.55			
<u>Amphiura filiformis</u>							0.4	0.70	
<u>Ophiura marsi</u>	0.2	0.63							
<u>Ophiura sp.</u>	0.1	0.32	0.2	0.63					
<u>Amage auricula</u>			0.1	0.32					
<u>Amphictesis gunneri</u>	0.1	0.32							
<u>Asynchis biceps</u>	0.4	0.70	0.7	0.95	0.9	0.74	0.8	0.92	
<u>Brachia villosa</u>	0.2	0.42			0.3	0.48			
<u>Chaetozone setosa</u>					0.2	0.63	0.1	0.32	
<u>Dasybranchus caducus</u>					0.1	0.32	0.1	0.32	
<u>Diplocirrus glaucus</u>	0.1	0.32							
Euclymenidae indet.	0.1	0.32	0.2	0.42					
<u>Glycera</u> sp.					0.1	0.32	0.1	0.32	
<u>Goniada maculata</u>							0.1	0.32	
<u>Heteromastus filiformis</u>	0.8	1.03	0.2	0.42	0.1	0.32	0.1	0.32	
<u>Leptochone borealis</u>					0.4	0.52			
<u>Lumbrineris fragilis</u>	0.6	0.52	1.8	1.03	1.0	0.82	0.6	0.84	
Maldanidae indet.			0.1	0.32					
<u>Myriochela</u> sp.	0.3	0.67	0.3	0.48	0.9	2.51	6.8	6.65	
<u>Nephtys incisa</u>	0.2	0.42			0.3	0.48	0.2	0.42	
<u>Nephtys paradoxa</u>	0.1	0.32	0.4	0.70					
<u>Omphis quadruscuspis</u>			0.1	0.32					
<u>Ophelina norvegica</u>							0.1	0.32	
<u>Paramphionome jeffreysi</u>	0.2	0.42	0.1	0.32	0.1	0.32			
<u>Pectinaria auricoma</u>					0.1	0.32			
<u>Pholoe minuta</u>	0.3	0.67			0.1	0.32			
<u>Phylo norvegicus</u>			0.9	0.88	0.3	0.48	1.1	1.45	
<u>Pista cristata</u>					0.1	0.32			
Polynoidae indet.							0.2	0.63	
<u>Polyphysia crassa</u>	0.2	0.42	0.2	0.42	0.2	0.42	0.4	0.84	
Rhodine loveni	0.4	0.77	0.5	0.71	0.5	0.71	0.7	0.48	
<u>Scoleoplos armiger</u>	1.6	0.84							
<u>Sphaerodorum gracilis</u>					0.1	0.32			
<u>Spirochaetopterus</u>			0.3	0.95					
Terebellidae indet.					0.1	0.32			
<u>Terebellides stroemi</u>			0.1	0.32	0.1	0.32			
<u>Abra nitida</u>	9.7	5.52	4.4	1.96	4.2	2.53	2.9	1.29	
Cardiidae indet.	0.3	0.48							
<u>Kellia</u> sp.	0.1	0.32							
<u>Kelliella miliaris</u>			0.2	0.42	0.5	0.71			
<u>Nucula nucleus</u>							0.5	0.53	
<u>Nucula</u> sp.	2.2	1.62							
<u>Nucula tumidula</u>			1.6	1.78	2.8	1.75			
<u>Pallidium vitreum</u>	0.1	0.32			0.2	0.63			
<u>Thyasira</u> sp.	3.5	2.46	1.3	1.42	2.3	2.00	1.2	0.92	
<u>Yoldieilla</u> sp.	1.6	1.35	0.7	0.82	1.3	1.16	0.9	0.88	
<u>Phoronis mülleri</u>					0.1	0.32			
<u>Entalia quinquangularis</u>					0.3	0.67			
<u>Oncinnesoma steenstrupi</u>	14.3	7.86	7.4	6.19	10.8	4.37	2.3	4.30	
Diversity:	λ	0.83		0.87		0.86		0.86	
	H	2.26		2.53		2.54		2.43	
	d	4.93		4.97		5.99		4.28	
Sampling efficiency:	0.964	$\leq S \leq 1$	0.957	$\leq S \leq 1$	0.944	$\leq S \leq 1$	0.931	$\leq S \leq 1$	

Table 29. Station SK 1, 20 m

	JUNE 1972		OCT.		JUNE 1973		OCT.		JUNE 1974		OCT.	
	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s	\bar{x}	s
<u><i>Virgularia mirabilis</i></u>									0.5	0.71		
<u><i>Astroidea</i> indet.</u>	0.1	0.32										
<u><i>Ctenodiscus crispatus</i></u>									0.1	0.32		
<u><i>Chaetoderma nitidulum</i></u>					0.4	0.52	0.3	0.67	0.7	1.06	0.8	0.92
<u><i>Amphipoda</i> indet.</u>	1.1	1.10	0.5	1.27	0.9	1.10				0.5	1.27	
<u><i>Cumacea</i> indet.</u>	0.2	0.63							0.2	0.42		
<u><i>Echinocardium cordatum</i></u>							0.1	0.32				
<u><i>Echinocardium flavescent</i></u>	0.1	0.32										
<u><i>Aporrhais pespelican</i></u>	0.1	0.32			0.2	0.63						
<u><i>Buccinum</i> sp.</u>			0.1	0.32					0.1	0.32		
<u><i>Buccinum undatum</i></u>	0.5	0.53			0.1	0.32						
<u><i>Menestho divisa</i></u>					0.1	0.32						
<u><i>Philine scabra</i></u>							0.2	0.42				
<u><i>Holothuroidea</i> indet.</u>	0.1	0.32			0.8	1.32	1.0	1.89	1.5	1.18	0.3	0.48
<u><i>Nemertini</i> indet.</u>			0.5	0.53	0.8	0.79	0.5	0.71	1.3	1.06	0.3	0.48
<u><i>Amphiura chiajei</i></u>			0.3	0.48	0.3	0.67	0.1	0.32	0.9	0.74	0.6	0.97
<u><i>Amphiura filiformis</i></u>	0.1	0.32	0.1	0.32	0.5	0.71	0.3	0.48	0.7	1.06	0.7	1.06
<u><i>Ophiura albida</i></u>							1.0	1.33	0.1	0.32	1.6	0.97
<u><i>Ophiura robusta</i></u>							0.2	0.42				
<u><i>Ophiura sarsi</i></u>			0.2	0.42	0.2	0.63						
<u><i>Ophiura</i> sp.</u>			0.1	0.32	1.8	1.40	0.3	0.48	0.5	0.85	0.5	0.85
<u><i>Ophiura texturata</i></u>							0.1	0.32				
<u><i>Ampharetidae</i> indet.</u>									0.2	0.42		
<u><i>Amphicteis gunneri</i></u>			0.1	0.32								
<u><i>Anaitides</i> sp.</u>	0.6	0.52	0.8	0.79	0.2	0.42	0.5	0.71			0.1	0.32
<u><i>Anobothrus gracilis</i></u>									0.1	0.32		
<u><i>Aphrodita aculeata</i></u>					0.1	0.32						
<u><i>Asychnis biceps</i></u>							0.1	0.32				
<u><i>Brada villosa</i></u>			0.2	0.63	0.3	0.67			0.3	0.67	0.2	0.63
<u><i>Chaetozone setosa</i></u>	0.1	0.32									0.2	0.63
<u><i>Cirratulus cirratus</i></u>			0.1	0.32					0.6	0.97		
<u><i>Diplocirrus glaucus</i></u>	0.3	0.48			0.2	0.42					1.5	3.47
<u><i>Euchone papillosa</i></u>									0.1	0.32	0.1	0.32
<u><i>Glycera alba</i></u>	1.0	0.82	0.8	0.79	0.9	1.45	1.1	1.29	1.0	0.67	0.5	0.53
<u><i>Glycera rouxi</i></u>									0.1	0.32		
<u><i>Goniada maculata</i></u>	1.0	1.05	1.4	1.07	1.9	1.29	1.2	0.63	1.2	0.79	1.5	0.97
<u><i>Heteromastus filiformis</i></u>											0.7	1.89
<u><i>Lanassa venusta</i></u>									0.5	0.71	0.1	0.32
<u><i>Laonice cirrata</i></u>	0.4	0.97	1.7	1.25	0.2	0.63	0.8	0.92	2.2	1.55	1.7	0.95
<u><i>Lumbrineris fragilis</i></u>	0.6	0.70	0.8	0.79	0.8	1.03	0.9	1.37	2.1	2.02	0.9	0.88
<u><i>Maldanidae</i> indet.</u>									0.4	0.52		
<u><i>Maldane sarsi</i></u>	0.2	0.42			1.6	2.22	0.2	0.63	9.1	4.86	1.1	1.52
<u><i>Myriochele</i> sp.</u>	16.3	13.52			15.0	12.04	10.9	11.69	15.6	15.89	20.0	18.47
<u><i>Neoamphitrite grayi</i></u>							0.1	0.32	0.1	0.32		

Table 29. cont.

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{x}	s										
<u>Nephtys ciliata</u>	0.1	0.32	0.1	0.32	0.3	0.48	0.4	0.52	0.3	0.48	0.6	0.84
<u>Nephtys hombergi</u>	0.1	0.32										
<u>Nephtys incisa</u>									0.1	0.32		
<u>Nephtys paradoxa</u>							0.1	0.32			0.3	0.67
<u>Nereimyra punctata</u>			0.1	0.32			0.1	0.32				
<u>Nereis</u> sp.					0.1	0.32						
<u>Nereis virens</u>	0.6	0.70	0.3	0.48								
<u>Ophelina acuminata</u>			0.2	0.42			0.5	0.97	0.3	0.48	0.1	0.32
<u>Ophiodromus flexuosus</u>			0.1	0.32			0.6	0.97	0.6	0.70	0.3	0.48
<u>Owenia fusiformis</u>			0.1	0.32					1.0	1.41	0.2	0.63
<u>Paramphinome jeffreysi</u>											0.4	1.26
<u>Paraonis gracilis</u>			0.1	0.32					0.1	0.32		
<u>Pectinaria auricoma</u>					0.3	0.67			0.7	0.95	0.1	0.32
<u>Pectinaria koreni</u>	0.9	1.45	0.9	1.29	1.7	1.42	1.3	1.06	1.6	0.97	11.7	8.92
<u>Pholoe minuta</u>	3.9	3.38	8.7	4.22	0.9	0.74	2.1	2.38	0.5	0.71	5.0	4.35
Phyllodocidae indet.											0.1	0.32
<u>Rhodine gracilior</u>			0.2	0.42					4.5	4.09	0.9	0.88
<u>Rhodine</u> sp.					0.5	0.71	0.1	0.32				
Sabellidae indet.							0.1	0.32				
<u>Scalibregma inflatum</u>											0.1	0.32
<u>Scoloplos armiger</u>	3.6	3.78	23.1	5.92	7.9	5.97	14.4	4.14	2.5	1.65	11.4	4.43
<u>Sphaerodorum gracilis</u>						0.1	0.32					
<u>Spio filicornis</u>			0.2	0.42							0.1	0.32
Spionidae indet.												
Syllidae indet.	0.6	1.07	0.1	0.32	0.4	0.70	0.1	0.32	0.3	0.32	0.48	
<u>Terebellides stroemii</u>									2.5	2.32	0.8	1.32
<u>Trichobranchus roseus</u>									1.1	0.99	0.3	0.48
<u>Abra alba</u>	1.5	1.65	4.8	2.25	11.0	11.18	3.8	2.86	1.2	1.55	5.1	3.96
<u>Abra nitida</u>	15.9	7.49	8.5	6.29	11.0	7.21	14.3	9.07	29.5	18.73	27.2	14.34
<u>Acanthocardia echinata</u>						0.1	0.32					
<u>Astarte montagui</u>						0.1	0.32					
<u>Corbula gibba</u>			0.2	0.42	0.3	0.48	0.4	0.70	0.2	0.42	0.7	0.82
<u>Ennucula tenuis</u>					0.1	0.32			0.1	0.32	0.1	0.32
<u>Hiatella arctica</u>											0.2	0.42
<u>Limatula sulcata</u>											0.5	1.58
<u>Macoma calcarea</u>	1.3	1.57	19.0	12.36	10.7	4.62	7.3	2.71	2.0	1.70	3.5	2.27
<u>Musculus niger</u>					0.1	0.32						
<u>Mya arenaria</u>						0.1	0.32					
<u>Mya</u> sp.	0.1	0.32	0.3	0.48			0.1	0.32				
<u>Mysella bidentata</u>			0.1	0.32	0.2	0.42	0.6	1.26	0.1	0.32		
<u>Spisula</u> sp.											0.1	0.32
<u>Thracia</u> sp.									0.1	0.32	0.4	0.52
<u>Thyasira</u> sp.	123.5	31.47	151.3	43.87	178.5	56.11	103.6	25.98	37.8	18.61	156.6	88.55
<u>Yoldiella</u> sp.					0.1	0.32			0.1	0.32	0.1	0.32
<u>Phoronis mülleri</u>	0.3	0.95										
<u>Onchnesoma steenstrupi</u>					0.4	0.52					0.1	0.32
<u>Phascolion strombi</u>			0.1	0.32	0.2	0.42			0.3	0.48		
Sipunculida indet.											0.1	0.32
Polycladida indet.					0.1	0.32					0.2	0.42
	λ	0.48		0.54		0.49		0.61		0.84		0.62
Diversity:	H	1.19		1.35		1.33		1.65		2.42		1.70
	d	3.62		4.79		5.11		5.51		6.85		6.48
Sampling efficiency:		0.991 < s < 1		0.990 < s < 1		0.994 < s < 1		0.987 < s < 1		0.983 < s < 1		0.993 < s < 1

Table 30. Station SK 2, 50 m

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	X	s	\bar{X}	s								
<u>Funiculina quadrangularis</u>	0.1	0.32										
<u>Stylatula elegans</u>			0.1	0.32			0.1	0.32			0.2	0.42
<u>Chaetoderma nitidulum</u>	0.6	0.97	0.2	0.42	0.9	0.88	0.5	0.53	2.1	1.45	0.6	0.84
Amphipoda indet.	1.5	2.32	1.3	0.82	0.3	0.67	1.2	1.03	1.4	1.51	4.8	2.53
<u>Calocharis macandreae</u>	0.1	0.32										
Cumacea indet.	0.9	1.29	1.0	1.05	4.3	4.19	0.3	0.67	2.5	2.01	2.8	3.61
<u>Geryon tridens</u>					0.1	0.32	0.1	0.32				
<u>Lunatia pallida</u>											0.1	0.32
<u>Philine quadrata</u>			0.4	0.70			0.1	0.32				
<u>Philine scabra</u>												
<u>Philine sp.</u>	0.1	0.32										
<u>Retusa umbilicata</u>			0.1	0.32								
Holothuroidea indet.									0.2	0.42		
Nemertini indet.			0.9	1.10	1.0	0.94	1.0	0.82	1.5	1.08	1.2	1.14
<u>Amphilepis norvegica</u>					0.2	0.42						
<u>Amphiura chiajei</u>			0.3	0.48	0.5	0.71	0.4	0.52	0.7	0.82	0.7	0.67
<u>Amphiura filiformis</u>	0.3	0.48			0.8	1.03	0.2	0.42			0.3	0.48
<u>Amphiura sp.</u>			0.1	0.32	0.1	0.32						
<u>Ophiura albida</u>	0.2	0.42					0.3	0.48			1.0	1.05
<u>Ophiura sarsi</u>			0.2	0.42	0.1	0.32	0.1	0.32			0.2	0.63
<u>Ophiura sp.</u>					0.5	0.97	0.1	0.32	0.3	0.67	0.7	0.95
Ampharetidae indet.	0.1	0.32										
<u>Antinoella sarsi</u>	0.3	0.48										
<u>Aphrodita aculeata</u>					0.1	0.32						
<u>Asychis biceps</u>	0.6	1.07	0.7	1.25			0.2	0.42			0.1	0.32
<u>Chaetozone setosa</u>	0.2	0.42	0.5	0.71					0.5	0.71	0.3	0.67
<u>Dasybranchus caducus</u>	0.2	0.42										
<u>Diplocirrus glaucus</u>	3.9	2.02	5.0	4.14	2.2	1.14	1.9	1.60	13.0	4.32	14.4	6.96
<u>Driloneris filum</u>				0.1	0.32		0.3	0.48	0.6	0.70	0.1	0.32
Euclymeninae indet.									1.2	1.14		
<u>Glycera alba</u>	0.1	0.32			0.3	0.48						
<u>Glycera rouxii</u>	0.9	0.88	0.2	0.42	0.1	0.32	0.4	0.70	0.3	0.48		
<u>Goniada maculata</u>	0.3	0.48	0.1	0.32	0.5	0.53	0.4	0.52	0.3	0.48	0.2	0.63
<u>Heteromastus filiformis</u>	0.5	0.71	0.5	0.85					4.8	3.29	3.0	2.40
<u>Laonice cirrata</u>	0.2	0.42	0.1	0.32	0.1	0.32					0.3	0.67
<u>Leiochone borealis</u>											0.4	0.70
<u>Lumbrineris fragilis</u>	1.1	1.45	0.1	0.32	0.2	0.63	0.2	0.42	0.5	0.71		
<u>Lumbriclymene minor</u>	0.1	0.32							0.2	0.42		
Maldanidae indet.					0.2	0.42			0.2	0.42		
<u>Maldane sarsi</u>	0.3	0.67	0.1	0.32	0.1	0.32			0.2	0.63	5.2	7.45
<u>Melinna cristata</u>	0.1	0.32					0.6	0.84				
<u>Myriochele sp.</u>	1.5	1.58	1.2	1.23	1.8	1.87	1.1	0.88	1.4	1.51	0.8	0.63
<u>Nephrys ciliata</u>	0.7	0.95	0.9	0.88	0.8	1.32	0.1	0.32				
<u>Nephrys incisa</u>									0.1	0.32		
<u>Nephrys paradoxa</u>	0.2	0.63	0.3	0.67	0.5	0.53	0.6	0.97	0.3	0.48	0.6	0.84
<u>Notomastus latericeus</u>							0.2	0.63				

Table 30. cont.

	1972				1973				1974			
	JUNE		OCT.		JUNE		OCT.		JUNE		OCT.	
	\bar{X}	s	\bar{X}	s	\bar{X}	s	\bar{X}	s	\bar{X}	s	\bar{X}	s
<u>Ophelina norvegica</u>							0.1	0.32				
<u>Ophiodromus flexuosus</u>	0.4	0.70	0.2	0.42	0.4	0.52	0.3	0.48	0.8	0.79	0.1	0.32
<u>Owenia fusiformis</u>						0.3	0.67			0.2	0.42	
<u>Paramphinome jeffreysi</u>	2.6	2.50	5.6	4.84	1.9	1.37	1.4	0.97	3.1	2.60	2.6	3.20
<u>Pectinaria koreni</u>						0.1	0.32					
<u>Pholoe minuta</u>	0.6	1.58	0.2	0.63	0.3	0.67	0.1	0.32	0.2	0.63	0.5	0.71
<u>Phylo norvegicus</u>					0.3	0.48	0.3	0.67	0.1	0.32	0.2	0.42
<u>Polycirrus</u> sp.	0.1	0.32										
Polynoidae indet.			0.4	0.70			0.4	0.52			0.5	0.85
<u>Praxillella gracilis</u>	0.1	0.32	0.3	0.48	0.8	0.79	0.2	0.42			0.4	0.52
<u>Praxillella praetermissa</u>	2.8	2.53	2.9	1.52	0.5	0.85	1.6	0.97			1.1	0.99
<u>Sabella penicillatus</u>	0.1	0.32										
<u>Samytha sexcirtata</u>			0.1	0.32								
<u>Scoloplos armiger</u>	0.3	0.48									0.1	0.32
Spionidae indet.												
<u>Spiophanes krøyeri</u>	0.3	0.48										
Syllidae indet.	0.1	0.32							0.6	1.07		
<u>Terebellides stroemi</u>	0.1	0.32	0.5	0.71	0.3	0.48	0.2	0.63	0.1	0.32	0.9	1.10
<u>Trichobranchus roseus</u>	0.2	0.42			0.2	0.42	0.1	0.32	0.3	0.67	0.1	0.32
<u>Abra alba</u>	7.9	7.48	12.2	3.05	10.6	4.40	20.3	10.46	6.5	2.84	13.4	5.52
<u>Abra nitida</u>	12.9	7.87	19.3	6.99	13.1	4.18	10.1	6.98	36.6	12.41	29.6	14.37
<u>Arctica islandica</u>				0.1	0.32					0.1	0.32	
Cardiidae indet.	0.1	0.32										
<u>Corbula gibba</u>					0.1	0.32						
<u>Ennucula tenuis</u>	0.1	0.32					0.2	0.42				
<u>Kelliella miliaris</u>			0.2	0.42	0.2	0.63						
<u>Limatula</u> sp.	0.2	0.63									0.1	0.32
<u>Limatula sulcata</u>			0.1	0.32			0.7	0.82	0.9	0.88	3.5	2.46
<u>Modiolus modiolus</u>			0.1	0.32								
<u>Mytilus edulis</u>				0.1	0.32							
<u>Parvicardium minimum</u>			0.4	0.70	0.2	0.42	0.5	0.71	0.6	1.07	2.0	1.94
<u>Thyasira</u> sp.	14.1	8.12	30.3	12.04	21.6	7.07	15.5	4.65	8.0	3.77	6.8	4.42
<u>Tropidomya abbreviata</u>											0.2	0.42
<u>Yoldiella</u> sp.	1.2	1.03	4.0	2.91	1.3	1.16	1.3	1.64	1.4	0.52	1.9	1.37
<u>Phoronis mülleri</u>			0.1	0.32								
<u>Onchnesoma steenstrupi</u>			0.4	0.97								
<u>Phascolion strombi</u>	0.2	0.42			0.3	0.48	0.2	0.42				
Polycladida indet.							0.1	0.32	0.1	0.32		
λ	0.87		0.82		0.83		0.82		0.80		0.87	
Diversity:	H	2.59		2.24		2.36		2.31		2.27		2.57
	d	7.04		5.86		6.13		6.65		4.84		5.92
Sampling efficiency:	0.964 < s < 1	0.979 < s < 1	0.975 < s < 1	0.972 < s < 1	0.990 < s < 1	0.985 < s < 1						

Table 31. Station VB 1, 20 m

	1974	
	JUNE	
	\bar{x}	s
<u>Funiculina quadrangularis</u>	0.1	0.32
<u>Ctenodiscus crispatus</u>	0.3	0.48
<u>Chaetoderma nitidulum</u>	0.4	0.52
Amphipoda indet.	0.3	0.95
Cumacea indet.	1.3	1.25
<u>Brissopsis lyrifera</u>	0.1	0.32
<u>Oenopota cancellata</u>	0.1	0.32
Holothuroidea indet.	0.1	0.32
Nemertini indet.	1.1	0.74
<u>Amphiura chiajei</u>	0.1	0.32
<u>Amphiura filiformis</u>	0.1	0.32
<u>Amphiura</u> sp.	0.1	0.32
<u>Ophiura</u> sp.	0.1	0.32
Ampharetidae indet.	0.1	0.32
<u>Asychis biceps</u>	0.9	1.37
<u>Chaetozone setosa</u>	1.0	1.15
Cirratulidae indet.	0.1	0.32
<u>Diplocirrus glaucus</u>	4.8	3.36
Euclymninae indet.	0.3	0.48
<u>Glycera alba</u>	0.1	0.32
<u>Goniada maculata</u>	0.6	0.70
<u>Heteromastus filiformis</u>	1.4	1.07
Maldanidae indet.	0.1	0.32
<u>Maldane sarsi</u>	1.2	1.14
<u>Myriochele</u> sp.	4.8	3.94
<u>Nephtys incisa</u>	0.1	0.32
Ophiodromus flexuosus	0.7	0.82
<u>Owenia fusiformis</u>	15.4	22.11
<u>Paramphinome jeffreysi</u>	0.1	0.32
<u>Scoloplos armiger</u>	0.1	0.32
<u>Spiochaetopterus typicus</u>	0.1	0.32
Spionidae indet.	0.1	0.32
Syllidae indet.	0.1	0.32
<u>Terebellides stroemi</u>	1.7	0.95
<u>Thelepus cincinnatus</u>	0.1	0.32
<u>Trichobranchus roseus</u>	0.1	0.32
<u>Abra nitida</u>	4.3	2.45
<u>Astarte sulcata</u>	0.1	0.32
<u>Parvicardium minimum</u>	0.1	0.32
<u>Teredo</u> sp.	0.2	0.63
<u>Thyasira</u> sp.	4.2	2.15
<u>Yoldiella</u> sp.	0.2	0.42
<u>Phoronis mülleri</u>	0.1	0.32
<u>Onchnesoma steenstrupi</u>	0.2	0.42
<u>Phascolion strombi</u>	0.2	0.42
	λ	0.86
Diversity:	H	2.55
	d	7.13
Sampling efficiency:	0.932 < S < 1	

Table 32. Station VB 2, 50 m

	1974			
	JUNE		OCT.	
	\bar{x}	s	\bar{x}	s
<u>Virgularia mirabilis</u>	0.1	0.32		
<u>Ctenodiscus crispatus</u>	0.1	0.32	0.4	0.52
<u>Chaetoderma nitidulum</u>	0.1	0.32	0.2	0.42
Amphipoda indet.	2.2	2.44	3.7	2.83
<u>Calocharis macandreae</u>			0.1	0.32
Cumacea indet.	0.2	0.42	0.5	0.71
<u>Brisaster fragilis</u>	0.1	0.32		
<u>Briassopsis lyrifera</u>	0.4	0.52		
Nemertini indet.	0.7	1.06	0.9	0.74
<u>Amphilepis norvegica</u>	0.3	0.67		
<u>Amphiura chiajei</u>	0.1	0.32		
<u>Amphiura filiformis</u>			0.1	0.32
<u>Asychis biceps</u>	0.3	0.48	0.1	0.32
<u>Chaetozone setosa</u>	2.3	1.70	10.6	6.40
<u>Diplocirrus glaucus</u>	1.2	1.14	0.3	0.48
<u>Driloneris filum</u>	0.1	0.32		
Euclymeninae indet.			0.1	0.32
<u>Heteromastus filiformis</u>	2.2	1.62	8.7	4.27
<u>Laonice cirrata</u>	0.2	0.42		
<u>Lumbineris fragilis</u>	0.4	0.52		
Maldanidae indet.	0.1	0.32		
<u>Maldane sarsi</u>	1.1	1.60	0.1	0.32
<u>Myriochele</u> sp.	1.1	1.91	0.5	0.97
<u>Nephtys paradoxa</u>			0.1	0.32
<u>Ophiodromus flexuosus</u>	0.2	0.42	0.1	0.32
<u>Owenia fusiformis</u>	0.1	0.32		
<u>Panthalis oerstedi</u>	0.1	0.32		
<u>Paramphinome jeffreysi</u>	0.1	0.32	0.4	0.70
<u>Pectinaria auricoma</u>			0.2	0.42
<u>Pectinaria belgica</u>	0.1	0.32		
<u>Pectinaria koreni</u>			0.2	0.42
<u>Phylo norvegicus</u>	0.1	0.32	1.6	1.07
<u>Praxillella gracilis</u>			0.4	0.70
<u>Praxillella praetermissa</u>			0.1	0.32
<u>Rhodine gracilior</u>	0.1	0.32		
<u>Scoloplos armiger</u>			0.1	0.32
<u>Streblosoma hairdi</u>	0.1	0.32		
<u>Terebellides stroemi</u>	1.2	1.48	0.1	0.32
<u>Abra alba</u>	0.2	0.42		
<u>Abra nitida</u>	1.8	1.32	0.5	0.71
<u>Thyasira</u> sp.	0.6	0.70	2.4	1.90
<u>Phascolion strombi</u>	0.1	0.32		
	λ	0.93		0.80
Diversity:	κ	2.88		2.08
	ζ	6.16		4.32
Sampling efficiency:	0.874	$< S < 1$	0.948	$< S < 1$

Table 33.

Qualitative similarity (cc)

BV1	BV2	BV3	FT1	FT2	FT3	KV2	KV3	KV4	OF1	OF2	OF3	OF4	SD1	SD2	SF1	SF2	SK1	SK2	TH1	TH2	TH3	VB1	VB2	VD1	VD2	YØ1	AF1	AF2	AF3
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

BV1	54	34	42	38	30	28	21	19	26	27	29	17	42	42	18	30	40	35	35	15	13	26	21	37	28	31	33	28	32
BV2	.53	40	37	36	30	34	25	26	31	32	35	24	34	45	23	33	37	37	30	18	14	33	33	40	36	34	37	34	33
BV3	.45	.49	32	34	32	32	35	32	26	24	28	25	33	37	27	35	29	42	29	26	19	35	34	35	36	35	40	28	26
FT1	.33	.31	.24	49	38	34	25	23	24	20	23	13	52	47	21	31	51	37	43	16	16	27	28	50	35	41	28	34	34
FT2	.11	.17	.12	.40	44	40	30	27	20	20	23	14	46	44	26	38	47	44	44	15	17	30	31	47	37	44	35	31	28
FT3	.21	.27	.31	.50	.29	45	40	35	20	14	18	13	40	39	32	45	39	45	50	16	18	28	28	35	39	47	34	27	27
KV2	.21	.21	.32	.21	.10	.34	36	30	21	18	19	15	30	31	26	40	38	48	43	17	14	29	27	36	40	49	27	28	27
KV3	.09	.10	.12	.07	.04	.28	.23	48	19	15	20	17	22	24	44	49	25	42	31	24	18	30	26	26	30	38	28	19	22
KV4	.11	.12	.15	.11	.05	.23	.30	.62	22	17	20	17	23	24	45	44	26	37	27	30	21	31	32	25	32	32	29	19	22
OF1	.17	.08	.08	.09	.03	.04	.05	.05	.06	41	35	29	24	30	19	20	25	22	20	16	11	22	23	23	28	22	19	24	26
OF2	.14	.07	.08	.08	.04	.05	.04	.05	.81	48	28	19	22	16	18	24	18	18	15	13	18	19	21	22	18	20	27	30	
OF3	.48	.42	.45	.23	.10	.21	.18	.05	.09	.45	.42	27	20	25	19	20	25	20	19	17	11	19	20	23	21	19	19	25	27
OF4	.03	.03	.08	.01	.00	.02	.05	.05	.05	.03	.05	.08	13	17	18	16	16	18	12	23	11	25	22	17	21	15	20	21	24
SD1	.37	.36	.24	.40	.19	.22	.24	.10	.18	.07	.06	.14	.02	48	22	31	47	36	40	16	14	25	25	43	36	36	33	33	31
SD2	.24	.34	.21	.43	.32	.24	.21	.08	.13	.05	.05	.13	.01	.61	21	32	47	39	40	16	14	32	34	49	43	37	42	38	38
SF1	.09	.07	.11	.05	.02	.15	.16	.37	.40	.05	.04	.05	.08	.07	.04	41	22	36	25	32	19	24	28	24	27	31	22	17	22
SF2	.24	.22	.24	.16	.07	.36	.37	.68	.60	.05	.04	.16	.04	.26	.17	.30	32	50	40	24	22	32	30	30	31	42	33	22	26
SK1	.22	.25	.17	.38	.58	.28	.14	.05	.08	.13	.11	.24	.01	.24	.24	.03	.11	40	45	15	13	33	28	50	47	40	38	39	40
Sk2	.32	.38	.38	.53	.30	.55	.34	.16	.21	.05	.06	.31	.02	.41	.43	.07	.26	.34	43	22	19	40	35	39	42	51	38	33	29
TH1	.35	.35	.46	.30	.16	.35	.37	.14	.16	.09	.06	.31	.02	.32	.24	.08	.25	.23	.44	16	16	26	23	39	36	46	32	30	31
TH2	.03	.02	.06	.01	.00	.02	.03	.09	.11	.02	.01	.02	.17	.02	.01	.16	.08	.01	.02	.03	29	21	29	20	20	18	18	13	11
TH3	.03	.02	.21	.01	.01	.10	.16	.05	.04	.02	.01	.02	.05	.02	.01	.07	.04	.01	.02	.24	.05	21	20	17	17	18	19	11	09
VB1	.09	.10	.14	.06	.04	.06	.14	.20	.22	.05	.05	.07	.13	.11	.09	.13	.17	.04	.12	.11	.08	.07	38	33	31	30	39	34	26
VB2	.07	.09	.15	.05	.03	.10	.16	.17	.23	.08	.12	.19	.29	.07	.06	.20	.17	.03	.13	.10	.10	.06	.36	31	43	28	33	30	23
VD1	.23	.18	.14	.36	.11	.14	.17	.13	.18	.06	.05	.06	.03	.47	.29	.10	.24	.14	.34	.23	.04	.03	.13	.12	44	39	36	42	39
VD2	.15	.25	.12	.31	.17	.16	.19	.11	.17	.06	.06	.07	.02	.58	.53	.06	.19	.14	.37	.19	.02	.02	.11	.15	.43	38	37	36	36
YØ1	.32	.25	.29	.29	.14	.33	.40	.33	.39	.07	.07	.21	.07	.48	.32	.18	.50	.16	.46	.41	.04	.07	.14	.18	.47	.39	34	28	26
AF1	.15	.14	.16	.10	.04	.09	.13	.15	.18	.05	.04	.06	.05	.30	.21	.07	.19	.10	.15	.15	.06	.05	.36	.21	.17	.12	.18	46	31
AF2	.16	.31	.13	.08	.05	.04	.08	.12	.13	.04	.04	.06	.07	.23	.16	.07	.11	.05	.08	.07	.05	.05	.24	.17	.18	.23	.08	.24	39
AF3	.18	.18	.21	.09	.05	.10	.21	.14	.18	.08	.15	.26	.17	.15	.11	.09	.22	.06	.16	.14	.04	.04	.29	.44	.16	.12	.20	.22	.25

Quantitative similarity (C_z)

LEGENDS TO THE FIGURES:

Figs. 2-31. Column diagrams showing number of individuals per m^2 (left scale) of the numerically dominant species, and plots of the values of the diversity indices (right scale).

Fig. 2. OF 1




Fig. 3. OF 2




Fig. 4. OF 3




Fig. 5. OF 4




Fig. 6.




Fig. 7.




Fig. 8.




Fig. 9.





Fig. 10.



TH 3




Fig. 11.

Melitaea cynthia

Fig. 12.

SF 1




Fig. 13.

SF 2




Fig. 14.

SD 1




Fig. 15. SD 2




Fig. 16.

ÅF 1




Fig. 17.

ÅF 2




Fig. 18.

ÅF 3




Fig. 19.

FT 1




Fig. 20. FT 2




Fig. 21. 2000





Fig. 22.



VD 2

Fig. 23.

VD 1




Fig. 24.




Fig. 25.

KV 2




Fig. 26.

KV 3




Fig. 27.

KV 4




Fig. 28.

VB 1




Fig. 29.

VB 2




Fig. 30.





Fig. 31. SK 2



Received April 1977
Printed August 1977